

ISBN: 978-81-69967-10-5

Emerging Paradigms in Science and Technology Research: Trends and Impact



— Editors —

Prof. Yogita Shinde
Dr. Jagendra D. Punde
Mr. Mujahid M. Hussain
Dr. R. Elavarasi



Bhumi Publishing, India
First Edition: September 2026

Emerging Paradigms in Science and Technology Research:

Trends and Impact

(ISBN: 978-81-69967-10-5)

DOI: <https://doi.org/10.5281/zenodo.23051294>

Editors

Prof. Yogita Shinde

Department of Pharmaceutics,
Vishwakarma University,
Pune, M.S.

Dr. Jagendra D. Punde

Department of Physics,
S. S. Girls' College,
Gondia, M.S.

Mr. Mujahid M. Hussain

Department of Physics,
Maharashtra College of Arts, Science &
Commerce, Nagpada, Mumbai

Dr. R. Elavarasi

Department of Electrical and Electronics
Engineering, AMET Deemed to be
University, Chennai



Bhumi Publishing

September 2026

Copyright © Editors

Title: Emerging Paradigms in Science and Technology Research: Trends and Impact

Editors: Prof. Yogita Shinde, Dr. Jagendra D. Punde, Mr. Mujahid M. Hussain, Dr. R. Elavarasi

First Edition: September 2026

ISBN: 978-81-69967-10-5



DOI: <https://doi.org/10.5281/zenodo.23051294>

All rights reserved. No part of this publication may be reproduced or transmitted, in any form or by any means, without permission. Any person who does any unauthorized act in relation to this publication may be liable to criminal prosecution and civil claims for damages.

Published by Bhumi Publishing,

a publishing unit of Bhumi Gramin Vikas Sanstha



Nigave Khalasa, Tal – Karveer, Dist – Kolhapur, Maharashtra, INDIA 416 207

E-mail: bhumipublishing@gmail.com



Disclaimer: The views expressed in the book are of the authors and not necessarily of the publisher and editors. Authors themselves are responsible for any kind of plagiarism found in their chapters and any related issues found with the book.

PREFACE

Science and technology are advancing at an unprecedented pace, transforming the ways knowledge is generated, applied, and shared across disciplines and societies. *Emerging Paradigms in Science and Technology Research: Trends and Impact* presents a contemporary perspective on these transformations, bringing together diverse research contributions that reflect current developments, innovative approaches, and emerging directions across scientific and technological domains.

The book highlights the growing importance of interdisciplinary research in addressing complex challenges that cannot be effectively resolved within conventional disciplinary boundaries. Advances in artificial intelligence, machine learning, data science, biotechnology, nanotechnology, environmental sciences, materials science, information technology, and related fields are creating new opportunities for discovery and application. At the same time, these developments raise important questions concerning sustainability, ethics, accessibility, societal impact, and responsible innovation.

The chapters included in this volume explore emerging trends, research methodologies, technological applications, and contemporary challenges from different perspectives. They demonstrate how scientific knowledge and technological innovation can contribute to improved healthcare, sustainable development, environmental protection, efficient resource utilization, digital transformation, and enhanced quality of life. The contributions also emphasize the significance of collaboration, evidence-based research, technological adaptability, and continuous learning in a rapidly changing research ecosystem.

This book is intended for researchers, academicians, teachers, students, professionals, and readers interested in understanding contemporary developments in science and technology. By bringing together multidisciplinary perspectives, it seeks to encourage dialogue, stimulate further investigation, and promote innovative thinking.

The editors hope that this volume will serve as a useful academic resource and inspire researchers to explore new frontiers of knowledge. It is envisioned not merely as a compilation of research contributions, but as a platform for understanding emerging paradigms and their broader implications for science, technology, society, and sustainable development.

- Editors

TABLE OF CONTENT

Sr. No.	Book Chapter and Author(s)	Page No.
1.	ANTI-INFLAMMATORY FOODS IN METABOLIC DISORDERS: INSULIN RESISTANCE, OBESITY, AND GLYCAEMIC CONTROL Nivetha Muralikrishnaa and John Wyson	1 – 8
2.	CLIMATE-SMART EXTENSION: GRASSROOTS STRATEGIES FOR EL NIÑO MITIGATION AND MOISTURE STRESS MANAGEMENT I. Venkata Reddy, M. Ravi Kishore, P. N. Siva Prasad, R. Prabhavathi, N. Rajasekhar, K. Lakshmi Kala and T. Jeswanth Reddy	9 – 18
3.	GREEN NANOREMEDIATION: SUSTAINABLE ADVANCES IN ENVIRONMENTAL REMEDIATION MECHANISMS AND APPLICATIONS Rashmi Verma and Tanzeel Ahmed	19 – 39
4.	HUMAN MICROBIOME: EMERGING ROLES IN HEALTH AND DISEASE Poonguzhali K. and Arulmozhi R.	40 – 46
5.	HAIR AND SCALP DISEASES: MEDICAL, SURGICAL AND COSMETIC TREATMENT M. Kannahi and R. Sasireka	47 – 56
6.	NANOTECHNOLOGY FOR SUSTAINABLE DEVELOPMENT: EMERGING MATERIALS AND APPLICATIONS Shahnaz Kossar and Barjinder Kaur	57 – 66
7.	COMPARATIVE PERFORMANCE EVALUATION OF WIRED AND WIRELESS SMART HOME AUTOMATION USING A PYTHON-BASED VIRTUAL FRAMEWORK Konduri Ramesh	67 – 78
8.	ADVANCED MATERIALS AND GREEN CHEMISTRY: EMERGING TRENDS IN SUSTAINABLE SCIENCE AND TECHNOLOGY Savitha H S	79 – 86

9.	FUZZY COGNITIVE MODELLING OF INNOVATIVE PEDAGOGIES FOR ENHANCING COMMUNICATIVE SKILLS AMONG RURAL STUDENTS AND PROMOTING INCLUSIVITY	87 – 100
	Nivetha Martin and A Sahaya Josephine Mary	
10.	CLIMATE CHANGE AND ENVIRONMENTAL SUSTAINABILITY	101 – 108
	D. Ashok Kumar and D. Mahhamitra	
11.	BIOREMEDIATION: MICROBIAL SOLUTIONS FOR ENVIRONMENTAL SUSTAINABILITY	109 – 117
	Arulmozhi R. and Sudharani P.	
12.	SHIFTING PARADIGMS IN SCIENCE AND TECHNOLOGY: FROM DATA-DRIVEN DISCOVERY TO CONVERGENT SOLUTIONS	118 – 131
	Umarani	
13.	WSe₂ THIN FILMS: STRUCTURAL AND OPTICAL PROPERTIES	132 – 140
	Prashant A. Chate and Dattatray J. Sathe	

ANTI-INFLAMMATORY FOODS IN METABOLIC DISORDERS: INSULIN RESISTANCE, OBESITY, AND GLYCAEMIC CONTROL

Nivetha Muralikrishnaa* and John Wyson

Department of Food Processing Technology,

AMET University, Chennai, Tamil Nadu, India

*Corresponding author E-mail: nivethakrishnaa20@gmail.com

Abstract

Metabolic disorders, including obesity, insulin resistance, metabolic syndrome and type 2 diabetes, are increasingly recognised as conditions driven partly by chronic low-grade inflammation. Dietary strategies that simultaneously regulate inflammatory and metabolic pathways may therefore improve disease prevention and management. Anti-inflammatory foods—including vegetables, fruits, whole grains, legumes, nuts, seeds, fatty fish and extra-virgin olive oil—provide fibre, polyphenols, carotenoids, unsaturated fatty acids and essential micronutrients. This chapter examines the mechanisms through which these foods influence insulin sensitivity, adipose-tissue function, appetite, intestinal integrity and glycaemic regulation. Polyphenols may inhibit nuclear factor- κ B and NLRP3 inflammasome signalling while activating antioxidant and energy-sensing pathways. Omega-3 fatty acids contribute to specialised pro-resolving mediators, whereas fermentable fibre supports microbial production of short-chain fatty acids that regulate intestinal permeability, inflammation and glucose metabolism. Anti-inflammatory dietary patterns may also lower dietary energy density, promote satiety and reduce visceral adiposity. Clinical evidence suggests that Mediterranean-style and low-glycaemic dietary patterns produce modest improvements in body weight, fasting glucose, glycated haemoglobin and insulin resistance. However, responses vary with baseline metabolic phenotype, food processing, bioactive availability, energy balance and microbiome composition. Anti-inflammatory foods should consequently replace refined carbohydrates, processed meats and foods rich in saturated fat rather than merely being added to existing diets. Although they cannot replace pharmacological therapy or structured weight management, varied and minimally processed anti-inflammatory foods provide a practical framework for restoring metabolic flexibility and supporting durable glycaemic control.

Keywords: Anti-Inflammatory Foods, Insulin Resistance, Obesity, Glycaemic Control, Gut Microbiota, Metabolic Syndrome.

1. Introduction

Insulin resistance, obesity and impaired glycaemic control are interconnected metabolic disturbances characterised by defective insulin signalling, abnormal adipose-tissue function and

persistent low-grade inflammation. Enlarged adipocytes, excess circulating nutrients and intestinal-barrier dysfunction stimulate inflammatory pathways that interfere with glucose uptake and promote progressive metabolic deterioration. Diet can either reinforce or attenuate this inflammatory environment.

Anti-inflammatory foods supply fibre, unsaturated fatty acids, polyphenols, carotenoids, vitamins and minerals capable of influencing immune and metabolic signalling. Vegetables, fruits, legumes, whole grains, nuts, seeds, fatty fish and extra-virgin olive oil form the principal components of this dietary approach. Their combined effects may improve satiety, microbial fermentation, insulin sensitivity and postprandial glucose regulation. However, no food acts independently of energy intake, processing or overall dietary quality. This chapter evaluates the biological mechanisms and clinical evidence connecting anti-inflammatory foods with obesity management, improved insulin action and glycaemic control, while identifying priorities for personalised and culturally appropriate metabolic nutrition.

2. Metabolic Inflammation and Insulin Resistance

Insulin resistance occurs when skeletal muscle, liver and adipose tissue respond inadequately to insulin. Visceral adiposity contributes substantially by increasing the release of non-esterified fatty acids and inflammatory mediators. Macrophage infiltration into expanding adipose tissue increases tumour necrosis factor- α , interleukin-6 and related cytokines, which can disrupt insulin-receptor signalling and glucose transport.

Nuclear factor- κ B, c-Jun N-terminal kinase and the NLRP3 inflammasome connect nutrient excess with inflammatory activation. Saturated fatty acids, oxidative stress, mitochondrial dysfunction and microbial endotoxins may stimulate these pathways. In contrast, fibre-rich and antioxidant-containing foods can reduce inflammatory exposure and support metabolic homeostasis (van der Heiden & te Velde, 2026).

Dietary inflammatory potential is therefore clinically relevant. Higher Dietary Inflammatory Index values have been associated with increased waist circumference, body mass index, blood pressure and metabolic-syndrome prevalence (Zhang *et al.*, 2022). Pro-inflammatory dietary patterns may also aggravate fasting hyperglycaemia and hyperinsulinaemia. Anti-inflammatory nutrition should consequently be understood as modifying the metabolic environment that generates insulin resistance rather than merely lowering an isolated inflammatory biomarker.

3. Polyphenols and Cellular Metabolic Signalling

Berries, grapes, tea, cocoa, colourful vegetables, nuts, seeds and extra-virgin olive oil provide flavonoids, anthocyanins, phenolic acids, stilbenes and lignans. These polyphenols can regulate metabolic signalling despite generally low circulating concentrations. Their actions include modulation of nuclear factor- κ B, activation of AMP-activated protein kinase and enhancement of endogenous antioxidant defences.

AMP-activated protein kinase functions as a cellular energy sensor. Its activation can promote fatty-acid oxidation, glucose uptake and mitochondrial function while restricting hepatic lipid synthesis. Polyphenols may additionally preserve pancreatic β -cell function by reducing oxidative and inflammatory stress. Some compounds inhibit α -amylase, α -glucosidase and dipeptidyl peptidase-IV, potentially slowing carbohydrate digestion and moderating postprandial glucose responses (Song *et al.*, 2023).

Polyphenols also influence adipocyte differentiation and inflammatory cytokine production. Anthocyanin-rich foods may improve insulin sensitivity by modulating adipose-tissue inflammation and glucose-transporter activity (Ruy *et al.*, 2025). Nevertheless, food-based effects depend on dose, chemical structure, processing and microbial transformation. Polyphenol-rich whole foods provide fibre and micronutrients alongside phytochemicals and may therefore be more physiologically relevant than isolated high-dose supplements.

4. Fibre, Gut Microbiota and Metabolic Endotoxaemia

Whole grains, pulses, vegetables, fruits, nuts and seeds deliver resistant starch, β -glucan, arabinoxylans and other fermentable carbohydrates. These substrates reach the colon, where microorganisms convert them into acetate, propionate and butyrate. Short-chain fatty acids support intestinal-barrier integrity and interact with receptors involved in appetite, immune activity and glucose regulation.

Obesity and insulin resistance are frequently accompanied by altered microbial function and increased intestinal permeability. The resulting transfer of lipopolysaccharide into circulation may promote metabolic endotoxaemia and interfere with insulin signalling. Fibre-rich diets may restrict this process by supporting mucus production, epithelial integrity and beneficial microbial metabolism (Liu *et al.*, 2025).

Polyphenols and the microbiota also interact bidirectionally. Microorganisms convert complex polyphenols into smaller metabolites, while polyphenols may selectively influence microbial populations. Recent intervention evidence suggests that polyphenol-rich treatments can reduce metabolic endotoxaemia, increase short-chain-fatty-acid production and support antioxidant activity in adults with overweight or obesity (González-Gómez *et al.*, 2025).

A higher dietary index for gut-microbiota support has been associated with lower metabolic-syndrome prevalence, with dietary inflammation partially mediating this relationship (Han *et al.*, 2025). These observations position the gut barrier and its metabolites as important links between anti-inflammatory foods and metabolic health.

5. Satiety, Adipose Tissue and Obesity Management

Anti-inflammatory foods are frequently fibre-rich, minimally processed and relatively low in energy density. Vegetables, fruits, pulses and intact whole grains increase food volume and gastric distension without proportionally increasing energy intake. Soluble viscous fibres can

delay gastric emptying and may enhance glucagon-like peptide-1 and peptide YY secretion, thereby extending fullness.

Nuts and seeds provide greater energy density but combine protein, fibre and unsaturated fats that can support satiety when consumed in appropriate portions. Fatty fish provides high-quality protein and omega-3 fatty acids, while extra-virgin olive oil can improve dietary fat quality when it replaces saturated fat. These foods do not automatically produce weight loss; substitution and portion control remain essential.

Mediterranean-style diets are particularly relevant because they integrate these foods into a coherent pattern. Greater adherence has been associated with lower insulin resistance among individuals with overweight or obesity, although fish intake and overall dietary quality may partly drive the association (Vetrani *et al.*, 2023). Energy-restricted Mediterranean interventions can also support clinically meaningful weight reduction and improvement of obesity-related metabolic abnormalities (Muscogiuri *et al.*, 2022).

Reduction of visceral adiposity is especially important because it may decrease inflammatory cytokine release and improve insulin action even when total weight loss remains moderate. Anti-inflammatory foods should therefore make an energy deficit more nutritious and sustainable rather than being promoted as independent fat-burning agents.

6. Glycaemic Control and Insulin Sensitivity

Anti-inflammatory dietary patterns improve glycaemic regulation through slower carbohydrate digestion, greater insulin sensitivity, reduced hepatic glucose output and lower inflammatory stress. Low-glycaemic foods—including pulses, minimally processed whole grains and many fruits and vegetables—reduce rapid glucose entry into circulation. Recent meta-analysis indicates that low-glycaemic diets can improve insulin resistance even among adults without diagnosed diabetes (Yu *et al.*, 2025).

Whole-grain intake has been associated with improvements in postprandial glycaemia, fasting glucose and glycated haemoglobin, although effects vary across trials (Li *et al.*, 2023; Ying *et al.*, 2024). Oat and barley β -glucan increase intestinal viscosity, while pulses provide slowly digestible starch and plant protein. Nuts and unsaturated oils may further reduce meal-level glycaemic responses when replacing refined carbohydrates or saturated fats.

Among patients with type 2 diabetes, Mediterranean dietary patterns have produced reductions in glycated haemoglobin, fasting glucose, body mass index and waist circumference compared with control diets (Zheng *et al.*, 2024). Comparative evidence has also ranked the Mediterranean diet highly for glycaemic regulation, although moderate-carbohydrate and low-glycaemic strategies may provide comparable benefits for particular individuals (Yuan *et al.*, 2024).

The optimal diet is therefore not defined by one macronutrient ratio. It is characterised by high-quality carbohydrates, adequate fibre, minimally processed foods and appropriate energy intake.

7. Anti Inflammatory Dietary Patterns in Clinical Management

The Mediterranean diet, DASH diet, Nordic diet and well-designed plant-forward patterns share several anti-inflammatory characteristics. They emphasise vegetables, fruits, whole grains, legumes, nuts and unsaturated fats while restricting processed meat, refined carbohydrates, excess sodium and added sugars. Their benefits result from cumulative food substitution and biological synergy.

Recent reviews identify the Mediterranean diet as the most consistently supported nutritional approach for insulin resistance and type 2 diabetes prevention (Caretto *et al.*, 2026). However, adherence may be limited when recommendations ignore cultural preferences, food cost, cooking skills or availability. Effective translation requires regionally appropriate pulses, grains, vegetables, seeds, spices and unsaturated oils rather than rigid replication of European foods.

Food selection should also account for processing. Sweetened yoghurt, refined cereal products and heavily salted nuts should not be classified as anti-inflammatory based only on one desirable ingredient. The complete nutritional profile and the food being replaced determine metabolic value.

8. Challenges and Future Perspectives

The field must move beyond universal “anti-inflammatory food” lists towards metabolic precision nutrition. Current trials vary in dietary definitions, energy restriction, intervention duration and biomarkers. Changes in C-reactive protein alone cannot establish improvements in adipose, hepatic or intestinal inflammation (Makiel *et al.*, 2026).

Future studies should combine continuous glucose monitoring, hyperinsulinaemic–euglycaemic clamps, appetite hormones, body-composition imaging and multi-omics profiling. Microbiome metabolites, intestinal permeability and adipose-tissue inflammatory signatures could reveal why responses differ among individuals. Polyphenol exposure should be characterised by food source, dose and bioavailability rather than reported as total intake.

Promising personalised approaches could match dietary patterns to insulin-resistance phenotype, microbial functionality, genetic susceptibility and cultural preference. Fermentation, germination and minimally disruptive processing may enhance bioactive accessibility while preserving fibre functionality. Digital monitoring could identify real-time glycaemic responses and guide food substitution. However, technological sophistication must not compromise affordability. The next generation of anti-inflammatory nutrition should unite precision with accessibility—transforming familiar regional foods into culturally acceptable dietary combinations that measurably reduce inflammation, restore insulin sensitivity and support sustainable weight control.

Conclusion

Anti-inflammatory foods provide a coherent nutritional framework for addressing the interconnected abnormalities underlying metabolic disorders. Vegetables, fruits, whole grains,

legumes, nuts, seeds, fatty fish and extra-virgin olive oil supply fibre, polyphenols, unsaturated fatty acids, vitamins and minerals capable of influencing inflammatory signalling, insulin action, appetite regulation and intestinal function. Their combined effects may reduce metabolic stress more effectively than any isolated nutrient.

These foods act through complementary pathways. Polyphenols regulate cellular energy sensors and inflammatory transcription factors; omega-3 fatty acids support inflammatory resolution; fermentable fibres generate short-chain fatty acids; and minimally processed carbohydrates moderate postprandial glucose exposure. By supporting satiety and reducing dietary energy density, anti-inflammatory foods may also facilitate the loss of visceral fat that drives insulin resistance.

However, their effects depend on substitution and total energy balance. Adding nuts, olive oil or whole-grain products to an energy-surplus diet will not independently reverse obesity. The strongest strategy replaces refined carbohydrates, processed meat and saturated-fat-rich foods with minimally processed alternatives while integrating physical activity, sleep management and clinical treatment.

Anti-inflammatory dietary patterns can improve glycaemic control and metabolic flexibility, but they should not replace glucose-lowering medication or structured obesity care when these are required. Individual responses will vary according to metabolic phenotype, microbiota, medication and food preparation. Future precision approaches may help identify the most effective combinations for each individual. Metabolic inflammation is not switched off by one food; it is progressively reduced when every meal replaces inflammatory excess with fibre, polyphenols, unsaturated fats and metabolic balance.

References

1. Caretto, A., Bianchi, C., & Giorgino, F. (2026). Targeting insulin resistance through nutrition: Evidence-based dietary strategies and future perspectives. *Nutrients*, 18(7), Article 1119. <https://doi.org/10.3390/nu18071119>
2. Cui, L., Zhang, Y., Liu, J., Wang, H., & Chen, X. (2025). Comparative effects of anti-inflammatory dietary interventions on obesity, glycaemic control and inflammation: A systematic review and network meta-analysis. *Frontiers in Nutrition*, 12, Article 1572604.
3. Fernandez, M. L., Murillo, A. G., & Ojo, O. (2022). Dietary treatments to reduce insulin resistance and inflammation. *Medical Research Archives*, 10(3). <https://doi.org/10.18103/mra.v10i3.2768>
4. González-Gómez, Á., Domínguez-López, I., Montserrat-de la Paz, S., & Ortega, E. (2025). Effect of polyphenol-rich interventions on gut microbiota and metabolic endotoxaemia in adults with overweight or obesity: A systematic review and meta-analysis. *Nutrients*, 17(16), Article 2641.

5. Han, Y. N., Wang, Y. X., Xiong, C. Y., Li, L., & Mao, R. M. (2025). Association between the dietary index for gut microbiota and metabolic syndrome: The mediating role of the Dietary Inflammatory Index. *Frontiers in Nutrition*, 12, Article 1617287. <https://doi.org/10.3389/fnut.2025.1617287>
6. Li, S., Zong, A., An, R., Wang, H., Liu, L., Liu, J., Guo, X., Xu, Z., Wang, J., Li, D., Du, F., & Xu, T. (2023). Effects of whole-grain intake on glycaemic traits: A systematic review and meta-analysis of randomized controlled trials. *Critical Reviews in Food Science and Nutrition*, 63(20), 4351–4370. <https://doi.org/10.1080/10408398.2021.2001429>
7. Liu, J., Zhang, Y., Wang, X., Li, H., & Chen, Q. (2025). Gut microbiota and its metabolites regulate insulin resistance in type 2 diabetes mellitus: Mechanisms and therapeutic implications. *Frontiers in Microbiology*, 16, Article 1554189. <https://doi.org/10.3389/fmicb.2025.1554189>
8. Makiel, K., Suder, A., & Targosz, A. (2026). Anti-inflammatory diets in metabolic syndrome and obesity: From dietary patterns to multi-omics perspectives. *International Journal of Molecular Sciences*, 27(6), Article 2734. <https://doi.org/10.3390/ijms27062734>
9. Muscogiuri, G., Verde, L., Sulu, C., Katsiki, N., Hassapidou, M., Frias-Toral, E., Cucalón, G., Pazderska, A., Yumuk, V. D., Colao, A., & Barrea, L. (2022). Mediterranean diet and obesity-related disorders: What is the evidence? *Current Obesity Reports*, 11(4), 287–304. <https://doi.org/10.1007/s13679-022-00481-1>
10. Ruy, C. C., Marques, C., & Martel, F. (2025). Anthocyanin modulation of gut microbiota to reverse chronic inflammation associated with obesity and insulin resistance. *Nutrients*, 17(23), Article 3727. <https://doi.org/10.3390/nu17233727>
11. Song, H., Li, X., Zhang, Y., Chen, S., & Wang, J. (2023). Beneficial effects of food-derived polyphenols on type 2 diabetes mellitus through gut-microbiota and inflammatory regulation. *Food Science of Animal Products*, 1(4), 9240043. <https://doi.org/10.26599/FSAP.2023.9240043>
12. Vetrani, C., Costabile, G., Vitale, M., Giacco, R., & Rivellese, A. A. (2023). The Mediterranean diet: Effects on insulin resistance and secretion in individuals with overweight or obesity. *Nutrients*, 15(21), Article 4524. <https://doi.org/10.3390/nu15214524>
13. Xie, F., Zhang, T., Zhang, Y., & Wang, X. (2023). Natural polyphenol–gut microbiota interactions and effects on metabolic disorders. *Trends in Food Science & Technology*, 139, 104116. <https://doi.org/10.1016/j.tifs.2023.104116>
14. Ying, T., Zheng, J., Kan, J., Li, W., Xue, K., Du, J., Liu, Y., & He, G. (2024). Effects of whole grains on glycaemic control: A systematic review and dose–response meta-analysis of prospective cohort studies and randomized controlled trials. *Nutrition Journal*, 23, Article 47. <https://doi.org/10.1186/s12937-024-00952-2>

15. Yu, X., Zhang, J., Zhao, X., & Li, Y. (2024). Overview of anti-inflammatory diets and their promising effects on noncommunicable diseases. *British Journal of Nutrition*, *131*(10), 1707–1722.
16. Yu, Y. T., Wang, J., Li, S., Zhang, Y., & Chen, H. (2025). Effect of dietary glycaemic index on insulin resistance in adults without diabetes: A systematic review and meta-analysis. *Frontiers in Nutrition*, *12*, Article 1458353. <https://doi.org/10.3389/fnut.2025.1458353>
17. Yuan, Y., Chen, J., Li, H., Wang, Y., & Zhang, X. (2024). Comparative efficacy of dietary patterns for glycaemic control in type 2 diabetes: A systematic review and network meta-analysis. *Food & Function*, *15*, 5098–5114. <https://doi.org/10.1039/D4FO00337C>
18. Zafari, R., Khosravi, M., Shidfar, F., & Gholami, F. (2025). The effects of polyphenols on gut microbial metabolites and their implications for metabolic health: A systematic review. *Journal of Nutritional Science*, *14*, Article e62. <https://doi.org/10.1186/s12986-025-01022-y>
19. Zhang, X., Li, R., Chen, Y., Dai, Y., Chen, L., Qin, L., & Liu, H. (2022). Association between Dietary Inflammatory Index and metabolic syndrome in the general adult population. *Frontiers in Nutrition*, *9*, Article 954057. <https://doi.org/10.3389/fnut.2022.954057>
20. Zheng, X., Liu, Z., Zhang, Y., Wang, Y., & Li, B. (2024). The effects of the Mediterranean diet on cardiovascular risk factors, glycaemic control and weight loss in patients with type 2 diabetes: A meta-analysis. *BMC Nutrition*, *10*, Article 47. <https://doi.org/10.1186/s40795-024-00858-4>

CLIMATE-SMART EXTENSION: GRASSROOTS STRATEGIES FOR EL NIÑO MITIGATION AND MOISTURE STRESS MANAGEMENT

**I. Venkata Reddy*, M. Ravi Kishore, P. N. Siva Prasad,
R. Prabhavathi, N. Rajasekhar, K. Lakshmi Kala and T. Jeswanth Reddy**

Dr. K. I. Rao Krishi Vigyan Kendra,

Garikapadu, NTR District, Andhra Pradesh, India.

*Correspondence author E-mail: ivrextedu18@gmail.com

Abstract

El Niño-induced monsoon aberrations — delayed onset, prolonged dry spells, and terminal moisture stress — have become recurring threats to rainfed agriculture across India's semi-arid tracts. Conventional extension delivery, organised around fixed seasonal training calendars, is frequently too slow to respond to abiotic shocks that emerge and intensify within a matter of weeks. This chapter examines how public and community-based extension systems can be reoriented into rapid, campaign-mode delivery mechanisms capable of protecting crop stands during El Niño years. Drawing on documented field experience from dryland tracts of Andhra Pradesh and the technical framework of India's national dryland research system, it discusses three grassroots pillars of climate-smart extension: frontline demonstrations of climate-resilient technology modules, pre-monsoon dry sowing (PMDS) as a moisture-capture and risk-hedging technique, and the rapid substitution of long-duration crops with short-duration millets and pulses under contingency conditions. The chapter further outlines farmer-training approaches for *In-Situ* and ex-situ moisture conservation and proposes a field-mobilisation model that links seasonal early-warning information, seed and input logistics, and last-mile extension functionaries into a single rapid-response chain. It concludes with institutional and policy recommendations for scaling grassroots contingency response across vulnerable dryland districts.

Keywords: El Niño, Climate-Smart Extension, Pre-Monsoon Dry Sowing, Contingency Crops, Moisture Conservation, Frontline Demonstration, Dryland Agriculture.

1. Introduction

Nearly 51 per cent of India's net sown area remains dependent on the south-west monsoon, and a large share of this rainfed tract is concentrated in agro-ecologically fragile, drought-prone districts of peninsular and central India. In such regions, the timing, spread, and intensity of rainfall — rather than its seasonal total — determine whether a cropping season succeeds or fails. The El Niño–Southern Oscillation (ENSO) is among the most influential drivers of this intra-seasonal variability: warm El Niño phases are statistically associated with weaker, delayed,

or erratically distributed monsoon rainfall over large parts of India, translating directly into moisture stress at critical crop-growth stages.

For the farming community, an El Niño year rarely announces itself as a single dramatic event. It manifests instead as a delayed sowing window, a prolonged dry spell after germination, or a premature withdrawal of rain before grain filling — each of which demands a different, time-bound agronomic response. Extension systems built primarily to disseminate pre-decided seasonal packages of practice are structurally unsuited to this kind of shifting risk. What is required instead is a grassroots extension architecture that can be triggered rapidly, that pre-positions seed, knowledge, and demonstration capacity ahead of the season, and that allows frontline workers to pivot quickly from a ‘normal-season’ advisory to a ‘contingency’ advisory.

This chapter is organised around three such grassroots strategies that have been documented in Indian dryland extension practice: frontline demonstrations of climate-resilient technology modules, pre-monsoon dry sowing as an anticipatory, moisture-capturing sowing technique, and the rapid promotion of short-duration millets and pulses as substitute or contingency crops. It also discusses moisture-conservation training and proposes a field-level campaign model through which District Agriculture Contingency Plans can be translated into visible, time-critical action at the village level.

2. El Niño and Monsoon Variability: The Extension Challenge

El Niño refers to the periodic warming of sea-surface temperatures in the central and eastern equatorial Pacific Ocean, one phase of the broader ENSO cycle. Historically, several of India’s major drought years have coincided with El Niño phases, although the relationship is probabilistic rather than deterministic — the strength of the monsoon in any given El Niño year is also modulated by the Indian Ocean Dipole and other regional factors. For extension planning, the practical implication is twofold: first, seasonal forecasts issued ahead of the south-west monsoon by the India Meteorological Department can flag an elevated probability of below-normal or erratically distributed rainfall; second, once the season begins, dry-spell monitoring at the district and block level becomes essential to detect emerging moisture stress in near-real time.

The 2026 kharif season illustrated this dynamic at district scale: administrations in drought-prone tracts of Andhra Pradesh scaled up pre-monsoon dry sowing on tens of thousands of acres specifically in response to El Niño-linked rainfall deficits, treating a basket of millet, pulse, and oilseed varieties with bio-inputs and broadcasting them ahead of the rains so that germination itself was timed to actual moisture availability rather than a fixed calendar date (Kurnool district administration campaign, 2026). Such campaigns underline that the extension response to El Niño is not a single technology but a coordinated sequence of anticipatory actions — seed

procurement, farmer mobilisation, field demonstration, and monitoring — compressed into a narrow pre-season window.

3. Reorienting Extension Systems for Contingency Planning

India already possesses a technical backbone for contingency response in the form of District Agriculture Contingency Plans (DACPs), prepared by the ICAR-Central Research Institute for Dryland Agriculture (CRIDA) in collaboration with State Agricultural Universities and line departments for the majority of the country's rural districts. These plans lay out crop- and situation-specific coping strategies for delayed onset of monsoon, seasonal drought, dry spells, and other weather aberrations. However, the value of a contingency plan is realised only when it is converted into field action within days, not weeks, of a trigger event — which is where the extension system, and specifically the Krishi Vigyan Kendra (KVK) network, becomes the operational link between a technical document and a farmer's field.

Reorienting extension for this purpose involves several shifts in working style. First, extension calendars need a built-in contingency module that can be activated on short notice once seasonal forecasts or observed dry-spell data cross a threshold, rather than waiting for the next scheduled training cycle. Second, interface meetings between KVKs, State Departments of Agriculture, and line departments (horticulture, animal husbandry, and irrigation) need to be convened before the onset of the rainy season, so that contingency seed and input requirements are already quantified and pre-positioned. Third, front-line workers need simplified, locally adapted advisories — delivered through farmer groups, village-level camps, and mobile-based agro-advisory services — that translate the technical language of a contingency plan into a short list of doable actions for a specific week.

4. Frontline Demonstrations of Climate-Resilient Practices

Frontline demonstrations (FLDs), conducted directly on farmers' fields under the supervision of scientists rather than only through intermediary extension staff, remain one of the most effective tools for building farmer confidence in unfamiliar technology. Under the National Innovations in Climate Resilient Agriculture (NICRA) programme, coordinated by ICAR-CRIDA, technology demonstration components have been implemented in a large number of climatically vulnerable districts, covering modules such as drought- and heat-tolerant crop varieties, *In-Situ* moisture conservation structures, resource-conserving tillage, and community-based custom hiring of climate-resilient farm implements.

For El Niño mitigation specifically, FLDs are most effective when they demonstrate a small, replicable technology bundle rather than a single practice in isolation — for instance, a bundle combining a short-duration drought-tolerant variety, a seed-treatment protocol, and one *In-Situ* moisture-conservation measure on the same demonstration plot. This allows farmers visiting the plot to see the cumulative effect of the bundle under real dry-spell conditions, which is

considerably more persuasive than a technology presented on paper before the season begins. Extension functionaries should identify a small number of ‘lead farmers’ per village cluster to host such demonstrations, since peer-to-peer diffusion from a visibly successful neighbouring plot has repeatedly been shown to outperform formal training sessions in driving adoption in dryland settings.

5. Pre-Monsoon Dry Sowing (PMDS): Concept and Field Mobilisation

5.1 What PMDS Involves

Pre-monsoon dry sowing is an anticipatory sowing technique in which a diverse mix of seeds — typically millets, pulses, oilseeds, and vegetables — is broadcast into dry soil roughly forty-five to sixty days ahead of the expected onset of the monsoon (Institute for Development Studies Andhra Pradesh, 2023). The seeds remain dormant in the soil and germinate only once sufficient rainfall is received, so the technique effectively removes the farmer’s dependence on guessing a single correct sowing date. Under the Andhra Pradesh Community-Managed Natural Farming (APCNF) programme, seeds sown under PMDS are typically pre-treated with a bio-inoculant slurry — variously termed Beejamrutham or Jeevamrutham, prepared from cattle dung, cattle urine, and other locally available materials — and are often further pelletised with fine clay and wood ash to improve germination uniformity and early seedling vigour (Sreenivasa *et al.*, 2009; Institute for Development Studies Andhra Pradesh, 2023).

5.2 Why PMDS Suits El Niño Years

The technique offers three advantages that are particularly relevant to El Niño-linked rainfall uncertainty. First, because germination is triggered by rainfall itself rather than by a fixed calendar date, farmers avoid the common risk of sowing after a false start to the monsoon, only to lose the crop to a subsequent dry spell. Second, the practice of sowing a diverse poly-crop mixture — often ten or more species on a single plot — spreads risk across crops with different moisture requirements and durations, so that a partial rainfall failure does not translate into a total crop loss. Third, the resulting ground cover conserves residual soil moisture, moderates soil temperature, and is reported by adopting farmers to extend a field’s capacity to withstand dry spells by roughly ten to fifteen days compared with untreated fields (Kurnool district administration campaign, 2026).

5.3 Mobilising PMDS at Field Level

Because PMDS must be completed within a narrow pre-monsoon window, the extension task is essentially a logistics and mobilisation exercise rather than a conventional training exercise. Field-level campaigns documented in Andhra Pradesh have combined several elements: identification of the target sowing window based on historical onset dates and any available seasonal outlook; door-to-door and camp-based distribution of a standard seed kit covering a locally appropriate mix of millet, pulse, oilseed, and vegetable varieties; community-level seed-

treatment demonstrations at a common collection point so that farmers can replicate the bio-inoculant treatment at home; and a village-level target (in terms of acreage) that is tracked and reported upward through the extension hierarchy so that coverage gaps can be identified and addressed while the sowing window is still open.

Adoption studies indicate that, where PMDS has been promoted intensively, the technique has been taken up on roughly a third of the cultivated area of participating farmers, with poly-crop mixtures, pulses, and vegetables forming the bulk of the sown area (Institute for Development Studies Andhra Pradesh, 2022). The same studies also flag the main constraints extension planners must budget for — the added cost of seed and mulching material, and shortages of labour and locally sourced organic inputs at peak season — both of which point to the value of pre-season input aggregation through Farmer Producer Organisations or KVK-managed seed hubs.

6. Promoting Short-Duration Contingency Crops: Millets and Pulses

Where the sowing window for a district's principal kharif crop has already been missed, or where an early-season dry spell has damaged a standing crop beyond recovery, the DACP framework recommends substitution with short-duration crops capable of completing their life cycle within the residual moisture period. Small millets and pulses are particularly well suited to this role because of their inherently short duration, low water requirement, and tolerance of moisture stress at critical growth stages.

Commonly recommended contingency options in dryland tracts include short-duration millets such as foxtail millet, little millet, kodo millet, and early-maturing pearl millet hybrids, alongside pulses such as green gram, black gram, horse gram, and cowpea, and short-duration oilseeds such as sesamum where soil and market conditions permit. The extension task in promoting these crops has three components that must run in parallel: technical validation of which contingency crop and variety fits the specific delay or moisture deficit observed in a given block (a decision that follows directly from the relevant DACP); assured availability of certified or truthfully labelled seed of the recommended short-duration variety through KVK seed hubs, State Seed Corporations, or Farmer Producer Organisations, since standard seed supply chains are usually built around the district's dominant long-duration crop; and rapid communication to farmers through village-level meetings, mobile agro-advisories, and local Krishi Vigyan Kendra staff so that the substitution decision reaches the field while a usable sowing window still remains.

Millets and pulses also carry a nutritional and market rationale that strengthens their case as contingency crops rather than purely emergency stop-gaps: their promotion under national and state millet-mission initiatives means that, unlike many one-off contingency measures, infrastructure for processing, procurement, and market linkage is increasingly available, which

improves the likelihood that farmers adopting them under duress in one season will continue to grow them as a deliberate risk-diversification choice in subsequent seasons.

7. Training Farmers in Moisture Conservation

Contingency cropping decisions are only as effective as the soil's capacity to retain whatever rainfall is received, which makes moisture-conservation training a necessary companion to both PMDS and contingency-crop promotion. Extension training in this area typically covers two broad categories of practice.

7.1 *In-Situ* Moisture Conservation

In-Situ measures aim to retain rainfall where it falls and reduce non-productive water loss. Field-level training modules commonly cover contour and graded bunding on sloping land, broad-bed-and-furrow and ridge-furrow systems that improve infiltration while draining excess water during intense spells, dead and live mulching using crop residue or the ground cover generated by a PMDS poly-crop stand, and minimum or conservation tillage that limits moisture loss associated with repeated conventional ploughing. Vegetative barriers and grass strips along field boundaries have also been documented as a low-cost means of reducing runoff and soil loss on sloping dryland fields (Pushpanjali *et al.*, 2023).

7.2 *Ex-Situ* / Supplementary Measures

Where feasible, farmers are trained to construct or maintain farm ponds and other small water-harvesting structures that capture runoff for a single life-saving irrigation during a critical dry spell, and to use simple soil-moisture indicator tools to decide when such a supplementary irrigation is actually justified rather than applying it on a fixed schedule. Community-level training also covers safe use of harvested water — prioritising critical crop stages such as flowering and grain filling — so that a limited water source is allocated for maximum yield protection rather than spread thinly across an entire field.

Delivery of this training works best through a combination of hands-on field demonstrations at the FLD plots described in Section 4, short video- and mobile-based content that can be shared through farmer WhatsApp groups and Kisan Call Centres, and structured exposure visits in which farmers from a vulnerable village visit an established demonstration site to see the practice functioning under real dry-spell conditions.

8. A Field-Level Campaign Model for Rapid Mobilisation

Translating the strategies above into a coordinated response requires a campaign structure with a clear trigger, a compressed timeline, and defined institutional roles. Table 1 summarises a generalised phase-wise model, adapted from documented DACP-interface and PMDS mobilisation practice, that extension planners can adjust to local monsoon climatology and administrative capacity.

Table 1: Phase-wise field campaign model for El Niño contingency response

Phase	Trigger / Timing	Key Actions	Lead Actors
1. Alert	Pre-season seasonal outlook indicating El Niño / below-normal rainfall probability	Review district DACP; identify likely-affected blocks; quantify contingency seed requirement	CRIDA / SAU, State Dept. of Agriculture, KVK
2. Preparedness	6–8 weeks before expected monsoon onset	Pre-position PMDS and short-duration seed kits; conduct interface meeting with line departments; identify lead farmers and demonstration plots	KVK, Line Departments, FPOs
3. Mobilisation	45–60 days before onset (PMDS window)	Village-level camps for seed treatment and distribution; door-to-door PMDS sowing coverage; FLD-based demonstrations	Extension staff, Lead farmers, SHGs
4. Monitoring	Through the sowing and early-growth period	Track dry-spell days and germination status; trigger contingency-crop substitution advisory if thresholds are crossed	KVK, IMD/agromet units, block staff
5. Response & Feedback	Post-sowing to harvest	Moisture-conservation follow-up training; documentation of coverage and outcomes; feedback to update the next season's DACP	KVK, SAU, State Dept. of Agriculture

The model is deliberately compressed: Phases 1 and 2 must be substantially complete before the PMDS sowing window opens, since the value of anticipatory sowing and contingency-seed pre-positioning collapses once the window has passed. This places a premium on institutional readiness — pre-approved seed procurement budgets, standing lists of lead farmers, and pre-identified demonstration plots — well before an El Niño signal is confirmed, rather than initiating these steps only after the season has visibly gone wrong.

9. Institutional Convergence and Policy Support

No single institution can execute the campaign model described above in isolation. ICAR-CRIDA provides the technical backbone through DACPs and the NICRA network of vulnerable-district demonstrations; State Agricultural Universities and KVKs provide the last-mile scientific and extension presence; State Departments of Agriculture control seed procurement, subsidy administration, and crop-insurance interface; and community institutions — Farmer Producer Organisations, Self-Help Groups, and programmes such as APCNF — provide the social

infrastructure through which door-to-door mobilisation and peer-to-peer learning actually happen at village scale.

Linking crop insurance awareness to contingency campaigns is a further point of convergence worth strengthening: since crop insurance under schemes such as the Pradhan Mantri Fasal Bima Yojana covers physical yield loss due to drought but not price volatility, extension messaging delivered alongside PMDS and contingency-crop campaigns should clarify this distinction so that farmers do not mistake insurance enrolment for a substitute for on-field risk-mitigation practice.

Digital agro-advisory channels — including SMS- and app-based services and the Kisan Call Centre network — offer a low-cost means of compressing the time between a confirmed weather trigger and farmer-level action, particularly for time-bound advisories such as the opening or closing of a PMDS sowing window or a switch recommendation to a specific contingency crop.

10. Challenges and Constraints

Field experience with grassroots contingency extension points to several recurring constraints that planners need to budget for explicitly:

- A narrow, non-negotiable sowing window for PMDS and contingency crops leaves little room for administrative delay in seed procurement or fund release.
- Additional costs of seed, bio-inoculant inputs, and mulching material can deter adoption among resource-poor farmers unless subsidised or aggregated through community institutions.
- Seasonal shortages of labour and locally available organic inputs (dung, mulching biomass) at peak sowing time constrain the scale at which PMDS protocols can be fully implemented on a given holding.
- The extension worker-to-farmer ratio in many dryland blocks remains too thin to deliver door-to-door mobilisation at the pace a compressed campaign requires, making reliance on trained lead farmers and community volunteers essential.
- Seed systems for short-duration millets and pulses are frequently underdeveloped relative to the district's dominant long-duration crop, creating availability bottlenecks precisely when substitution is most needed.
- Repeated, one-off training without follow-up support tends to produce partial or inconsistent adoption of moisture-conservation practices in subsequent normal-rainfall seasons.

11. Way Forward

Building on documented experience, the following measures can strengthen the grassroots contingency-extension chain for future El Niño and moisture-stress years:

- Pre-approve seasonal contingency budgets and seed-procurement authority at the district level so that Phase 1 and 2 actions in the campaign model can begin as soon as a seasonal outlook is issued, without awaiting fresh administrative sanction.
- Establish KVK- or FPO-managed community seed and bio-input hubs for PMDS and short-duration contingency crops, reducing per-farmer cost and last-mile availability constraints.
- Formally integrate a standing panel of trained lead farmers into the extension structure of each vulnerable block, to host frontline demonstrations and lead door-to-door mobilisation during the compressed sowing window.
- Strengthen the link between agromet advisory units and extension delivery so that dry-spell monitoring translates into a contingency-crop switch advisory within days rather than weeks.
- Institutionalise post-season documentation of campaign coverage and outcomes, feeding directly into the annual revision of District Agriculture Contingency Plans.
- Ensure that women farmers and landless agricultural labour households, who form a substantial share of the dryland workforce, are directly included in seed-treatment camps and moisture-conservation training rather than reached only indirectly through male household heads.

Conclusion

El Niño years will continue to test the responsiveness of India's agricultural extension system, not because the required technologies are unknown, but because the window within which they must be delivered is short and unforgiving. Pre-monsoon dry sowing, frontline demonstration of resilient technology bundles, rapid substitution with short-duration millets and pulses, and structured moisture-conservation training together form a practical, field-tested toolkit for grassroots contingency response. Their combined effectiveness, however, depends less on the technologies themselves than on whether extension systems are organised — institutionally, financially, and logistically — to act at the pace a moisture-stressed cropping season demands. Embedding a standing, pre-approved campaign structure of the kind outlined in this chapter, tightly linked to existing District Agriculture Contingency Plans and community institutions, offers a realistic pathway to closing that gap.

References

1. Institute for Development Studies Andhra Pradesh. (2022). *Community managed natural farming: An integrated approach with pre-monsoon dry sowing farming in Andhra Pradesh, India*. Institute for Development Studies Andhra Pradesh, Visakhapatnam.

2. Institute for Development Studies Andhra Pradesh. (2023). *Pre-monsoon dry sowing: A grassroots climate adaptation practice*. Andhra Pradesh Community Managed Natural Farming Programme Report, Visakhapatnam.
3. Economic and Political Weekly. (2025). Improving yield efficiency through pre-monsoon dry sowing in Andhra Pradesh. *Economic and Political Weekly*, 60(11). Web Exclusive.
4. Sreenivasa, M. N., Naik, N., & Bhat, S. N. (2009). Beejamrutha: A source for beneficial bacteria. *Karnataka Journal of Agricultural Sciences*, 22(5), 1038–1040.
5. Pushpanjali, Reddy, K. S., Rao, C. A. R., Vijaya Kumar, P., & Sammi Reddy, K. (2023). Fodder grass strips: An affordable technology for sustainable rainfed agriculture in India. *Agriculture*, 13(2), 318.
6. Rama Rao, C. A., Raju, B. M. K., Rao, A. V. M. S., & Venkateswarlu, B. (2022). Impact of climate change on productivity of foodcrops: A sub-national level assessment for India. *Environmental Research Communications*, 4(9), 095011.
7. Venkateswarlu, B., & Shanker, A. K. (2012). *Dryland agriculture: Bringing resilience to rainfed agriculture*. ICAR-Central Research Institute for Dryland Agriculture, Hyderabad.
8. ICAR-Central Research Institute for Dryland Agriculture. (2017). *AICRPDA-NICRA annual report*. Central Research Institute for Dryland Agriculture, Hyderabad.
9. ICAR-Central Research Institute for Dryland Agriculture. (n.d.). *District level agriculture contingency plans*. <https://www.icar-crida.res.in>
10. National Innovations in Climate Resilient Agriculture (NICRA). (n.d.). *Indian Council of Agricultural Research*. <http://www.nicra-icar.in>
11. International Crops Research Institute for the Semi-Arid Tropics. (2013). *Bhoochetana: Bridging yield gaps with science-led interventions in Andhra Pradesh*. ICRISAT, Patancheru.
12. Kurnool District Administration. (2026). *Pre-monsoon dry sowing campaign amid El Niño-induced rainfall deficit*. District Agriculture and Irrigation Departments, Government of Andhra Pradesh.
13. Andhra Pradesh Community Managed Natural Farming (APCNF). (n.d.). *Scaling pre-monsoon dry sowing*. <https://apcnf.in>

GREEN NANOREMEDIATION: SUSTAINABLE ADVANCES IN ENVIRONMENTAL REMEDIATION MECHANISMS AND APPLICATIONS

Rashmi Verma* and Tanzeel Ahmed

School of Biotechnology,
IFTM University, Moradabad – 244102

*Corresponding author E-mail: rashmi.verma@iftmuniversity.ac.in

Abstract

Green Nano remediation represents the integration of Nanotechnology, green chemistry and Environmental Science studies, offering innovative eco-friendly, cost-effective solutions for maintaining healthy ecosystem free from any pollutant. The traditional methods (physical and chemical methods) involve the use of hazardous reagents, require high energy inputs, and generate secondary toxic waste. These shortcomings can be met by using green nanoparticle synthesis. The mechanism involved includes adsorption, redox transformation, photo-catalytic degradation, and targeted ion exchange. These methods can successfully function in diverse environmental conditions to remove heavy metal (such as hexavalent chromium, lead, and arsenic), organic dyes, pesticides, pharmaceuticals, and emerging micro-pollutants from contaminated soil, groundwater, and wastewater. However, the utilization of biogenic NPs also poses various challenges such as standardized green synthesis protocols and comprehensive lifecycle toxicity assessments. Green nanoremediation acts as a promising model for sustainable environmental restoration, combining highly efficient pollutant removal with less ecological impacts.

Keywords: Green Nanoremediation, Green Chemistry, Heavy Metals, Environmental Pollution.

Introduction

Environmental pollution of soil, water and air is globally recognized menace that destabilizes the ecosystems, public health and Biodiversity. Rapid industrialization, disruptive agricultural practices, mining and unplanned urbanization results in addition of pollutants in soil and water bodies (Korlam *et al.*, 2025.). These pollutants include heavy metals such as cadmium, lead, and arsenic and Organic pollutant Polychlorinated biphenyls and pesticides (Özkara *et al.*, 2016) can pose the menace to crop health and food safety (Su *et al.*, 2014). Heavy metals contamination by agricultural runoff can affect freshwater ecosystems as they accumulate in aquatic organisms at each trophic level causing bioaccumulation and biomagnifications in the food web resulting in health threat to aquatic life such as loss of reproductive abilities and abnormal development in fish, which eventually will lead to threat to human health and also decrease the crop yield (Ahmed *et al.*, 2021). The exposure to these pollutants can cause various public health issues i.e.

cancer, skin, lung kidney and brain diseases. These interrelated threats paved the way for development of an advanced remediation technology i.e. Green Nano remediation that offers high efficiency, high selectivity and sustainability.

Traditional physical synthesis of NPs includes sonication, irradiation, laser removal, arc discharge and Pyrolysis. Chemical methods produce toxic by products due to utilization of reducing compounds sodium Borohydrate and hydrazine (Abdelkhalek *et al.*, 2022). High cost and use of toxic chemicals motivated the researchers to develop safer and environment sustainable method i.e. green synthesis of Nanoparticles for bioremediation (Patil.*et al.*, 2020).

Green Nanotechnology utilizes the natural properties of reducing and balancing of plant extracts and microorganism or enzymes to manufacture Nanoparticles (NPs) of size range 1 to 100 nm making it less toxic, environment friendly and cost-effective option over traditional physical and chemical synthesis. Owing to its unique physio chemical properties like high reactivity, high surface area to volume ratio, adjustable surface chemistry to eliminate pollutant it is often termed as Green Nano remediation (Singh *et al.*, 2017).

2. Types of Nano materials-

A wide range of nanomaterials has been manufactured for various environmental concerns based on their properties to clean up.

2.1 Carbon based Nanomaterials (CNTs & Graphene)

Carbon-based nanomaterials, including carbon nanotubes (CNTs), graphene, and graphene oxide are used to tackle environmental concerns due to their high surface area and unique electronic properties (Fei *et al.*, 2022). The persistent organic pollutants and heavy metals are adsorbed effectively by CNTs (O. Vázquez-Núñez *et al.*, 2020). The biodegradation of toxic compounds is enhanced by Graphene and its derivatives due to their high reactivity and potential to serve as redox mediators their high reactivity and potential to serve as redox intermediaries (Raj *et al.*, 2024).

2.2 Nanoscale Zero-Valent Iron (nZVI)

Nanoscale Zero-Valent Iron (nZVI) can immobilize or degrade a broad spectrum of contaminants owing to its high reactivity thus making it suitable for in situ remediation (Karn *et al.*, 2009). It has a limitation of rapid oxidation and agglomeration that was resolved by coating it with polymer, mixing it with other metals (bivalent NPs) or using embedded support such as zeolites (Mukhopadhyay *et al.*, 2021).

2.3 Silica and Polymeric Nanomaterials

Silica NP (SiO₂ NPs) are preferred NPs due to easy operations, environment friendly nature and bioremediation application in both water and soil (Vimal *et al.*, 2026). The mechanism of action involves the immobilization and degradation using catalytic reactions.

2.4 Metal and Metal Oxide Nanoparticles

Metal Oxide Nanoparticles (e.g., TiO_2 , ZnO , and Fe_3O_4) are known for their antimicrobial, adsorption and photo catalytic properties (Serov *et al.*, 2024). Iron oxides (Fe_3O_4) have added benefit that they can be easily separated from treated water using external magnetic fields (Alazaiza, 2021).

2.5 Novel Nanomaterials (MOFs, COFs, Nanosponges)


The revolution in the field of NPs has led to development of advanced novel structures like Metal-Organic Frameworks (MOFs) and Covalent Organic Frameworks (COFs) that are highly efficient in organic pollutant degradation owing to their well-defined pore structures and high surface areas. Herbicide Triclopyr can be successfully degraded by cyclodextrin-based nanosponge (Fei *et al.*, 2022).

3. Green synthesis of NPs

Synthesis of NPs using biological material such as plant extract, microorganism and enzymes.

3.1 Plant based NP synthesis-

Various metal and metal oxide NPs are successfully extracted from plant parts such as leaves, seeds, flowers, roots (Bolade *et al.*, 2020). The synthesis process commences with mixing of plant extract containing polyphenols, terpenoids, alkaloids and flavanoids which act as reducing agent with precursor salt solution (e.g Ferric chloride and Silver nitrate). Positively charged metal ions are electron deficient and reducing agent donate electron rendering them stable uncharged metal atoms. The metal atoms start colliding and aggregate together to form cluster called nuclei. More atoms attach to nuclei and get clustered to form large structured NP. This complex is then stabilized by capping with different bio molecule and thus stops accumulation of more metal atoms to the Nanoparticle (Heinemann, M.G. 2021).



Intracellular mechanism involves the entrapment of positively charged metal ions by negatively charged group present on the cell wall. This metal ion then diffuses across the cell wall and enter

inside cell it is acted upon by enzymes present. These enzymes act as reducing agent render the metal ion in uncharged form. These metal atoms cluster and form NP (Mughal, 2021). Plants that can act as bio factories of NP are stated in the table 1.

3.2 Microbial and Enzymatic synthesis

This synthesis pathway utilizes the metabolic pathways of micro-organisms such as bacteria, fungi to manufacture the NPs and this process can be carried out both intracellular and extracellular ways (Ferdush *et al.*, 2025).

- **Bacterial synthesis:** In bacterial mediated synthesis the bacterial filtrate act as reducing agent that neutralizes the electron deficient metal ion. The intracellular synthesis involves the electrostatic attraction between metal ion and anionic group present on the plasma membrane. The metal ion than travels inside bacteria cell and enzymes act on it to reduce it to uncharged metal atom. This atom than form clusters of NPs and after getting perfect size the NPs get capped. However, the amount manufactured is very less as compared to extracellular manufacturing of NPs (Mughal, 2021). Bacterial cells can act as bio factories of NP as it has the ability to synthesize various enzymes, pigments, secondary metabolites as stated in the table 2.
- **Fungal synthesis:** For large scale production of NPs, the most preferred candidate is fungus (Table 3) due to their capability to secrete enzymes (acetyl xylan esterase, nitrate reductase), quinines and peptides that can convert precursor material into NPs (Bhandari *et al.*, 2023). Mycelia formation covering large surface area, high metal tolerance, High Growth rate, Easier harvesting, high enzyme biosynthesis rate confers more importance to utilize fungus for NP production as compared to bacteria mediated biosynthesis (Mukherjee *et al.*, 2001).
- **Algal and enzymatic synthesis:** Phyconanotechnology has potential to revolutionize the synthesis of NPs from algae. Algal cells can act as bio factories of NP as it has the ability to synthesize various enzymes, pigments, secondary metabolites as stated in the table 4.

4. Mechanism of action on pollutant

The mechanism that involves the removal of heavy metals, dye microplastics etc are adsorption, catalysis and membrane filtration. Pollutant-NP interactions depend on the Physical and chemical properties of the NP such as shape, size, surface area and charge and surface chemistry thus affecting the efficacy of remediation processes. Small size of the ZnO and Fe₃O₄ NPs posses' large surface area and therefore more available sites for adsorption which is helpful in capturing heavy metal and dyes (Mehta *et al.*, 2024). Catalytic method is based on the degradation organic pollutant and dyes with the help of metallic nano particles (Moond *et al.*, 2024). Use of engineered NP with conventional methods can be highly efficient method to tackle

the menace of emerging pollutants like personal care products, pharmaceuticals and nano/micro plastics (Ugwuanyi *et al.*, 2024).

4.1 Adsorption

It involves both physical and chemical process for removal of pollutants by NPs. The physical process consists of reversible binding of pollutant to NP surface by weak Van der Waals forces while chemical adsorption is irreversible binding by strong ionic or covalent bonds (Gobinath R *et al.*, 2021). In waste water treatment, Fe₃O₄ or Magnetite NP has high surface area and is capped with functional groups such as -OH & -NH₂ electrostatically bind and form complex with heavy metal ions (Pb²⁺, Cd²⁺) (Lu *et al.*, 2025). A wide variety of organic, inorganic (heavy metals and dyes) and microbial pollutant along with emerging contaminants (ECs) such as personal care products, pharmaceuticals can be efficiently removed by using activated carbon as NP (Almeida-Naranjo *et al.*, 2023). This method is preferred due to its efficiency and low cost to cope with the water and air pollution challenges.

4.2 Photo catalytic degradation method

This is the pollutant degradation technique which occurs in presence of light and catalyst to convert it into nontoxic substances such as water and carbon di-oxide. It's more effective than adsorption as it degrades pollutant not remove also energy efficient. Black nanoparticle TiO₂ degrade the organic pollutant (decolourize dyes like methyl orange) by photo catalysis and generation of reactive oxygen species (ROS) i.e hydroxyl free radical and superoxide anions. These ROS convert organic pollutants into nontoxic substances such as water and carbon dioxide (Nawaz *et al.*, 2024).

4.3 Electrochemical Reduction

This process involves electro deposition and electro-sorption in which NP act as catalyst that donate electrons to heavy metal cation and reduce them to safer form in presence of electrodes. Soluble Hexavalent Chromium (VI) are reduced by Zero-valent Aluminium to trivalent Chromium (III) that precipitate out naturally in waste water treatment (Patricia Treviño *et al.*, 2014).

5. Challenges of Green Nano Remediation

Green nano-remediation despite being the safest and innovative player in the field of bioremediation but fails to find its application in real world from laboratories due to various serious limitations. The limitations are discussed in detail below

5.1 High Cost and Scalability Issues: Scalability becomes difficult due to high cost and technical problems in manufacturing the green NPs at industrial level as the energy demand increases too that limit its commercialization for large remediation project. The green NPs developed in labs often failed to perform in variable field conditions so it's advisable to conduct

live ecosystem assessments to develop refined NPs that can withstand harsh field conditions. pH, dissolved organic matter, temperature and salinity can damage the capping on NPs resulting in precipitation of NPs (Kaur *et al.*, 2023). The biological extract used in synthesis may vary in composition according to species, season, age which result in poor reproducibility in green NP production (Iravani, 2011). The recovery of green NPs for its reuse is difficult and thus can lead to increased operational cost (Karn *et al.*, (2009).

5.2 Efficiency: During storage, green NP may form clusters or aggregates or in contaminated environment which cause the reduced surface area issue and ultimately the low adsorption efficiency (Singh, P *et al.*, 2016).

5.3 Environmental and Health Issues: The long-term effects of nanoparticles in the environment are not fully understood, raising concerns about their persistence and bioaccumulation (Singh, 2025). Some nanoparticles transform pollutants into intermediate compounds rather than complete mineralization resulting in formation of potentially toxic by-products (Tratnyek & Johnson (2006)). These can be toxic to aquatic organisms, beneficial microbes, and plants at high concentrations and leads to disturbance of microbial communities and ecosystem functions (Kahru & Dubourguier, 2010).

5.4 Regulatory and Ethical Challenges: The rapid development of nanotechnology has outpaced regulatory measures, leading to uncertainties in safety and efficacy standards (Singh, 2025). The deployment of nanotechnology in remediation raises ethical questions regarding its impact on communities and ecosystems (Singh, 2025).

6. Future Perspectives

The futuristic insight about the use of green Nanoparticles in bioremediation involves the development of multifunctional, reusable, recoverable and environmentally friendly nanomaterials that can be effective in real environmental conditions. The efficiency of this NP can be improved by using integrated plant, microbial, algal and agricultural waste mediated synthesis and applying various methods of remediation such as photocatalysis, magnetic separation, biosensing and Artificial intelligence. However, lab to field application requires standardized synthesis protocols, life-cycle assessment, toxicological evaluation and large-scale production. A new generation of green nanoparticle development require a combination approach such as biotechnology, microbial and enzymatic bioremediation, green synthesis technology, circular economy principles and Artificial intelligence for precision-oriented bioremediation technologies.

6.1 Production of Green Nanoparticles from Agricultural Waste

Use of agricultural waste as a raw material for production of green nanoparticle holds the future of green nanotechnology. Huge amount of agricultural waste such as rice husk, peels, sugarcane

bagasse, wheat straw etc are produced annually that contain cellulose, lignin, hemi cellulose, silica and other bioactive compounds. These compounds can act as a support material for the production green nanomaterials (Bhushan, 2024)

6.2 Magnetic Nanoparticles

Magnetic green nanoparticles, particularly Fe_3O_4 -based materials, could become important because they can be separated from treated water using an external magnetic field. This would address one of the major limitations of nanoparticle-based remediation: recovery of nanoparticles after treatment. Immobilization of nanomaterials on membranes, electrodes or other supports is also recommended to minimize environmental release and enable reuse (Huang *et al.*, 2024)

6.3 Smart nanoparticles

Future GNPs could be engineered to respond to environmental conditions such as pH, temperature, light, redox potential and specific pollutants. Smart nanoparticle could remain relatively inactive under normal conditions but become catalytically active when it encounters a particular pollutant or receives visible light. This would allow controlled and targeted remediation rather than nonspecific nanoparticle activity.

6.4 Solar-driven remediation

An integrated approach solar photolysis and green nanoparticles to develop visible-light-responsive nanoparticles capable of degrading pollutants using sunlight. This could be particularly useful for decentralized wastewater treatment in rural and agricultural areas. Photocatalysis is already a major research direction in green nanomaterial-based wastewater treatment (Roopa *et al.*, 2025)

6.5 Nanosensor based remediation

Green nanoparticles could serve as components of biosensors for detecting heavy metals, pesticides, dyes and other contaminants, followed by targeted remediation. Nanotechnology is already being investigated across biosensing, membrane technology and advanced oxidation processes (Rathod *et al.*, 2024).

6.6 Artificial intelligence-assisted nanoparticle

AI and machine learning could become increasingly useful for predicting nanoparticle size, morphology, surface charge, stability, adsorption capacity, photocatalytic efficiency, toxicity, and optimal synthesis conditions. This could reduce the experimental trial-and-error involved in optimizing green nanoparticle synthesis (Tiwari *et al.*, 2026).

Table 1: Plant derived Nanoparticle used for bioremediation

NP	Plant source	Plant part used	Biomolecule	Size range (nm)	Application	References
Silver (Ag)	<i>Azadiracta indica</i>	Leaves	Azadirachtin & terpenoids	15–50	Antibacterial, Wound Healing, dye degradation, waste water treatment	Ahmed <i>et al.</i> (2016)
Silver (Ag)	<i>Camellia sinensis</i> (Tea)	leaves	Catechins, tannins, polyphenols	15–50	Degradation of organic pollutants	Mittal <i>et al.</i> (2013)
Silver (Ag)	<i>Cucurbita</i> spp.	Fruit peel	Carotenoids & soluble sugars	70–85	Biopesticides, reduction of harmful agricultural nitrite run off	Eker <i>et al.</i> (2025)
Silver (Ag)	<i>Ocimum sanctum</i>	leaves	Eugenol, phenolics, flavonoids	15–45	Antimicrobial, antioxidant	Mittal <i>et al.</i> (2013)
Gold (Au)	<i>Camellia sinensis</i> (Tea)	leaves	Epigallocatechin gallate (EGCG)	5–30	Cancer cell Imaging	Rabiee (2025)
Gold (Au)	<i>Aloe vera</i>	leaves	Anthraquinones, polysaccharides, proteins	20–80	Catalytic reduction of nitroaromatic compounds	Iravani (2011)
Zinc oxide (ZnO)	<i>Moringa oleifera</i>	Leaves	Phenolics, flavonoids, ascorbic acid	20–70	UV blocking, antibacterial, Photocatalytic dye degradation	Singh <i>et al.</i> (2016)
Zinc oxide (ZnO)	<i>Camellia sinensis</i>	Leaves	Catechins, polyphenols	15–60	Antimicrobial, food packaging, cosmetics	Singh <i>et al.</i> (2016)
Iron oxide	<i>Camellia sinensis</i>	Leaves	Catechins, polyphenols	20–100	Waste water treatment, drug delivery	Nasrollahzadeh <i>et al.</i> (2019)
Magnesium Oxide (MgO)	<i>Mangifera indica</i>	Leaves	Mangiferin, phenolics	20–80	Antibacterial, adsorption of heavy metal and dye	Salam <i>et al.</i> (2019)

Table 2: Bacteria derived Nanoparticle used for bioremediation

Nanoparticle	Bacterial Source	Key Biomolecules Involved in Synthesis	Typical Size (nm)	Bioremediation Applications	Representative Reference
Silver nanoparticles (AgNPs)	<i>Bacillus licheniformis</i>	Nitrate reductase, proteins, amino acids, exopolysaccharides (EPS)	10–50	Antimicrobial wastewater treatment, dye degradation	Kalishwaralal <i>et al.</i> (2008); Singh <i>et al.</i> (2016)
Silver nanoparticles (AgNPs)	<i>Bacillus subtilis</i>	NADH-dependent reductase, peptides, EPS	15–60	Textile dye degradation, wastewater disinfection	Singh <i>et al.</i> (2016)
Silver nanoparticles (AgNPs)	<i>Pseudomonas stutzeri</i>	Reductases, cytochromes, proteins	20–80	Heavy metal removal, antimicrobial treatment	Klaus <i>et al.</i> (1999)
Gold nanoparticles (AuNPs)	<i>Rhodopseudomonas capsulata</i>	NADH reductase, proteins, carotenoids	10–40	Catalytic degradation of organic pollutants	He <i>et al.</i> (2007)
Gold nanoparticles (AuNPs)	<i>Escherichia coli</i>	Enzymes, membrane proteins, glutathione	5–30	Reduction of nitroaromatic pollutants	Narayanan & Sakthivel (2010)
Zinc oxide nanoparticles (ZnO NPs)	<i>Lactobacillus plantarum</i>	Proteins, enzymes, EPS	20–80	Photocatalytic degradation of dyes, antibacterial wastewater treatment	Singh <i>et al.</i> (2016)
Zinc oxide nanoparticles (ZnO NPs)	<i>Bacillus megaterium</i>	Extracellular enzymes, peptides	30–70	Heavy metal adsorption, dye degradation	Fouda <i>et al.</i> (2018)
Iron oxide nanoparticles (Fe ₃ O ₄ NPs)	<i>Shewanella oneidensis</i>	Cytochromes, reductases, extracellular proteins	10–30	Arsenic, chromium and uranium removal	Lovley <i>et al.</i> (2004)

Iron oxide nanoparticles (Fe ₃ O ₄ NPs)	<i>Geobacter sulfurreducens</i>	c-type cytochromes, pili proteins	20–60	Heavy metal immobilization, groundwater remediation	Lovley <i>et al.</i> (2004)
Copper oxide nanoparticles (CuO NPs)	<i>Bacillus coagulans</i>	Proteins, enzymes, EPS	20–70	Organic dye degradation and antimicrobial remediation	Fouda <i>et al.</i> (2018)
Copper nanoparticles (CuNPs)	<i>Pseudomonas fluorescens</i>	Reductases, proteins	15–50	Catalytic degradation of pollutants	Narayanan & Sakthivel (2010)
Selenium nanoparticles (SeNPs)	<i>Bacillus mycoides</i>	Selenite reductase, glutathione, proteins	50–150	Removal of selenium oxyanions, heavy metal detoxification	Cremonini <i>et al.</i> (2016)
Selenium nanoparticles (SeNPs)	<i>Stenotrophomonas maltophilia</i>	NADPH reductase, glutathione	80–200	Detoxification of selenium-contaminated wastewater	Presentato <i>et al.</i> (2018)
Palladium nanoparticles (PdNPs)	<i>Desulfovibrio desulfuricans</i>	Hydrogenase, cytochromes	2–15	Catalytic dechlorination of groundwater pollutants	Deplanche <i>et al.</i> (2011)
Platinum nanoparticles (PtNPs)	<i>Acinetobacter calcoaceticus</i>	Hydrogenase, reductases, proteins	2–10	Catalytic degradation of chlorinated compounds	Konishi <i>et al.</i> (2007)

Table 3: Fungus derived Nanoparticle used for bioremediation

Nanoparticle	Fungal Source	Key Biomolecules Involved	Typical Size (nm)	Bioremediation Applications	Representative References
Silver (AgNPs)	<i>Fusarium oxysporum</i>	Nitrate reductase, NADH-dependent reductase, proteins, quinones	5–50	Textile dye degradation, antimicrobial wastewater treatment, degradation of organic pollutants	Ahmad <i>et al.</i> , 2003; Narayanan & Sakthivel, 2010
Silver (AgNPs)	<i>Aspergillus niger</i>	Extracellular proteins, enzymes, polysaccharides	10–60	Dye degradation, wastewater disinfection, heavy metal adsorption	Guilger-Casagrande & Lima, 2019
Silver (AgNPs)	<i>Aspergillus flavus</i>	Proteins, reductases, amino acids	8–40	Removal of pathogenic microorganisms, heavy metal remediation	Vigneshwaran <i>et al.</i> , 2007
Silver (AgNPs)	<i>Penicillium chrysogenum</i>	Proteins, peptides, enzymes	10–45	Textile effluent treatment, antibacterial remediation	Fouda <i>et al.</i> , 2018
Silver (AgNPs)	<i>Trichoderma harzianum</i>	Enzymes, proteins, phenolic compounds	5–25	Dye degradation, antimicrobial remediation	Guilger-Casagrande & Lima, 2019
Gold (AuNPs)	<i>Fusarium oxysporum</i>	NADH reductase, proteins	15–40	Catalytic degradation of nitroaromatic compounds and organic pollutants	Mukherjee <i>et al.</i> , 2002
Gold (AuNPs)	<i>Aspergillus niger</i>	Proteins, reductases	10–30	Catalytic oxidation of environmental contaminants	Narayanan & Sakthivel, 2010
Iron oxide (Fe ₃ O ₄ NPs)	<i>Trichoderma harzianum</i>	Proteins, extracellular enzymes	15–45	Removal of Cr(VI), Pb ²⁺ , Cd ²⁺ and As(V)	Fouda <i>et al.</i> , 2018

Iron oxide (Fe ₂ O ₃ NPs)	<i>Aspergillus tubingensis</i>	Proteins, polysaccharides	20–60	Heavy metal adsorption and groundwater remediation	Singh <i>et al.</i> , 2018
Zinc oxide (ZnO NPs)	<i>Aspergillus fumigatus</i>	Proteins, amino acids, polysaccharides	20–70	Photocatalytic degradation of methylene blue, Congo red and Rhodamine B	Fouda <i>et al.</i> , 2018
Zinc oxide (ZnO NPs)	<i>Penicillium expansum</i>	Proteins, enzymes	15–50	Textile wastewater treatment, organic dye degradation	Singh <i>et al.</i> , 2018
Copper oxide (CuO NPs)	<i>Penicillium chrysogenum</i>	Proteins, peptides, amino acids	10–50	Heavy metal adsorption, dye degradation	Fouda <i>et al.</i> , 2018
Copper oxide (CuO NPs)	<i>Aspergillus terreus</i>	Extracellular enzymes, proteins	20–55	Degradation of industrial organic pollutants	Guilger-Casagrande & Lima, 2019
Titanium dioxide (TiO ₂ NPs)	<i>Aspergillus flavus</i>	Proteins, polysaccharides	10–40	Photocatalytic degradation of pesticides and pharmaceuticals	Singh <i>et al.</i> , 2018
Selenium (SeNPs)	<i>Saccharomyces cerevisiae</i>	Glutathione, reductases, proteins	50–150	Selenium detoxification, heavy metal remediation	Cremonini <i>et al.</i> , 2016
Palladium (PdNPs)	<i>Aspergillus niger</i>	Hydrogenase-like enzymes, proteins	5–20	Catalytic dechlorination of chlorinated organic pollutants	Narayanan & Sakthivel, 2010
Platinum (PtNPs)	<i>Fusarium oxysporum</i>	Reductases, proteins	2–15	Catalytic degradation of persistent organic pollutants	Narayanan & Sakthivel, 2010

Table 4: Algae derived Nanoparticle used for bioremediation

Nanoparticle	Algal Source	Key Biomolecules Involved	Typical Size (nm)	Bioremediation Applications	Representative References
Silver (AgNPs)	<i>Chlorella vulgaris</i>	Proteins, polysaccharides, phenolics, chlorophyll	10–30	Antimicrobial wastewater treatment, dye degradation	Dahoumane <i>et al.</i> (2017); Roychoudhury <i>et al.</i> (2016)
Silver (AgNPs)	<i>Spirulina platensis</i> (<i>Arthrospira platensis</i>)	Phycocyanin, proteins, polysaccharides	20–50	Wastewater disinfection, degradation of organic pollutants	Govindaraju <i>et al.</i> (2009)
Silver (AgNPs)	<i>Sargassum muticum</i>	Fucoidan, alginate, polyphenols	15–40	Heavy metal removal, antimicrobial remediation	Singh <i>et al.</i> (2016)
Gold (AuNPs)	<i>Sargassum wightii</i>	Fucoidan, alginate, phenolics	8–20	Catalytic degradation of nitroaromatic pollutants	Singaravelu <i>et al.</i> (2007)
Gold (AuNPs)	<i>Turbinaria conoides</i>	Sulfated polysaccharides, proteins	10–35	Catalytic reduction of organic contaminants	Dahoumane <i>et al.</i> (2017)
Iron oxide (Fe ₃ O ₄ NPs)	<i>Spirulina platensis</i>	Phycocyanin, proteins, carbohydrates	20–60	Adsorption of Cr(VI), Pb ²⁺ , Cd ²⁺ and As(V)	El-Naggar <i>et al.</i> (2020)
Iron oxide (Fe ₂ O ₃ NPs)	<i>Chlorella vulgaris</i>	Proteins, chlorophyll, polysaccharides	20–70	Groundwater remediation, heavy metal removal	Singh <i>et al.</i> (2018)
Zinc oxide (ZnO NPs)	<i>Ulva lactuca</i>	Sulfated polysaccharides, proteins	20–70	Photocatalytic degradation of methylene blue and Congo red	El-Naggar <i>et al.</i> (2020)

Zinc oxide (ZnO NPs)	<i>Gracilaria edulis</i>	Agar, proteins, polyphenols	15–60	Textile dye degradation and wastewater treatment	Singh <i>et al.</i> (2018)
Copper oxide (CuO NPs)	<i>Gracilaria corticata</i>	Agar, carrageenan, phenolics	20–60	Heavy metal adsorption, antimicrobial wastewater treatment	El-Naggar <i>et al.</i> (2020)
Titanium dioxide (TiO ₂ NPs)	<i>Chlamydomonas reinhardtii</i>	Proteins, carbohydrates, pigments	10–50	Photocatalytic degradation of pesticides and pharmaceuticals	Dahoumane <i>et al.</i> (2017)
Titanium dioxide (TiO ₂ NPs)	<i>Ulva fasciata</i>	Sulfated polysaccharides, proteins	15–45	Degradation of organic pollutants	Singh <i>et al.</i> (2018)
Selenium (SeNPs)	<i>Chlorella vulgaris</i>	Glutathione, proteins, antioxidants	50–100	Selenium detoxification and heavy metal remediation	El-Naggar <i>et al.</i> (2020)
Palladium (PdNPs)	<i>Sargassum bovinum</i>	Fucoidan, alginate, phenolics	5–20	Catalytic dechlorination of chlorinated compounds	Dahoumane <i>et al.</i> (2017)
Magnesium oxide (MgO NPs)	<i>Ulva lactuca</i>	Sulfated polysaccharides, proteins	20–80	Adsorption of dyes and toxic metal ions	El-Naggar <i>et al.</i> (2020)

References

1. Ahmed, S., Ahmad, M., Swami, B. L., & Ikram, S. (2016). A review on plants extract mediated synthesis of silver nanoparticles for antimicrobial applications. *Journal of Advanced Research*, 7(1), 17–28. <https://doi.org/10.1016/j.jare.2015.02.007>
2. Eker, F., Akdaşçi, E., Duman, H., Bechelany, M., & Karav, S. (2025). Green Synthesis of Silver Nanoparticles Using Plant Extracts: A Comprehensive Review of Physicochemical Properties and Multifunctional Applications. *International Journal of Molecular Sciences*, 26(13), 6222. <https://doi.org/10.3390/ijms26136222>
3. Iravani, S. (2011). Green synthesis of metal nanoparticles using plants. *Green Chemistry*, 13(10), 2638–2650. <https://doi.org/10.1039/c1gc15386b>
4. Mittal, A. K., Chisti, Y., & Banerjee, U. C. (2013). Synthesis of metallic nanoparticles using plant extracts. *Biotechnology Advances*, 31(2), 346–356. <https://doi.org/10.1016/j.biotechadv.2013.01.003>
5. Nasrollahzadeh, M., Sajadi, S. M., Sajjadi, M., & Issaabadi, Z. (2019). Green nanotechnology. *Interface Science and Technology*, 28, 145–198. <https://doi.org/10.1016/B978-0-12-817812-6.00005-0>
6. Rabiee, N. (2025). Green Synthesized Nanomaterials: Principles, Progress, and Perspectives. *ACS Symposium Series*. <https://doi.org/10.1021/bk-2025-1520.ch001>
7. Salam, H. A., et al. (2019). Green synthesis of metal oxide nanoparticles and their environmental applications. *SN Applied Sciences*, 1, 1094.
8. Singh, P., Kim, Y. J., Zhang, D., & Yang, D. C. (2016). Biological synthesis of nanoparticles from plants and microorganisms. *Trends in Analytical Chemistry*, 86, 1–15. <https://doi.org/10.1016/j.trac.2016.08.019>
9. Abdelkhalek, A., Al-Askar, A. A., & Behiry, S. I. (2022). Green synthesized zinc oxide nanocrystals using Pelargonium odoratissimum extract: Characterization and their potent antibacterial, antifungal, and antiviral activities. *Materials*, 15(13), 4445. <https://doi.org/10.3390/ma15134445>
10. Ahmad, A., Mukherjee, P., Senapati, S., Mandal, D., Khan, M. I., Kumar, R., & Sastry, M. (2003). Extracellular biosynthesis of silver nanoparticles using the fungus *Fusarium oxysporum*. *Colloids and Surfaces B: Biointerfaces*,
11. Abdelkhalek, A., Al-Askar, A. A., & Behiry, S. I. (2022). Green synthesized zinc oxide nanocrystals for combating plant pathogenic microbes: Current status and future prospects. *Frontiers in Microbiology*, 13, Article 877478. <https://doi.org/10.3389/fmicb.2022.877478>
12. Ahmad, A., Senapati, S., Khan, M. I., Kumar, R., Ramani, R., Srinivas, V., & Sastry, M. (2003). Intracellular synthesis of gold nanoparticles by a novel alkalotolerant fungus,

- Trichothecium sp. *Nanotechnology*, 14(7), 824–828. <https://doi.org/10.1088/0957-4484/14/7/323>
13. Ahmed, T., Noman, M., Gardner, M., Jiang, C., Tuncsoy, B. S., & Li, B. (2021). Current status and perspective on the application of engineered nanomaterials in wastewater treatment: A comprehensive review. *Journal of Hazardous Materials*, 418, Article 126359. <https://doi.org/10.1016/j.jhazmat.2021.126359>
 14. Alazaiza, M. Y. D., Albahnasawi, A., Ali, G. A. M., Bashir, M. J. K., Coptly, N. K., Amr, S. S. A., & Abushammala, M. F. M. (2021). Application of magnetic nanoparticles for water and wastewater treatment: A review. *Environmental Nanotechnology, Monitoring & Management*, 16, Article 100596. <https://doi.org/10.1016/j.enmm.2021.100596>
 15. Almeida-Naranjo, C. E., Guerrero, V. H., & Villamar-Ayala, C. A. (2023). Activated carbon-based nanomaterials for wastewater treatment: A review. *Nanomaterials*, 13(8), Article 1354. <https://doi.org/10.3390/nano13081354>
 16. Bhandari, G., Baghyalakshmi, K., Kaur, P., & Ganguly, R. (2023). Fungal-mediated synthesis of nanoparticles and their potential applications in environmental remediation: A review. *Environmental Science and Pollution Research*, 30(3), 5275–5295. <https://doi.org/10.1007/s11356-022-24329-8>
 17. Bhushan, D., Sharma, R., & Singla, M. L. (2024). Agricultural waste-derived nanomaterials: Green synthesis and applications in environmental remediation. *Biomass Conversion and Biorefinery*, 14(5), 6121–6140. <https://doi.org/10.1007/s13399-022-03112-4>
 18. Bolade, O. P., Oladipo, A. A., & Bodede, O. (2020). Green synthesis of metal and metal oxide nanoparticles using plant extracts: A review of mechanisms and biomedical applications. *Journal of Molecular Structure*, 1203, Article 127438. <https://doi.org/10.1016/j.molstruc.2019.127438>
 19. Cremonini, E., Zonaro, E., Donini, M., Lampis, S., Boaretti, M., D'Annibale, A., & Vallini, G. (2016). Biogenic selenium nanoparticles: Current status and future prospects in environmental and biomedical fields. *Biotechnology Advances*, 34(8), 1279–1296. <https://doi.org/10.1016/j.biotechadv.2016.09.004>
 20. Dahoumane, S. A., Mechke, C., Wijesekera, D., & Filipe, C. D. M. (2017). Algae-mediated biosynthesis of inorganic nanomaterials: A review. *Bioresource Technology*, 224, 1–16. <https://doi.org/10.1016/j.biortech.2016.10.038>
 21. Deplanche, K., Caldelari, I., Mikheenko, I. P., Sargent, F., & Macaskie, L. E. (2011). Involvement of hydrogenases in axial and peripheral palladium nanoparticle deposition on *Desulfovibrio desulfuricans*. *Microbiology*, 157(Pt 11), 3244–3253. <https://doi.org/10.1099/mic.0.052670-0>

22. El-Naggar, N. E. A., Hussein, M. H., & El-Sawah, A. A. (2020). Phycobiliprotein-mediated green synthesis of metal and metal oxide nanoparticles and their potential applications: An eco-friendly approach. *International Journal of Biological Macromolecules*, 165, 3121–3139. <https://doi.org/10.1016/j.ijbiomac.2020.10.180>
23. Fei, X., Wang, Y., Lu, J., & Zhao, C. (2022). Application of carbon-based nanomaterials and cyclodextrin nanosponges in the degradation of persistent organic pollutants: A review. *Chemosphere*, 307, Article 135841. <https://doi.org/10.1016/j.chemosphere.2022.135841>
24. Ferdush, N., Islam, M. T., & Uddin, M. E. (2025). Microbial fabrication of functional nanomaterials: Mechanisms, challenges, and environmental remediation perspectives. *Environmental Technology & Innovation*, 37, Article 103982. <https://doi.org/10.1016/j.eti.2024.103982>
25. Fouda, A., Hassan, S. E. D., Eid, A. M., & Ewais, E. E. D. (2018). Biotechnological applications of metal and metal oxide nanoparticles synthesized by fungal and bacterial endophytes: A comprehensive review. *Journal of Advanced Research*, 13, 61–74. <https://doi.org/10.1016/j.jare.2018.05.012>
26. Gobinath, R., Rajeshkumar, S., & Santhoshkumar, J. (2021). Mechanisms of heavy metal and dye adsorption using green synthesized nanomaterials: A review. *Water, Air, & Soil Pollution*, 232(5), Article 201. <https://doi.org/10.1007/s11270-021-05151-2>
27. Govindaraju, K., Basha, S. K., Kumar, V. G., & Singaravelu, G. (2009). Silver, gold and bimetallic nanoparticles production using single-cell protein (*Spirulina platensis*) Geitler. *Journal of Materials Science*, 44(4), 976–985. <https://doi.org/10.1007/s10853-008-3193-4>
28. Guilger-Casagrande, M., & Lima, R. (2019). Synthesis of silver nanoparticles mediated by fungi: A review. *Frontiers in Bioengineering and Biotechnology*, 7, Article 287. <https://doi.org/10.3389/fbioe.2019.00287>
29. He, S., Guo, Z., Zhang, Y., Yan, S., Dahl, J., & Gu, N. (2007). Biosynthesis of gold nanoparticles using the bacterium *Rhodospseudomonas capsulata*. *Materials Letters*, 61(18), 3984–3987. <https://doi.org/10.1016/j.matlet.2007.01.018>
30. Heinemann, M. G. (2021). Plant extract-mediated synthesis of metal nanoparticles: Nucleation kinetics and biomolecular capping dynamics. *Nanotechnology Reviews*, 10(1), 841–858. <https://doi.org/10.1515/ntrev-2021-0060>
31. Huang, X., Zhang, K., & Liu, Y. (2024). Magnetic green nanoparticles in environmental remediation: Facile magnetic recovery, regeneration, and lifecycle assessment. *Journal of Cleaner Production*, 434, Article 140212. <https://doi.org/10.1016/j.jclepro.2023.140212>
32. Kahru, A., & Dubourguier, H.-C. (2010). From ecotoxicology to nanoecotoxicology. *Toxicology*, 269(2–3), 105–119. <https://doi.org/10.1016/j.tox.2009.08.016>

33. Kalishwaralal, K., Deepak, V., Ram Kumar Pandian, S. B., Kottaisamy, M., BarathManiKanth, S., Kartikeyan, B., & Gurunathan, S. (2008). Biosynthesis of silver and gold nanoparticles using *Brevibacterium casei*. *Colloids and Surfaces B: Biointerfaces*, 77(2), 257–262. <https://doi.org/10.1016/j.colsurfb.2010.02.007>
34. Karn, B., Kuiken, T., & Otto, M. (2009). Nanotechnology and in situ remediation: A review of the benefits and potential risks. *Environmental Health Perspectives*, 117(12), 1823–1831. <https://doi.org/10.1289/ehp.0900793>
35. Kaur, J., Sharma, M., & Singh, R. (2023). Environmental aging and stability of green nanoparticles in complex aquatic matrices: A review of challenges and field applicability. *Science of The Total Environment*, 862, Article 160821. <https://doi.org/10.1016/j.scitotenv.2022.160821>
36. Klaus, T., Joerger, R., Olsson, E., & Granqvist, C. G. (1999). Silver-based crystalline nanoparticles, microbially fabricated. *Proceedings of the National Academy of Sciences*, 96(24), 13611–13614. <https://doi.org/10.1073/pnas.96.24.13611>
37. Konishi, Y., Tsukiyama, T., Ohno, K., Saitoh, N., Nomura, T., & Nagamine, S. (2007). Intracellular recovery of platinum by *Shewanella* algae. *Hydrometallurgy*, 85(2–4), 188–193. <https://doi.org/10.1016/j.hydromet.2006.08.013>
38. Korlam, V. R., Kumar, P., & Reddy, S. (2025). Anthropogenic contamination of agro-ecosystems: Sources, pathways, and global management strategies. *Environmental Pollution*, 340, Article 122841. <https://doi.org/10.1016/j.envpol.2024.122841>
39. Lovley, D. R., Holmes, D. E., & Nevin, K. P. (2004). Dissimilatory Fe(III) and Mn(IV) reduction. *Advances in Microbial Physiology*, 49, 219–286. [https://doi.org/10.1016/S0065-2911\(04\)49005-5](https://doi.org/10.1016/S0065-2911(04)49005-5)
40. Lu, N., Chen, Z., Tang, H., & Zhou, W. (2025). Surface-engineered magnetic magnetite nanoparticles for efficient aqueous heavy metal sequestration: Interfacial mechanism and performance. *Chemical Engineering Journal*, 482, Article 148902. <https://doi.org/10.1016/j.cej.2024.148902>
41. Mehta, P., Sharma, A., & Verma, K. (2024). Surface area kinetics and adsorption mechanisms of green metal oxide nanoparticles in toxic dye removal. *Journal of Water Process Engineering*, 57, Article 104672. <https://doi.org/10.1016/j.jwpe.2023.104672>
42. Moond, M., Kumar, D., & Rani, P. (2024). Catalytic and photocatalytic degradation of industrial organic pollutants by metallic and bimetallic nanoparticles: A critical review. *Environmental Research*, 241, Article 117621. <https://doi.org/10.1016/j.envres.2023.117621>

43. Mughal, B., Zaidi, S. Z. J., Zhang, X., & Hassan, S. U. (2021). Biogenic nanoparticles: Synthesis, mechanism of action, and environmental applications. *Frontiers in Nanotechnology*, 3, Article 776144. <https://doi.org/10.3389/fnano.2021.776144>
44. Mukherjee, P., Ahmad, A., Mandal, D., Senapati, S., Sainkar, S. R., Khan, M. I., Parishcha, R., Ajaykumar, P. V., Alam, M., Kumar, R., & Sastry, M. (2001). Fungus-mediated synthesis of silver nanoparticles and their characterization. *Nano Letters*, 1(10), 515–519. <https://doi.org/10.1021/nl0155274>
45. Mukherjee, P., Senapati, S., Mandal, D., Ahmad, A., Khan, M. I., Kumar, R., & Sastry, M. (2002). Extracellular synthesis of gold nanoparticles by the fungus *Fusarium oxysporum*. *ChemBioChem*, 3(5), 461–463. [https://doi.org/10.1002/1439-7633\(20020503\)3:5<461::AID-CBIC461>3.0.CO;2-X](https://doi.org/10.1002/1439-7633(20020503)3:5<461::AID-CBIC461>3.0.CO;2-X)
46. Mukhopadhyay, S., Mukherjee, S., & Bhowmick, S. (2021). Zero-valent iron nanoparticles (nZVI) for environmental remediation: Advances in surface functionalization, stabilization, and reactive barriers. *Journal of Environmental Management*, 281, Article 111867. <https://doi.org/10.1016/j.jenvman.2020.111867>
47. Narayanan, K. B., & Sakthivel, N. (2010). Biological synthesis of metal nanoparticles by microbes. *Advances in Colloid and Interface Science*, 156(1–2), 1–13. <https://doi.org/10.1016/j.cis.2010.02.001>
48. Nawaz, R., Ali, A., & Khan, M. (2024). Visible light responsive TiO₂ black nanoparticles for advanced photocatalytic degradation of hazardous industrial dyes: Reactive oxygen species dynamics. *Solar Energy*, 269, Article 112349. <https://doi.org/10.1016/j.solener.2024.112349>
49. Özkara, A., Akyıl, D., & Konuk, M. (2016). Pesticides, environmental pollution, and health, the science of pesticides. In M. L. Larramendy & S. Soloneski (Eds.), *Environmental health risk: Hazardous factors to living species* (pp. 3–28). IntechOpen. <https://doi.org/10.5772/63094>
50. Patil, S., Chandrasekaran, R., & Vithanage, M. (2020). Green synthesis of nanoparticles: Sustainable methodologies for contaminant remediation. *Current Opinion in Environmental Science & Health*, 16, 22–30. <https://doi.org/10.1016/j.coesh.2020.02.003>
51. Presentato, A., Cappelletti, M., Sansone, A., Ferreri, C., & Turner, R. J. (2018). Bacterial production of selenium nanoparticles and their potential biotechnological applications. *Microbial Biotechnology*, 11(6), 1084–1098. <https://doi.org/10.1111/1751-7915.13289>
52. Raj, S., Sharma, M., & Das, S. (2024). Graphene-based nanostructures as redox mediators for enhanced biodegradation of xenobiotic pollutants. *Bioresource Technology Reports*, 25, Article 101742. <https://doi.org/10.1016/j.biteb.2023.101742>

53. Rathod, S., Pathak, A., & Joshi, P. (2024). Green nanosensors: Recent trends and applications in environmental pollution monitoring. *Trends in Environmental Analytical Chemistry*, 41, Article e00223. <https://doi.org/10.1016/j.teac.2023.e00223>
54. Roopa, M. C., Prasad, N., & Kumar, H. (2025). Solar-driven photocatalytic detoxification of wastewater using green-synthesized semiconductors: A roadmap from laboratory to decentralized field installations. *Journal of Cleaner Production*, 442, Article 141029. <https://doi.org/10.1016/j.jclepro.2024.141029>
55. Roychoudhury, P., Ghosh, S., & Webster, T. J. (2016). Algal synthesis of silver nanoparticles and their antibacterial activity against multi-drug resistant pathogens. *International Journal of Nanomedicine*, 11, 5233–5244. <https://doi.org/10.2147/IJN.S114751>
56. Salam, H. A., Rajiv, P., Kamaraj, M., Jagadeeswaran, P., Gunalan, S., & Sivaraj, R. (2019). Green synthesis of metal oxide nanoparticles and their environmental applications. *SN Applied Sciences*, 1(9), Article 1094. <https://doi.org/10.1007/s42452-019-1126-7>
57. Serov, D. A., Belyi, M. D., & Burmistrov, D. E. (2024). Photocatalytic and antimicrobial properties of metal oxide nanoparticles in water purification: Current status and mechanisms. *Nanomaterials*, 14(4), Article 341. <https://doi.org/10.3390/nano14040341>
58. Singaravelu, G., Arockiamary, J. S., Kumar, V. G., & Govindaraju, K. (2007). A novel extracellular synthesis of monodisperse gold nanoparticles using marine alga, *Sargassum wightii* Greville. *Colloids and Surfaces B: Biointerfaces*, 57(1), 97–101. <https://doi.org/10.1016/j.colsurfb.2007.01.010>
59. Singh, J., Dutta, T., Kim, K. H., Rawat, M., Samddar, P., & Kumar, P. (2018). ‘Green’ synthesis of metals and their oxide nanoparticles: Applications for environmental remediation. *Journal of Nanobiotechnology*, 16(1), Article 84. <https://doi.org/10.1186/s12951-018-0408-3>
60. Singh, R. (2025). Safe-by-design frameworks and ethical dilemmas in environmental nanotechnology deployment. *Environmental Science & Policy*, 153, Article 103681. <https://doi.org/10.1016/j.envsci.2024.103681>
61. Singh, R., Shedbalkar, U. U., Wadhwani, S. A., & Chopade, B. A. (2017). Bacteriogenic metallic nanoparticles: Synthesis, properties, and applications. In S. K. Ghosh & T. Webster (Eds.), *Nanobiotechnology: Human Health and the Environment* (pp. 111–138). CRC Press. <https://doi.org/10.1201/9781315153285-5>
62. Su, C., Jiang, L., & Zhang, W. (2014). A review on heavy metal contamination in the soil worldwide: Situation, impact and remediation techniques. *Environmental Skeptics and Critics*, 3(2), 24–38.

63. Tiwari, A., Patel, N. K., & Sharma, V. (2026). Artificial intelligence and machine learning in predictive nano-engineering: Optimizing green synthesis and remediation kinetics. *Computers & Chemical Engineering*, 181, Article 108529. <https://doi.org/10.1016/j.compchemeng.2025.108529>
64. Tratnyek, P. G., & Johnson, R. L. (2006). Nanotechnologies for environmental cleanup. *Nano Today*, 1(2), 44–48. [https://doi.org/10.1016/S1748-0132\(06\)70048-2](https://doi.org/10.1016/S1748-0132(06)70048-2)
65. Treviño, P., Rodríguez, M., & García, E. (2014). Electrochemical reduction of hexavalent chromium in industrial effluents using zero-valent aluminium nanoparticles. *Journal of Applied Electrochemistry*, 44(8), 915–923. <https://doi.org/10.1007/s10800-014-0708-4>
66. Ugwuanyi, E. D., Emenike, P. C., & Omole, D. O. (2024). Engineered and green nanomaterials in managing emerging contaminants in water: Removal mechanisms and lifecycle prospects. *Environmental Pollution*, 345, Article 123491. <https://doi.org/10.1016/j.envpol.2024.123491>
67. Vázquez-Núñez, O., Molina-Guerrero, C. E., Peña-Castro, J. M., Fernández-Luqueño, F., & de la Rosa-Álvarez, M. G. (2020). Use of carbon nanotubes for the removal of toxic organic compounds and heavy metals from wastewater: A review. *Environmental Technology & Innovation*, 18, Article 100762. <https://doi.org/10.1016/j.eti.2020.100762>
68. Vigneshwaran, N., Ashtaputre, N. M., Varadarajan, P. V., Nachane, R. P., Paralikar, K. M., & Balasubramanya, R. H. (2007). Biological synthesis of silver nanoparticles using the fungus *Aspergillus flavus*. *Materials Letters*, 61(6), 1413–1418. <https://doi.org/10.1016/j.matlet.2006.07.042>
69. Vimal, A., Kumar, A., & Singh, R. K. (2026). Functionalized silica nanoparticles (SiO₂ NPs) for catalytic soil and groundwater remediation: Operational mechanisms and eco-safety. *Journal of Hazardous Materials Advances*, 13, Article 100412. <https://doi.org/10.1016/j.hazadv.2025.100412>

HUMAN MICROBIOME: EMERGING ROLES IN HEALTH AND DISEASE

Poonguzhali K. and Arulmozhi R.*

PG and Research Department of Microbiology,
Sengamala Thayaar Educational Trust Women's College (Autonomous),
Sundarakkottai, Mannargudi, Thiruvavur (dt) - 614 016, Tamil Nadu

*Corresponding author E-mail: amozhi72@gmail.com

Abstract

The human microbiome is a dynamic community of bacteria, archaea, fungi, viruses and other microorganisms that inhabit distinct anatomical sites and interact continuously with host tissues. Modern sequencing and multi-omics approaches have shifted the view of microorganisms from isolated pathogens to components of a complex biological ecosystem that contributes to metabolism, epithelial barrier function, immune development, colonization resistance and host signalling. Although the gut microbiome has received substantial attention, microbial communities at the oral cavity, skin, respiratory tract and reproductive tract also participate in health and disease. Alterations in microbial composition or function, commonly described as dysbiosis, have been associated with gastrointestinal disorders, metabolic disease, inflammatory conditions, cancer and neurological disorders. However, association does not necessarily establish causation, and a universal definition of a 'healthy' microbiome remains difficult because microbiome structure varies with age, geography, diet, medications and other host and environmental factors. Emerging interventions include dietary modulation, probiotics, prebiotics, synbiotics, postbiotics, fecal microbiota transplantation and more targeted live biotherapeutic approaches. This chapter summarizes the organization and functions of the human microbiome, its emerging roles in health and disease, major determinants of microbiome variation, current therapeutic strategies and important directions for future research and clinical translation.

Keywords: Human Microbiome, Gut Microbiota, Dysbiosis, Host–Microbe Interaction, Immune System, Microbiome Therapeutics, Health and Disease.

Introduction

The human body is inhabited by complex microbial communities that occupy different ecological niches, including the gastrointestinal tract, oral cavity, skin, respiratory tract and urogenital tract. The term microbiota refers to the microorganisms present in a particular habitat, whereas microbiome is commonly used to describe the microbial community together with its genetic and functional potential. The Human Microbiome Project helped establish a systematic framework for characterizing microbial communities across body sites and highlighted the substantial diversity of microorganisms associated with healthy humans (Turnbaugh *et al.*, 2007).

Microbiome research has subsequently moved beyond cataloguing microbial species toward understanding microbial functions and interactions with the host. Microorganisms contribute to the digestion and transformation of dietary components, production of metabolites, maintenance of epithelial barriers, regulation of immune responses and protection against invading organisms. These functions arise from interactions among many microbial species rather than from a single organism acting in isolation (Flint *et al.*, 2012; Ma *et al.*, 2024).

The concept of a healthy microbiome is increasingly understood as a range of functional states rather than a fixed list of organisms. Microbial communities vary between individuals and can change over time in response to age, diet, geography, lifestyle, disease and medication exposure. Consequently, microbiome research increasingly emphasizes ecological stability, functional capacity and host context rather than simply microbial diversity (Kunath *et al.*, 2024).

Table 1: Human Phenotype Microbiome

Body Site	Major Ecological Features	Important Functions	Examples of Disease Links
Gut	High density and metabolic diversity	Fermentation, metabolite production, barrier and immune regulation	Inflammatory bowel disease, metabolic disorders, colorectal cancer
Oral cavity	Biofilms on teeth, tongue and mucosa	Colonization resistance, nitrate metabolism and local immune interactions	Dental caries, periodontal disease and systemic associations
Skin	Site-specific communities shaped by moisture and sebum	Barrier support, antimicrobial activity and immune signalling	Acne, atopic dermatitis and other inflammatory skin disorders
Respiratory tract	Low biomass compared with gut and site-specific communities	Immune education and interaction with inhaled/environmental microbes	Respiratory infection and inflammatory conditions
Vaginal tract	Community structure influenced by pH and host hormones	Maintenance of local environment and pathogen resistance	Bacterial vaginosis and reproductive health conditions

Composition and Distribution of the Human Microbiome

Microbial communities are distributed unevenly throughout the human body. The colon contains one of the most densely populated microbial ecosystems, while the oral cavity contains diverse communities adapted to surfaces such as the tongue, teeth and gingiva. The skin microbiome varies according to moisture, temperature, sebaceous activity and anatomical location. Microorganisms are also present in the upper respiratory and female reproductive tracts, where local environmental conditions shape community composition (Chaudhary *et al.*, 2024). At the

taxonomic level, bacteria are the best characterized component of the human microbiome, but bacteriophages and other viruses, archaea and fungi are also important. Importantly, the presence of a microorganism does not by itself indicate whether it is beneficial or harmful. The effect of a microbe can depend on microbial community structure, available nutrients, host immunity, epithelial integrity and environmental conditions.

Microbial communities can therefore be considered ecological systems. Species compete for nutrients and attachment sites, exchange metabolites and communicate through chemical signals. Some organisms produce compounds that inhibit competitors or pathogens, whereas others transform dietary or host-derived substrates into metabolites that influence distant organs. These interactions contribute to the overall functional phenotype of the microbiome.

Roles of the Microbiome in Human Health

Metabolic Functions

One of the most established roles of the gut microbiome is the transformation of dietary components that escape digestion in the upper gastrointestinal tract. Microbial fermentation of complex carbohydrates generates metabolites, including short-chain fatty acids (SCFAs), that can influence epithelial function, energy metabolism and immune signalling. The microbiome also participates in the transformation of bile acids and other host-derived compounds (Flint *et al.*, 2012). Microorganisms contribute to the bioavailability and metabolism of several nutrients and can synthesize or transform vitamins and other small molecules. These activities mean that host nutrition is influenced not only by the nutrient content of food but also by the metabolic capabilities of the microbial community.

Intestinal Barrier and Colonization Resistance

The intestinal epithelium forms a critical interface between the host and the external environment. Microbial metabolites and microbe–host interactions can support epithelial integrity and influence mucus production and barrier function. At the same time, resident microbial communities can limit pathogen establishment through competition for nutrients and ecological niches, production of inhibitory molecules and stimulation of host defence mechanisms. Colonization resistance is therefore an important component of microbiome-mediated protection. Disruption of microbial communities by factors such as antibiotics can reduce this ecological resistance and create conditions in which opportunistic pathogens can expand. The clinical relevance of this concept is particularly clear in recurrent *Clostridioides difficile* infection, where microbiota restoration has an established therapeutic role (Lee *et al.*, 2024).

Immune Development and Regulation

The microbiome interacts continuously with innate and adaptive immune pathways. Microbial products and metabolites can influence antigen-presenting cells, epithelial signalling and

lymphocyte responses. Conversely, host immune responses shape microbial community composition and help maintain separation between microorganisms and host tissues. This bidirectional relationship is central to intestinal homeostasis (Iebba *et al.*, 2012).

Disruption of this relationship may contribute to inappropriate inflammatory responses. Nevertheless, microbiome–immune associations should be interpreted carefully because the direction of causality may vary by disease and because host inflammation can itself alter microbial ecology.

Gut–Brain and Other Organ Axes

Microbial metabolites, immune mediators, endocrine signals and neural pathways provide routes through which the gut microbiome can influence organs beyond the intestine. The gut–brain axis has attracted particular interest because microbial activity can affect metabolites and signalling pathways relevant to nervous-system function. Similarly, gut–liver, gut–immune and oral–gut interactions illustrate how microbiomes at different anatomical sites may communicate indirectly or directly (Kunath *et al.*, 2024; Chaudhary *et al.*, 2024).

These axes are biologically plausible, but the strength of evidence differs among diseases and mechanisms. Current research increasingly combines longitudinal human studies, mechanistic experiments and intervention trials to distinguish correlation from causal effects.

Microbiome in Disease

Gastrointestinal Diseases

Changes in the intestinal microbiome have been described in inflammatory bowel disease, irritable bowel syndrome and other gastrointestinal disorders. Reported alterations include changes in microbial abundance, community structure and metabolite production. These observations have encouraged investigation of dietary interventions, probiotics, prebiotics and microbiome restoration approaches (Noguera-Fernández *et al.*, 2024).

Fecal microbiota transplantation (FMT) has demonstrated the clearest clinical efficacy in recurrent *C. difficile* infection. Evidence for other gastrointestinal conditions is developing, but outcomes vary according to disease, donor characteristics, preparation, route of administration and study design. Standardization and long-term safety remain important research priorities (Lee *et al.*, 2024; Allegretti *et al.*, 2024).

Metabolic Disorders

Microbiome alterations have been associated with obesity, insulin resistance, type 2 diabetes and metabolic dysfunction-associated liver disease. Potential mechanisms include changes in energy harvest, SCFA production, bile-acid signalling, intestinal permeability and low-grade inflammation. However, human metabolic phenotypes are influenced by diet, physical activity, medications, genetics and socioeconomic factors, making it difficult to attribute disease risk to the microbiome alone (Hu & Shen, 2024).

Intervention studies are therefore important. Research on microbiome modulation includes dietary fibre, prebiotics, probiotics, synbiotics and FMT. Results are heterogeneous, and treatment effects may depend on baseline microbiome configuration and host characteristics.

Cancer

Microorganisms can influence carcinogenesis through effects on inflammation, epithelial integrity, metabolite production and immune responses. The microbiome is also being studied in relation to response and adverse effects during cancer immunotherapy. These findings have raised the possibility of using microbial features as biomarkers or therapeutic targets, although clinical application requires careful validation.

Neurological and Psychiatric Conditions

Associations between the gut microbiome and neurological or psychiatric conditions have generated substantial interest. Proposed pathways include microbial metabolites, immune signalling, vagal and endocrine communication, and modulation of the gut barrier. Current evidence supports biological interaction but does not justify treating every microbiome difference as a direct cause of neurological disease. Well-controlled longitudinal and intervention studies remain necessary.

Factors That Shape the Human Microbiome

Microbiome composition is influenced by a combination of host, environmental and behavioural factors. Early-life exposures, mode of delivery, infant feeding, age, diet, geography, sanitation, occupation, stress, physical activity and medication use can all contribute to microbial variation. Antibiotics are especially important because they can produce substantial and sometimes prolonged changes in community structure.

Diet is a major modifiable determinant. Long-term dietary patterns influence substrate availability for microbial metabolism, while short-term dietary changes can also alter microbial composition and function. Dietary fibre and diverse plant-derived substrates can support fermentation pathways, but responses are individualized because different microbiomes have different metabolic capacities (Flint *et al.*, 2012).

These sources of variation complicate the definition of a universal 'healthy microbiome'. A microbial configuration associated with health in one population may not be identical to that observed in another. Consequently, microbiome research increasingly emphasizes functional characteristics and ecological resilience rather than a single ideal community.

Methods Used to Study the Human Microbiome

Culture-independent sequencing has transformed microbiome research. Amplicon sequencing, particularly 16S rRNA gene profiling for bacteria, is useful for describing community composition, whereas shotgun metagenomic sequencing provides broader information about

microbial genes and species. Metatranscriptomics, metabolomics and proteomics can provide complementary information about microbial activity and host responses.

However, microbiome measurements are sensitive to sampling site, collection method, storage, DNA extraction, sequencing platform and bioinformatic processing. Best-practice approaches therefore require careful experimental design, appropriate controls and transparent reporting. Functional validation is particularly important when associations are interpreted as potential mechanisms.

Emerging Microbiome-Based Therapeutics

Probiotics, Prebiotics and Synbiotics

Probiotics are live microorganisms administered with the intention of providing a health benefit, while prebiotics are substrates selectively utilized by host microorganisms to confer a health benefit. Synbiotics combine probiotic organisms with suitable substrates. Their effects are strain, formulation- and condition-dependent, so results from one product cannot automatically be generalized to all products.

Clinical studies have investigated these approaches across gastrointestinal and other conditions. Evidence varies by disease and intervention, and treatment should be evaluated according to defined clinical outcomes rather than changes in microbial diversity alone (Noguera-Fernández *et al.*, 2024).

Fecal Microbiota Transplantation

FMT transfers processed donor-derived intestinal microbiota to a recipient with the aim of restoring microbial functions. Its strongest established indication is recurrent *C. difficile* infection. For other disorders, clinical evidence is still developing and varies considerably. Key issues include donor screening, pathogen transmission, manufacturing consistency, dosing, route of administration and long-term monitoring (Lee *et al.*, 2024).

Live Biotherapeutic Products and Precision Microbiome Therapies

Next-generation approaches seek to replace broad community transfer with defined microbial consortia or live biotherapeutic products. Other emerging strategies include engineered microorganisms, targeted phage approaches, metabolite-based interventions and personalized dietary modulation. These approaches may allow more precise control over microbial functions while reducing some uncertainties associated with whole-community transfer (Allegretti *et al.*, 2024; Kim *et al.*, 2024).

Conclusion

The human microbiome is an integral component of human biology rather than a collection of microorganisms that merely coexist with the host. Its members participate in metabolism, immune regulation, barrier maintenance, colonization resistance and inter-organ signalling. Microbiome alterations are associated with a broad range of diseases, but the relationship

between microbial change and disease is often complex and bidirectional. Current evidence supports established clinical applications in selected settings, particularly recurrent *C. difficile* infection, while many other microbiome-based interventions remain under active investigation. Future progress will depend on combining ecological understanding with rigorous clinical research. Defining microbial functions, identifying causal mechanisms and developing safe, standardized and personalized interventions may enable the microbiome to become an increasingly useful component of preventive medicine, diagnostics and therapeutics.

References

1. Allegretti, J. R. (2024). The progression of microbiome therapeutics for the management of gastrointestinal diseases and beyond. *Gastroenterology*, 167(4), 1025–1043.
2. Chaudhary, P. P., Kaur, M., & Myles, I. A. (2024). Does “all disease begin in the gut”? The gut-organ cross talk in the microbiome. *Applied Microbiology and Biotechnology*, 108, 1–16.
3. Flint, H. J., Scott, K. P., Louis, P., & Duncan, S. H. (2012). The role of the gut microbiota in nutrition and health. *Nature Reviews Gastroenterology & Hepatology*, 9(10), 577–589.
4. Hu, C., & Shen, H. (2024). Microbes in health and disease: Human gut microbiota. *Applied Sciences*, 14(23), 11354.
5. Iebba, V., Nicoletti, M., & Schippa, S. (2012). Gut microbiota and the immune system: An intimate partnership in health and disease. *International Journal of Immunopathology and Pharmacology*, 25(4), 823–833.
6. Kim, D.-Y., Lee, S.-Y., Lee, J.-Y., Whon, T. W., Lee, J.-Y., Jeon, C. O., & Bae, J.-W. (2024). Gut microbiome therapy: Fecal microbiota transplantation vs live biotherapeutic products. *Gut Microbes*, 16(1), 2412376.
7. Kunath, B. J., De Rudder, C., Laczny, C. C., Letellier, E., & Wilmes, P. (2024). The oral gut microbiome axis in health and disease. *Nature Reviews Microbiology*, 22, 791–805.
8. Lee, J. Y., Kim, Y., Kim, J., & Kim, J. K. (2024). Fecal microbiota transplantation: Indications, methods, and challenges. *Journal of Microbiology*, 62(12), 1057–1074.
9. Lynch, S. V., & Pedersen, O. (2016). The human intestinal microbiome in health and disease. *New England Journal of Medicine*, 375(24), 2369–2379.
10. Ma, Z., Zuo, T., Frey, N., et al. (2024). A systematic framework for understanding the microbiome in human health and disease: From basic principles to clinical translation. *Signal Transduction and Targeted Therapy*, 9, 237.

HAIR AND SCALP DISEASES: MEDICAL, SURGICAL AND COSMETIC TREATMENT

M. Kannahi* and R. Sasireka

PG and Research Department of Microbiology,
STET Women's College (Autonomous), Sundarakkottai-614016
(Affiliated to Bharathidasan University, Tiruchirappalli)
Mannargudi, Thiruvarur, Tamil Nadu, India.

*Corresponding author E-mail: kannahiamf@gmail.com

Abstract

Hair and scalp diseases are common disorders that can affect the hair shaft, hair follicle, sebaceous glands, and scalp skin. They range from relatively mild conditions such as dandruff and seborrheic dermatitis to infectious diseases, inflammatory disorders, alopecia, and scarring conditions. Hair disorders may result from genetic factors, hormonal changes, nutritional deficiencies, infections, autoimmune mechanisms, medications, systemic diseases, or mechanical and cosmetic practices. Accurate diagnosis requires a detailed history, clinical examination, scalp and hair assessment, dermoscopy or trichoscopy, microscopic examination, fungal culture, laboratory investigations, and occasionally scalp biopsy. *Trichophyton* and *Microsporum* species are important causes of tinea capitis, while *Malassezia* species are associated with seborrheic dermatitis and dandruff.

Management depends on the underlying cause. Medical treatment includes antifungal, anti-inflammatory, antimicrobial and hair-growth therapies where clinically appropriate. Surgical procedures such as hair transplantation may be considered for selected stable forms of permanent hair loss. Cosmetic management includes appropriate hair cleansing, conditioning, camouflage techniques and careful selection of hair-care products. Early recognition and appropriate management are particularly important in infections and scarring alopecia because delayed treatment can sometimes result in permanent hair loss.

Keywords: Hair Diseases, Scalp Diseases, Alopecia, Dandruff, Tinea Capitis, Medical Treatment, Surgical Treatment, Cosmetic Treatment.

1. Introduction

Hair is a keratinized structure that develops from hair follicles embedded within the skin. In addition to its protective role, hair contributes to thermoregulation, sensory perception and physical appearance. The scalp contains numerous hair follicles and sebaceous glands, making it particularly susceptible to disorders involving keratinization, inflammation, infection and abnormal hair growth. Hair and scalp diseases may occur at any age and can produce symptoms

such as itching, scaling, redness, tenderness, excessive shedding, broken hairs and localized or diffuse hair loss. Hair loss is medically known as alopecia and can be classified broadly into nonscarring and scarring forms. In nonscarring alopecia, the follicular structures remain potentially capable of producing hair, whereas scarring alopecia involves destruction of follicles and may result in permanent loss. Common forms include androgenetic alopecia, alopecia areata and telogen effluvium. Scarring disorders include conditions such as lichen planopilaris frontal fibrosing alopecia and discoid lupus erythematosus.

Scalp infections are another important group of diseases. Tinea capitis is a dermatophyte infection affecting the scalp and hair, particularly in children. Important causative organisms include *Trichophyton tonsurans*, *Trichophyton violaceum* and *Microsporum canis*. The clinical presentation can vary from mild scaling to inflammatory lesions, broken hairs, alopecia and kerion formation.

2. Anatomy and Physiology of Hair and Scalp

The scalp consists of skin, connective tissue, the Epicranial aponeurosis, loose connective tissue and pericranium. Hair follicles extend from the epidermis into the dermis and, in some locations, the subcutaneous tissue. Each follicle passes through a continuous cycle of growth and regression. The hair cycle is traditionally divided into three major phases: anagen, catagen and telogen. Anagen is the active growth phase during which matrix cells proliferate and produce the hair shaft. Catagen is a relatively short transitional phase involving regression of the lower follicle. Telogen is the resting phase, after which the old hair is shed and a new cycle begins.

Disturbances in this cycle can result in excessive shedding or reduced hair density. Sebaceous glands associated with hair follicles produce sebum, which contributes to lubrication of the hair and scalp. Excessive or altered sebum production can contribute to certain scalp disorders. The scalp also possesses a complex microbial community.

Organisms including *Malassezia* species normally inhabit sebaceous areas, although changes in host and environmental factors may be associated with inflammatory scalp conditions.

3. Classification of Hair Scalp Diseases

Hair and scalp diseases can be classified according to their cause and clinical presentation (Messenger *et al.*, 2023).

3.1 Infectious Diseases

Important infectious conditions include tinea capitis, bacterial folliculitis and pediculosis capitis. Fungal infections of the scalp are particularly important because they may produce scaling and hair loss.

3.2 Inflammatory Diseases

Seborrheic dermatitis, psoriasis and eczema may involve the scalp. These disorders commonly produce scaling, itching, erythema and irritation.

3.3 Hair-loss Disorders

Hair-loss disorders include androgenetic alopecia, alopecia areata, telogen effluvium and anagen effluvium. They differ in their pattern, duration, underlying mechanism and prognosis.

3.4 Scarring Alopecia

Scarring alopecia are inflammatory disorders that can permanently damage hair follicles. Examples include lichen planopilaris frontal fibrosing alopecia and discoid lupus erythematosus.

3.5 Hair-Shaft Disorders

Hair-shaft disorders alter the structure and strength of hair. Excessive heat, chemical treatments, mechanical trauma and inherited abnormalities can result in hair breakage and altered hair texture.

4. Dandruff and Seborrheic Dermatitis

Dandruff is a common scalp condition characterized mainly by visible flakes without significant inflammation. Seborrheic dermatitis is a more inflammatory disorder involving erythema, scaling and sometimes itching. Sebaceous areas such as the scalp, eyebrows and areas around the nose are frequently affected.

The pathogenesis is multifactorial and involves sebum, host susceptibility, epidermal barrier function and the scalp microbiome. *Malassezia* species are commonly associated with seborrheic dermatitis, although their presence alone does not establish disease.

Management generally involves regular scalp cleansing and medicated shampoos selected according to the severity of symptoms. Active ingredients used in dandruff management can include antifungal agents and keratolytic or anti-inflammatory preparations. Persistent or severe scalp scaling should be assessed by a dermatologist because psoriasis, eczema and fungal infection can sometimes resemble dandruff.




Figure 1: Dandruff and Hair loss

Tinea Capitis

Tinea capitis is a dermatophyte infection of the scalp and hair shaft. It occurs predominantly in children but can also affect adults. Important organisms include *Trichophyton tonsurans*,

Trichophyton violaceum and *Microsporum canis*. The epidemiology of causative organisms differs according to geographical location (Alkeswani *et al.*, 2019; Mayser *et al.*, 2020).

Clinical manifestations include scalp scaling, itching, broken hairs and localized alopecia. Inflammatory forms can produce painful swelling and pustular lesions. Kerion is a highly inflammatory presentation that may result in scarring if treatment is delayed.

Diagnosis may involve direct microscopic examination using potassium hydroxide preparation, fungal culture and, where available, molecular testing. Trichoscopy can provide additional information about characteristic hair and scalp findings.

Treatment requires systemic antifungal therapy because topical treatment alone generally does not adequately penetrate infected hair shafts. Depending on the causative organism, age and clinical circumstances, medicines such as griseofulvin, terbinafine, itraconazole or fluconazole may be selected by a physician. Adjunctive antifungal shampoos may help reduce fungal spores and transmission but should not be regarded as a replacement for appropriate systemic therapy.

5. Bacterial Infections of the Scalp

Bacterial infections may involve hair follicles and surrounding skin. Folliculitis is characterized by inflammation around hair follicles and may produce small papules or pustules. Common bacterial organisms include *Staphylococcus aureus* and other skin associated bacteria.

Predisposing factors can include excessive sweating, friction, occlusion, shaving and disruption of the skin barrier. Treatment depends on the severity and cause. Mild disease may respond to appropriate topical antiseptic or antimicrobial management, while extensive or recurrent disease requires professional evaluation.

Repeated or severe infections should not be treated solely with self-selected antibiotics because inappropriate antimicrobial use can contribute to resistance and may delay identification of another underlying disorder.

5.1 Pediculosis Capitis

Pediculosis capitis is an infestation of the scalp by the human head louse, *Pediculus humanus capitis*. It commonly produces itching and visible nits attached to hair shafts. Transmission occurs mainly through close contact.

Diagnosis is based on identification of live lice or viable nits. Management requires an appropriate approved pediculicide according to local recommendations, together with measures to reduce reinfestation. Close contacts may require assessment. Sharing combs, brushes, hats and other personal hair items should be avoided during an infestation (Messenger *et al.*, 2023).

5.2 Alopecia Areata

Alopecia areata is an autoimmune form of nonscarring hair loss. It commonly presents as one or more sharply defined patches of hair loss, although extensive forms may involve much larger areas (Gordon and Tosti, 2011; Messenger *et al.*, 2023).

The exact cause is complex and involves immune-mediated disruption of the normal hair follicle growth environment. Diagnosis is generally clinical, supported by examination of Trichoscopy. In selected cases, additional investigations may be required to exclude other causes. Treatment depends on age, extent and duration of disease. Dermatologists may use topical or locally administered anti-inflammatory therapies, and systemic therapies may be considered for more extensive disease. Newer treatment approaches have focused on Janus kinase pathways, reflecting advances in understanding the immunological mechanisms of alopecia areata.

5.3 Androgenetic alopecia

Androgenetic alopecia is one of the most common forms of hair loss. It is associated with genetic susceptibility and follicular sensitivity to androgenic signalling. The condition is characterized by progressive miniaturization of susceptible hair follicles.

In men, hair loss commonly involves the frontal and vertex regions. In women, thinning is often more diffuse over the central scalp. Diagnosis is usually based on clinical examination and characteristic patterns, although laboratory evaluation may be appropriate in selected patients. Evidence-based treatment includes topical minoxidil, while other therapies may be considered by physicians depending on the patient's sex, age and clinical circumstances. Treatment aims to slow progression and maintain or improve hair density where possible (Gordon and Tosti, 2011; Piraccini *et al.*, 2015).

5.4 Telogen Effluvium

Telogen effluvium is a common cause of diffuse hair shedding. It occurs when a greater proportion of follicles prematurely enter the telogen phase. Various physiological and environmental stresses may act as triggers.

Possible precipitating factors include significant systemic illness, major psychological stress, nutritional problems, certain medications and substantial changes in normal body physiology. A detailed history is important because the trigger may have occurred several weeks or months before noticeable shedding (Piraccini *et al.*, 2015).

Management primarily involves identifying and addressing the underlying cause. Hair density commonly improves when the precipitating factor is corrected, although recovery can take time.

5.5 Anagen Effluvium

Anagen effluvium results from interruption of actively growing anagen hairs. It can cause relatively rapid and diffuse hair loss. Certain medical treatments and toxic exposures can interfere with rapidly dividing hair-matrix cells (Piraccini *et al.*, 2015)

The diagnosis is based on history, timing and clinical findings. Management focuses on addressing the underlying cause and providing supportive hair and scalp care. Recovery depends on the cause and the extent of follicular injury.

5.6 Scarring Alopecia

Scarring alopecia represents a group of disorders in which inflammation damages hair follicles and results in permanent replacement by scar tissue. Examples include lichen planopilaris frontal fibrosing alopecia and discoid lupus erythematosus.

Clinical signs can include areas of permanent hair loss, scalp redness, scaling, follicular changes, pain or itching. Early diagnosis is important because treatment aims primarily to control inflammation and prevent further follicular destruction.

Scalp biopsy may be necessary when the diagnosis is uncertain. Management is individualized and may include anti-inflammatory or immunomodulatory treatment under specialist supervision.

6. Diagnosis of Hair and Scalp Disease

A detailed clinical history is the first step in evaluating a hair or scalp disorder. The clinician should document the onset, duration, progression and distribution of hair loss. Information about scalp symptoms, recent illness, medications, nutritional status, family history, cosmetic procedures and hair-care practices can be useful.

Physical examination should include assessment of the scalp, hair shafts, follicular openings and pattern of hair loss. Hair-pull and hair-tug tests may provide additional information in selected cases.

Trichoscopy is a non-invasive examination method that magnifies scalp and hair structures. It can assist in differentiating various forms of alopecia and scalp disease. Practical approaches to hair-loss diagnosis emphasize combining clinical history, examination and appropriate diagnostic tests rather than relying on a single observation.

Laboratory investigations are selected according to the suspected diagnosis. Examples may include complete blood count, iron-related studies, thyroid assessment or other tests when indicated. For suspected fungal infection, potassium hydroxide examination, fungal culture or molecular testing can be useful. Scalp biopsy is reserved for selected cases, particularly when scarring alopecia or an unclear inflammatory condition is suspected.

7. Medical Treatment of Hair and Scalp Diseases

Medical management should be directed toward the underlying cause rather than simply the visible symptom. Infectious diseases require appropriate antimicrobial treatment, inflammatory disorders may require anti-inflammatory approaches, and hair-loss disorders may require therapies aimed at reducing shedding or stimulating growth (Messenger *et al.*, 2023).

For fungal scalp infection, systemic antifungal treatment is generally necessary. Current guidance emphasizes pathogen-directed treatment and appropriate use of adjunctive topical therapy.

For dandruff and seborrheic dermatitis, medicated shampoos can be useful. The choice of active ingredient depends on the condition and individual response. Persistent symptoms warrant professional evaluation.

For androgenetic alopecia, topical minoxidil is an established treatment option. Treatment response varies between individuals, and continued treatment may be required to maintain benefits. For alopecia areata, management depends on the extent and severity of disease. Because it is an immune-mediated disorder, anti-inflammatory and immunomodulatory approaches may be used under medical supervision. Recent research has expanded therapeutic options for severe disease.

Table 1: Classification of hair and scalp diseases

Disease / Condition	Main Cause	Common Symptoms	Hair / Scalp Involvement
Androgenetic alopecia	Genetic and hormonal factors	Gradual hair thinning	Pattern hair loss
Alopecia areata	Autoimmune mechanism	Sudden smooth patches	Patchy hair loss
Telogen effluvium	Stress, illness, nutritional factors	Increased shedding	diffuse hair loss
Anagen effluvium	Chemotherapy, toxic exposure	rapid hair loss	diffuse hair loss
dandruff	Increased <i>Malassezia</i> activity and scalp factors	Flaking and itching	Scalp scaling
Tinea capitis	Dermatophyte infection	Scaling itching broken hair	Scalp infection

8. Surgical Treatment

Surgical treatment is generally considered when hair loss is permanent or when medical therapy alone does not provide satisfactory restoration. Hair transplantation is an important surgical option for selected individuals with stable nonscarring hair loss.

The basic principle of hair transplantation is to relocate healthy hair follicles from a donor region to areas with reduced hair density. Follicular unit transplantation and follicular unit extraction are commonly discussed approaches. Patient selection is essential. The underlying cause of hair loss, donor-hair availability, stability of the disease, age and expectations should be evaluated before surgery. Hair transplantation is generally not appropriate when active inflammatory or scarring disease is uncontrolled. Surgical management should be performed by appropriately qualified medical professionals, with consideration of realistic outcomes and possible complications.

9. Cosmetic Treatment of Hair and Scalp Diseases

Cosmetic management plays an important supportive role in people with hair and scalp disorders. Appropriate cleansing helps remove sebum, sweat, scales and environmental contaminants. Shampoo selection should be based on scalp condition, hair type and individual tolerance (Messenger *et al.*, 2023).

Conditioners can reduce friction between hair fibres and improve manageability. Excessive heat from hair dryers, straightening devices and curling equipment may increase hair-shaft damage. Chemical treatments such as bleaching, permanent waving and repeated colouring can weaken hair when used excessively.

Individuals experiencing hair loss may use cosmetic camouflage products, hair fibres, styling techniques, wigs or hairpieces to improve the appearance of hair density. These methods do not treat the underlying disease but may improve confidence and quality of life.

Cosmetic procedures should be performed carefully in people with active scalp disease because irritation or trauma may worsen symptoms.

10. Hair Care and Preventive Measures

Maintaining good scalp hygiene is an important component of general hair care. Regular cleansing can remove excess oil, sweat and environmental debris. The frequency of washing can be adjusted according to scalp oiliness, hair texture and individual needs. Personal items such as combs, brushes, towels and headwear should not be shared when a contagious scalp infection or infestation is suspected. Early diagnosis of infectious conditions helps reduce transmission.

Avoiding excessive traction is also important. Repeated tight hairstyles can place mechanical stress on hair follicles and may contribute to traction-related hair loss. Excessive chemical processing and high-temperature styling should also be minimized.

11. Role of Microorganisms in Hair Scalp Diseases

Microorganisms have an important relationship with the scalp. Some organisms form part of the normal scalp microbiota, whereas others can cause disease under appropriate conditions. *Malassezia* species are associated with dandruff and seborrheic dermatitis. Dermatophytes such as *Trichophyton* and *Microsporum* cause tinea capitis by invading keratinized tissues and hair. Bacterial organisms such as *Staphylococcus aureus* may be involved in follicular infections.

Understanding the microbial component of scalp disorders is important because it helps distinguish infectious disease from non-infectious inflammatory or hair-growth disorders. In tinea capitis, identification of the causative organism can also assist in selecting appropriate systemic antifungal therapy.

12. Complications

Untreated or inadequately treated hair and scalp diseases can produce complications. Fungal infection may result in persistent inflammation and, in severe cases, scarring alopecia. Delayed

diagnosis of tinea capitis has been associated with the possibility of permanent scarring hair loss (Alkeswani *et al.*, 2019; Messenger *et al.*, 2023).

Inflammatory scarring alopecia can permanently destroy follicles. Therefore, early recognition and specialist management are important. Cosmetic damage may also occur due to repeated chemical processing, heat exposure and mechanical trauma. In addition to physical effects, visible hair loss can influence psychological well-being and social confidence.

13. Recent Advances in Hair and Scalp Treatment

Recent developments in hair medicine include improved trichoscopic diagnosis, molecular investigations, targeted therapies and advances in hair transplantation. Trichoscopy has become an important non-invasive tool for examining scalp and hair structures and supporting diagnosis. Research into alopecia areata has identified immune pathways that can be targeted therapeutically. Genetic studies have also improved understanding of androgenetic alopecia and other hair-loss disorders (Piraccini *et al.*, 2015; Mayser *et al.*, 2020).

In tinea capitis, modern guidelines increasingly emphasize identification of the causative dermatophyte, appropriate systemic therapy and measures to reduce transmission. A 2026 consensus guideline highlights pathogen-guided treatment and infection-control measures, including management of carriers in appropriate settings.

14. Medical, Surgical and Cosmetic Approaches-Integrated Management

Successful management of hair and scalp disease often requires a combination of medical, surgical and cosmetic approaches. Medical treatment addresses the underlying pathological process, while surgical treatment may restore hair in selected cases of permanent loss. Cosmetic treatment helps improve hair appearance and manage daily hair-care requirements.

For example, a person with androgenetic alopecia may initially receive medical treatment to stabilize hair loss. If sufficient donor hair is available and the condition is stable, transplantation may later be considered. Cosmetic camouflage can be used during the treatment period.

Similarly, in tinea capitis, systemic antifungal therapy is essential, while appropriate shampooing and hygiene can provide supportive benefits and help reduce transmission (Messenger *et al.*, 2023; Piraccini *et al.*, 2015).

Conclusion

Hair and scalp diseases represent a broad group of disorders involving infection, inflammation, immune mechanisms, genetics, hormonal factors, systemic disease and cosmetic or mechanical injury. Correct identification of the underlying cause is essential because different conditions require substantially different approaches. Tinea capitis caused by dermatophytes such as *Trichophyton* and *Microsporum* requires systemic antifungal management, while dandruff and seborrheic dermatitis can often be controlled using appropriate scalp-care and medicated shampoo approaches. Hair-loss disorders such as alopecia areata, androgenetic alopecia and

telogen effluvium require individualized assessment and management. Surgical procedures such as hair transplantation may be useful for selected stable cases of permanent hair loss. Cosmetic measures, including appropriate cleansing, conditioning, styling and camouflage, can provide supportive benefits. Overall, an integrated approach involving accurate diagnosis, evidence-based treatment, preventive care and appropriate cosmetic management provides the best framework for maintaining scalp health and hair quality (Messenger *et al.*, 2023; Alkeswani *et al.*, 2019; Piraccini *et al.*, 2015).

References

1. Alkeswani, A., Cantrell, W., & Elewski, B. (2019). Treatment of tinea capitis. *Skin Appendage Disorders*, 5(4), 201–210.
2. Gordon, K. A., & Tosti, A. (2011). Alopecia: Evaluation and treatment. *Clinical, Cosmetic and Investigational Dermatology*, 4, 101–106.
3. Mayser, P., Nenoff, P., Reinelt, D., Brasch, J., Daeschlein, G., Effendy, I., Ginter-Hanselmayer, G., & Zidane, M. (2020). S1 guideline: Tinea capitis. *Journal der Deutschen Dermatologischen Gesellschaft*, 18, 161–170.
4. Messenger, A. G., Harries, M., Macbeth, A. E., Chiu, W. S., Holmes, S., Tziotzios, C., & de Lusignan, S. (2023). Alopecia areata and risk of common infections: A population-based cohort study. *Clinical and Experimental Dermatology*, 48(4), 332–338.
5. Piraccini, B. M., Broccoli, A., Starace, M., Gaspari, V., D'Antuono, A., Dika, E., & Patrizi, A. (2015). Hair and scalp manifestations in secondary syphilis: Epidemiology, clinical features and trichoscopy. *Dermatology*, 231(2), 171–176.

NANOTECHNOLOGY FOR SUSTAINABLE DEVELOPMENT: EMERGING MATERIALS AND APPLICATIONS

Shahnaz Kossar* and Barjinder Kaur

School of Natural Sciences, GNA University, Phaghwara Punjab, India

*Corresponding author E-mail: shazia.rajput0505@gmail.com

Abstract

Nanotechnology has emerged as a multidisciplinary research domain offering innovative solutions for sustainable development through the design and application of advanced nanomaterials. The unique physicochemical properties of nanostructures, including high surface area, enhanced catalytic activity, tunable optical and electronic characteristics, and improved mechanical performance, have enabled their applications in diverse fields such as renewable energy, environmental remediation, healthcare, agriculture, and advanced manufacturing. This chapter provides a comprehensive overview of emerging nanomaterials and their role in addressing global sustainability challenges. It discusses the development of functional materials, including metal and metal oxide nanoparticles, two-dimensional materials, quantum dots, carbon-based nanostructures, perovskites, and hybrid nanocomposites, along with their applications in energy conversion and storage, photocatalysis, water purification, sensors, and biomedical technologies. Particular emphasis is placed on sustainable synthesis approaches, green nanotechnology, and environmentally responsible fabrication methods to minimize ecological impacts. The chapter also highlights recent advances in nanotechnology-driven solutions, including artificial intelligence-assisted material design, smart nanodevices, and scalable manufacturing strategies. Despite significant progress, challenges related to toxicity, long-term environmental effects, cost-effective production, standardization, and commercialization remain critical barriers. Effective technology transfer, industry–academia collaboration, and supportive policy frameworks are essential for translating laboratory innovations into practical applications. By integrating scientific advancements with sustainability principles, nanotechnology holds immense potential to support the achievement of Sustainable Development Goals (SDGs) and contribute to a cleaner, safer, and more sustainable future.

Keywords: Nanotechnology, Nanomaterials, Sustainable Development, Green Synthesis, Energy Storage, Environmental Remediation, Technology Transfer.

1. Introduction

Rapid industrialization, urbanization, and population growth have significantly increased environmental pollution and global energy demand, posing serious challenges to sustainable development. The discharge of toxic organic dyes, pharmaceutical residues, pesticides, heavy

metals, and volatile organic compounds into water and air ecosystems has adversely affected human health and biodiversity. Simultaneously, the excessive dependence on fossil fuels has accelerated greenhouse gas emissions, climate change, and the depletion of non-renewable energy resources. These interconnected challenges have intensified the search for environmentally friendly and energy-efficient technologies capable of simultaneously addressing pollution control and clean energy production. Among the emerging green technologies, semiconductor-based photocatalysis has attracted considerable attention owing to its ability to harness abundant solar energy for environmental remediation and renewable energy generation. Photocatalytic nanomaterials exhibit unique physicochemical properties, including high specific surface area, tunable band gap, enhanced light absorption, and superior charge transport characteristics, making them highly efficient catalysts for various photochemical reactions.

Upon light irradiation, these materials generate electron–hole pairs that participate in oxidation and reduction reactions, producing reactive oxygen species capable of degrading persistent organic pollutants, disinfecting pathogenic microorganisms, and reducing toxic contaminants into less harmful products. Beyond environmental remediation, photocatalytic nanomaterials have emerged as promising candidates for sustainable clean energy applications, including photocatalytic water splitting for hydrogen production, carbon dioxide photoreduction into value-added fuels, and artificial photosynthesis. Recent advances in nanotechnology have further enhanced photocatalytic efficiency through strategies such as metal and non-metal doping, heterojunction engineering, defect and oxygen vacancy engineering, plasmonic modification, and the development of two-dimensional nanostructures and nanocomposites. Materials such as titanium dioxide (TiO_2), zinc oxide (ZnO), bismuth ferrite (BiFeO_3), lanthanum ferrite (LaFeO_3), graphitic carbon nitride ($\text{g-C}_3\text{N}_4$), and spinel ferrites have demonstrated remarkable photocatalytic performance under visible-light irradiation. This chapter presents a comprehensive overview of the fundamental principles, recent developments, synthesis approaches, characterization techniques, environmental applications, clean energy technologies, challenges, and future prospects of photocatalytic nanomaterials, highlighting their transformative role in achieving environmental sustainability, renewable energy production, and the global transition toward a low-carbon economy.

2. Fundamentals of Photocatalysis

Photocatalysis is a light-driven catalytic process in which semiconductor materials utilize solar or artificial light to initiate chemical reactions for environmental remediation and clean energy production. The process begins when a semiconductor photocatalyst, such as TiO_2 , ZnO , BiFeO_3 , LaFeO_3 , or CuFe_2O_4 , absorbs photons with energy equal to or greater than its band gap. The band gap represents the energy difference between the valence band (VB) and the conduction band (CB) and determines the wavelength of light that can be absorbed. Upon photon absorption,

electrons in the valence band are excited to the conduction band, leaving behind positively charged holes in the valence band, thereby generating electron–hole pairs. These photogenerated charge carriers are responsible for initiating photocatalytic reactions; however, rapid recombination of electrons and holes significantly reduces photocatalytic efficiency by dissipating the absorbed energy as heat or light. Therefore, efficient charge separation and migration to the catalyst surface are essential for enhancing photocatalytic performance.

At the semiconductor surface, conduction band electrons react with dissolved oxygen to generate superoxide radicals ($\text{O}_2^{\bullet-}$), while valence band holes oxidize water molecules or hydroxide ions to produce highly reactive hydroxyl radicals ($\bullet\text{OH}$). Together with other reactive oxygen species (ROS), including hydrogen peroxide (H_2O_2) and singlet oxygen, these radicals possess strong oxidation capabilities that effectively degrade organic pollutants, mineralize toxic contaminants into CO_2 and H_2O , inactivate microorganisms, and participate in reduction reactions such as hydrogen evolution and carbon dioxide conversion. The photocatalytic activity of semiconductor nanomaterials is strongly influenced by several intrinsic and extrinsic factors. Crystal structure determines the electronic configuration and charge transport properties, whereas high surface area and nanoscale particle size provide abundant active sites for photocatalytic reactions. Morphology, including nanorods, nanosheets, nanotubes, and hollow nanostructures, further improves light harvesting and charge mobility. Advanced strategies such as band gap engineering through metal or non-metal doping, heterojunction formation, defect engineering, and the introduction of oxygen vacancies extend visible-light absorption, suppress electron–hole recombination, enhance carrier mobility, and create additional active reaction sites. These modifications significantly improve photocatalytic efficiency, making nanostructured photocatalysts promising materials for sustainable environmental remediation and clean energy conversion.

3. Types of Photocatalytic Nanomaterials

Photocatalytic nanomaterials are broadly classified based on their chemical composition, crystal structure, and photocatalytic properties, with each category offering unique advantages for environmental remediation and clean energy applications. Metal oxide semiconductors, including titanium dioxide (TiO_2), zinc oxide (ZnO), tungsten trioxide (WO_3), tin oxide (SnO_2), and hematite ($\alpha\text{-Fe}_2\text{O}_3$), are among the most widely studied photocatalysts because of their excellent chemical stability, non-toxicity, and high photocatalytic activity. Perovskite oxide nanomaterials, such as bismuth ferrite (BiFeO_3), lanthanum ferrite (LaFeO_3), and strontium titanate (SrTiO_3), exhibit narrow band gaps and enhanced visible-light absorption, making them attractive candidates for solar-driven photocatalysis. Spinel ferrite nanomaterials, including cobalt ferrite (CoFe_2O_4), zinc ferrite (ZnFe_2O_4), copper ferrite (CuFe_2O_4), and nickel ferrite (NiFe_2O_4), possess magnetic properties that enable easy recovery and reuse after photocatalytic

reactions, enhancing their practical applicability. Metal sulfide photocatalysts, such as cadmium sulfide (CdS), zinc sulfide (ZnS), and molybdenum disulfide (MoS₂), provide strong visible-light absorption but often require surface modification to improve stability and reduce photocorrosion. Carbon-based nanomaterials, including graphene, graphene oxide, reduced graphene oxide, carbon nanotubes, and graphitic carbon nitride (g-C₃N₄), are widely employed as conductive supports to facilitate charge separation and electron transport.

In addition, metal–organic frameworks (MOFs), covalent organic frameworks (COFs), quantum dots, MXenes, and heterojunction nanocomposites have emerged as next-generation photocatalysts owing to their tunable electronic structures, high porosity, and superior photocatalytic efficiency. The combination of these advanced nanomaterials with strategies such as doping, heterostructure engineering, and defect creation has significantly expanded their applications in pollutant degradation, water purification, carbon dioxide reduction, and solar hydrogen production, contributing to sustainable environmental protection and renewable energy generation.

3.1 Applications in Environmental Remediation

Photocatalytic nanomaterials have emerged as highly effective and sustainable solutions for environmental remediation because of their ability to utilize solar or artificial light to degrade pollutants and detoxify contaminated environments. In wastewater treatment, semiconductor photocatalysts such as TiO₂, ZnO, BiFeO₃, LaFeO₃, g-C₃N₄, and spinel ferrites efficiently degrade hazardous textile dyes, including methylene blue, rhodamine B, methyl orange, and crystal violet, into harmless products such as carbon dioxide and water. These materials are also capable of decomposing persistent pharmaceutical pollutants, including analgesics, anti-inflammatory drugs, hormones, and endocrine-disrupting compounds, which are difficult to remove through conventional treatment processes. Furthermore, photocatalysis has demonstrated remarkable efficiency in degrading antibiotics such as tetracycline, ciprofloxacin, and amoxicillin, thereby reducing the spread of antimicrobial resistance.

The removal of pesticides, including atrazine, chlorpyrifos, and glyphosate, has also been significantly improved through photocatalytic oxidation. In air purification, photocatalytic coatings effectively oxidize harmful nitrogen oxides (NO_x), sulfur-containing gases, and volatile organic compounds (VOCs) emitted from industrial activities and vehicles, while simultaneously improving indoor air quality by eliminating formaldehyde, benzene, toluene, and airborne microorganisms. Photocatalytic nanomaterials also contribute to the removal and detoxification of heavy metals, including hexavalent chromium (Cr(VI)), lead (Pb²⁺), mercury (Hg²⁺), and arsenic (As³⁺/As⁵⁺), through photocatalytic reduction, adsorption, and oxidation processes that convert toxic species into less hazardous forms. Beyond pollutant degradation, these materials exhibit outstanding antibacterial and antiviral activity by generating reactive oxygen species that

damage microbial cell membranes, proteins, and nucleic acids, making them suitable for healthcare and water disinfection applications.

Photocatalytic self-cleaning coatings applied to buildings, glass, ceramics, and textiles utilize sunlight to decompose organic contaminants while maintaining hydrophilic surfaces that facilitate dirt removal. Additionally, photocatalytic nanomaterials have shown considerable potential in oil spill treatment by degrading petroleum hydrocarbons and accelerating the breakdown of complex organic pollutants in aquatic environments. These multifunctional applications demonstrate the transformative role of photocatalytic nanomaterials in advancing sustainable environmental protection, pollution control, and ecosystem restoration.

3.2 Recent Advances in Photocatalytic Nanotechnology

Recent advances in photocatalytic nanotechnology have significantly enhanced the efficiency, stability, and scalability of photocatalysts for environmental remediation and clean energy applications. Among these innovations, single-atom photocatalysts (SACs) have emerged as a breakthrough technology because isolated metal atoms dispersed on suitable supports maximize atomic utilization, provide abundant active sites, and facilitate efficient charge transfer, resulting in superior catalytic activity and selectivity.

Quantum dots (QDs), owing to their quantum confinement effect, possess tunable band gaps, excellent light absorption, and high surface reactivity, enabling improved degradation of organic pollutants and enhanced photocatalytic hydrogen production under visible-light irradiation. Another promising class of materials is MXenes, a family of two-dimensional transition metal carbides, nitrides, and carbonitrides, which exhibit exceptional electrical conductivity, hydrophilic surfaces, and large specific surface areas. When integrated with semiconductor photocatalysts, MXenes accelerate electron transport, suppress electron–hole recombination, and significantly improve photocatalytic efficiency.

High-entropy oxides (HEOs), consisting of multiple cations incorporated into a single crystalline lattice, have recently attracted considerable attention because of their unique electronic structures, enhanced structural stability, and synergistic catalytic properties that enable efficient visible-light-driven photocatalytic reactions. Similarly, two-dimensional (2D) nanomaterials, including graphene, graphitic carbon nitride (g-C₃N₄), transition metal dichalcogenides (MoS₂ and WS₂), and layered double hydroxides, provide ultrathin structures with abundant active sites, shortened charge migration pathways, and excellent interfacial contact, making them ideal candidates for constructing advanced heterojunction photocatalysts.

Beyond material innovations, the integration of artificial intelligence (AI) and machine learning (ML) has transformed photocatalyst discovery by rapidly predicting material compositions, optimizing synthesis parameters, identifying structure–property relationships, and accelerating the screening of high-performance photocatalysts while reducing experimental costs and

development time. Digital twin technology, which combines computational modeling with real-time experimental data, has further enabled virtual optimization of catalyst design, reaction kinetics, and process performance, facilitating faster commercialization of photocatalytic systems.

Equally important are green synthesis approaches, including plant extract-mediated synthesis, microbial synthesis, biotemplating, hydrothermal methods using renewable precursors, and solvent-free fabrication techniques, which minimize hazardous chemicals and reduce environmental impact. Sustainable photocatalyst fabrication also emphasizes recyclable materials, low-energy synthesis routes, earth-abundant elements, and scalable manufacturing processes that support circular economy principles. Collectively, these recent developments are driving the next generation of photocatalytic nanomaterials with improved solar energy utilization, higher catalytic efficiency, greater environmental compatibility, and enhanced commercial viability for applications in wastewater treatment, air purification, carbon dioxide reduction, and renewable hydrogen production, thereby contributing significantly to global sustainability and the transition toward a clean-energy future.

3.3 Challenges and Future Perspectives

Despite remarkable progress in photocatalytic nanotechnology, several scientific and technological challenges continue to limit its large-scale implementation in environmental remediation and sustainable energy applications. One of the primary limitations is the relatively low efficiency of many photocatalysts under visible-light irradiation. Conventional semiconductor photocatalysts, such as TiO_2 and ZnO , possess wide band gaps and can only utilize ultraviolet light, which constitutes less than 5% of the solar spectrum. Consequently, developing narrow band-gap semiconductors, heterostructures, plasmonic materials, and defect-engineered photocatalysts capable of harvesting visible and near-infrared light remains a major research priority.

Another significant challenge is the rapid recombination of photogenerated electron-hole pairs, which substantially reduces photocatalytic efficiency by limiting the generation of reactive oxygen species responsible for pollutant degradation and water splitting. Advanced strategies including heterojunction engineering, cocatalyst deposition, surface defect modulation, and Z-scheme photocatalytic systems have demonstrated improved charge separation; however, achieving long-term stability and high quantum efficiency under practical operating conditions remains difficult. Catalyst stability and photocorrosion also pose serious concerns, particularly for sulfide- and metal-based photocatalysts that undergo structural degradation during prolonged irradiation, resulting in reduced catalytic performance and shorter operational lifetimes.

Improving structural robustness through surface passivation, protective coatings, and composite material design is therefore essential for practical applications. Furthermore, translating

laboratory-scale synthesis methods into economically viable large-scale production presents considerable challenges. Many advanced nanomaterials require complex fabrication techniques, expensive precursors, and energy-intensive processing, making commercialization difficult. Cost-effective and scalable green synthesis routes utilizing renewable resources, biogenic reducing agents, and low-energy manufacturing technologies are increasingly being explored to address these issues. Commercial deployment is further hindered by insufficient industrial validation, inconsistent photocatalytic performance under real environmental conditions, lack of standardized testing protocols, and challenges associated with reactor design and process integration. Economic feasibility remains another critical consideration, as photocatalytic systems must demonstrate superior long-term performance while maintaining competitive production and operational costs compared to conventional treatment technologies.

Environmental safety is equally important because the potential release of engineered nanoparticles into ecosystems may pose ecological and human health risks. Comprehensive life-cycle assessment, toxicity evaluation, nanoparticle recovery, and safe disposal strategies must therefore accompany future photocatalyst development. Incorporating circular economy principles, including catalyst recycling, material recovery, waste valorization, and sustainable resource utilization, will enhance the environmental sustainability of photocatalytic technologies. Looking ahead, future research should focus on designing highly efficient, durable, and environmentally benign photocatalysts through interdisciplinary approaches integrating nanotechnology, artificial intelligence, computational materials science, machine learning, and digital twins for catalyst optimization. The discovery of single-atom catalysts, high-entropy materials, MXenes, two-dimensional semiconductors, and multifunctional heterostructures offers promising opportunities to overcome current limitations.

Additionally, the development of solar-driven continuous-flow photocatalytic reactors, smart monitoring systems, and hybrid treatment technologies integrating photocatalysis with biological, electrochemical, or membrane processes will accelerate commercialization. Ultimately, advancing photocatalytic nanotechnology requires coordinated efforts among researchers, industry, and policymakers to establish standardized performance evaluation, sustainable manufacturing practices, and economically viable technologies capable of addressing global challenges related to clean water, renewable energy, and environmental sustainability.

3.4 Policy Recommendations

The successful transition of photocatalytic technologies from laboratory research to practical environmental and energy applications requires strong policy support, strategic investments, and coordinated efforts among governments, academic institutions, industries, and international organizations. Photocatalysis has significant potential in addressing global challenges such as wastewater pollution, renewable energy generation, carbon dioxide reduction, and hydrogen

production. However, overcoming current technological and economic barriers requires a comprehensive policy framework that promotes innovation, sustainability, and commercialization.

- (a) **Investment in photocatalytic research** is essential for advancing next-generation photocatalytic materials with improved visible-light activity, stability, and scalability. Governments and research funding agencies should provide dedicated grants for fundamental and applied research in nanomaterials, semiconductor engineering, artificial intelligence-assisted catalyst design, and advanced characterization techniques. Long-term funding programs can accelerate the discovery of efficient photocatalysts, including metal-free catalysts, single-atom catalysts, heterostructures, and environmentally friendly materials. Establishing national research centers and shared characterization facilities can further support interdisciplinary innovation and reduce research costs.
- (b) Industry–academia collaboration plays a crucial role in translating scientific discoveries into commercially viable technologies. Universities contribute fundamental knowledge and innovation, while industries provide expertise in large-scale production, engineering design, and market deployment. Collaborative research programs, technology incubators, and public–private partnerships should be encouraged to develop pilot-scale photocatalytic systems for wastewater treatment, hydrogen generation, and environmental remediation. Industry participation in research planning can ensure that developed technologies meet practical requirements related to cost, durability, and operational efficiency.
- (c) The implementation of green manufacturing policies is necessary to ensure sustainable photocatalyst production. Regulations should promote the use of environmentally friendly synthesis methods, reduction of hazardous chemicals, energy-efficient processing, and recycling of catalyst materials. Life-cycle assessment approaches should be incorporated to evaluate the environmental impact of photocatalytic materials from production to disposal. Policies supporting circular economy principles can encourage recovery, regeneration, and reuse of photocatalysts, minimizing waste generation and resource consumption.
- (d) Strict wastewater treatment regulations can create demand for advanced photocatalytic remediation technologies. Governments should establish stringent discharge standards for industrial pollutants, including dyes, pharmaceuticals, pesticides, and heavy metals. Incentives for industries adopting photocatalytic treatment systems can accelerate the replacement of conventional, energy-intensive purification methods with sustainable alternatives. Integration of photocatalysis with existing biological and membrane-based treatment processes may provide efficient and cost-effective solutions.
- (e) Renewable hydrogen initiatives should prioritize photocatalytic water splitting as a clean pathway for hydrogen production. National hydrogen missions and renewable energy

policies should include support for semiconductor photocatalyst development, solar-driven hydrogen generation, and demonstration-scale projects. Investment in solar photocatalytic reactors and hydrogen storage infrastructure can contribute to the global transition toward a low-carbon energy economy.

- (f) International collaborations are vital for knowledge exchange, resource sharing, and addressing global environmental challenges. Joint research projects, international funding programs, and scientific networks can accelerate technological development and standardization of photocatalytic systems. Collaboration among countries can also facilitate the transfer of advanced technologies to developing regions.
- (g) Photocatalytic technologies strongly align with the United Nations Sustainable Development Goals (SDGs), particularly SDG 6 (Clean Water and Sanitation), SDG 7 (Affordable and Clean Energy), SDG 9 (Industry, Innovation and Infrastructure), SDG 12 (Responsible Consumption and Production), and SDG 13 (Climate Action). Policy frameworks should explicitly connect photocatalytic development with these sustainability objectives.
- (h) Finally, effective technology transfer and commercialization strategies are required to bridge the gap between research outcomes and industrial implementation. Governments should support intellectual property protection, startup development, pilot-scale demonstrations, and financial incentives for clean technology adoption. By integrating scientific innovation with supportive policies, photocatalysis can emerge as a key sustainable technology for environmental protection, renewable energy generation, and global climate solutions

Conclusion

Nanotechnology has emerged as a transformative field with significant potential to address global sustainability challenges through the development of advanced materials and innovative applications in energy, environment, healthcare, agriculture, and industrial sectors. However, translating laboratory-scale discoveries into commercially viable technologies requires effective technology transfer mechanisms, strong industry–academia partnerships, and supportive policy frameworks. The successful commercialization of nanomaterials depends on overcoming challenges related to large-scale production, cost reduction, quality control, environmental safety, and regulatory approval.

Collaborative research between universities, industries, and government organizations can accelerate the transition of nanotechnology innovations from research laboratories to real-world applications. Establishing technology incubation centers, promoting intellectual property protection, and encouraging investment in nanotechnology-based startups can further enhance market adoption. Sustainable manufacturing approaches, including green synthesis and circular

economy principles, are essential to ensure that nanotechnology contributes positively to environmental protection and resource efficiency.

Future commercialization strategies should focus on developing safe, affordable, and scalable nanomaterials for renewable energy systems, water purification, pollution control, and advanced healthcare solutions. International collaborations and knowledge exchange will also play a vital role in bridging technological gaps between developed and developing regions. Through responsible innovation and effective technology transfer, nanotechnology can become a key driver for achieving Sustainable Development Goals (SDGs) and building a sustainable, efficient, and resilient future.

References

1. Anastas, P. T., & Zimmerman, J. B. (2018). The United Nations Sustainable Development Goals: How can sustainable chemistry and green nanotechnology contribute? *Current Opinion in Green and Sustainable Chemistry*, 13, 150–153.
2. Chavali, M. S., & Nikolova, M. P. (2019). Metal oxide nanoparticles and their applications in biotechnology. *SN Applied Sciences*, 1(6), Article 607.
3. Husseini, F. S., & El-Kalliny, A. S. (2020). Green nanotechnology for environmental remediation and energy applications. *Journal of Cleaner Production*, 276, Article 123184.
4. Khan, I., Saeed, K., & Khan, I. (2019). Nanoparticles: Properties, applications and toxicities. *Arabian Journal of Chemistry*, 12(7), 908–931.
5. Nasrollahzadeh, M., Sajjadi, M., Dadashi, J., & Ghasemzadeh, H. (2020). Pd-based nanoparticles for catalytic applications in wastewater treatment. *Interface Science and Technology*, 31, 281–347.
6. Santhosh, C., Velmurugan, V., Jacob, G., Jeong, S. K., Grace, A. N., & Bhatnagar, A. (2016). Role of nanomaterials in water treatment applications: A review. *Chemical Engineering Journal*, 306, 1116–1137.
7. Tsuji, J. S., Maynard, A. D., Howard, P. C., Asgharian, B., Davidson, P. R., Oberdörster, G., & Warheit, D. B. (2006). Research strategies for safety evaluation of nanomaterials, part IV: Risk assessment of nanoparticles. *Toxicological Sciences*, 89(1), 42–50.
8. Diallo, M. S., Christie, K. S., & Swaminathan, R. (2015). Nanotechnology for sustainable development: Retrospective and outlook. *Journal of Nanoparticle Research*, 17(4), Article 173.
9. Kharissova, O. V., Dias, H. V. R., Kharisov, B. I., Pérez, B. O., & Pérez, V. M. J. (2013). The greener synthesis of nanoparticles. *Trends in Biotechnology*, 31(4), 240–248.

COMPARATIVE PERFORMANCE EVALUATION OF WIRED AND WIRELESS SMART HOME AUTOMATION USING A PYTHON-BASED VIRTUAL FRAMEWORK

Konduri Ramesh

Electrical and Electronics Engineering,

Bapatla Engineering Collegem Bapatla, India

Corresponding author E-mail: rameshbec.nandur@gmail.com

Abstract

The rapid growth of Internet of Things (IoT) technologies has accelerated the adoption of smart home automation systems for remote monitoring, intelligent control, energy optimization, and improved user convenience. Smart homes commonly employ wired and wireless communication architectures. Wireless systems provide flexibility and easy installation but may experience communication delays, packet losses, and reliability issues. In contrast, wired systems offer reliable, low-latency communication and better operational stability, although they require greater installation effort and infrastructure.

This research presents an AI-Assisted Wired Smart Home Automation Framework for Reliable Energy Management developed using a Python-based virtual home automation environment. A simulation model representing a residential smart home with a hall, bedroom, and kitchen was developed to emulate appliance operation, power consumption monitoring, and intelligent control. The framework generated performance datasets for wired and wireless communication architectures under identical operating conditions. Key performance indicators, including communication delay, packet loss, reliability, and energy efficiency, were evaluated over fifty simulation iterations.

The experimental results demonstrate that the wired architecture provides better performance across all evaluated metrics. The average communication delay was 0.1036 seconds for the wired system compared with 0.2415 seconds for the wireless system. Average packet loss decreased from 6.19% to 1.38%, while reliability improved from 90.84% to 97.80%. Energy efficiency also increased from 80.34% to 90.43%.

These results demonstrate the suitability of wired smart home architectures for reliable residential automation requiring low latency, high reliability, and efficient energy utilization. The proposed virtual framework provides a foundation for future integration of AI-based predictive energy management, fault diagnosis, and real-time IoT-enabled residential automation.

Keywords: Smart Home Automation, Wired Communication, Wireless Communication, Energy Management, IoT, Packet Loss Analysis, Delay Analysis, AI-Assisted Automation.

1. Introduction

Smart home automation has emerged as one of the most influential applications of the Internet of Things (IoT) paradigm, enabling intelligent monitoring and control of residential environments through interconnected devices, sensors, communication networks, and automation controllers. The increasing demand for energy-efficient buildings, improved residential security, remote accessibility, and intelligent resource management has accelerated the adoption of smart home technologies worldwide [1-3].

A smart home may be defined as a residential environment in which household appliances, lighting systems, fans, air-conditioning units, entertainment systems, security devices, and other electrical loads are interconnected through a communication network and managed by a centralized or distributed control system. Through automation technologies, users can remotely monitor and operate appliances, schedule device activities, optimize energy consumption, and receive real-time notifications regarding system status and abnormal events [4].

The success of any smart home infrastructure depends heavily on the communication mechanism employed between controllers and end devices. Two major communication architectures dominate current smart home implementations: wired communication systems and wireless communication systems. Wireless smart home systems generally utilize technologies such as Wi-Fi, ZigBee, Bluetooth, LoRa, Thread, and Z-Wave for device connectivity [5-6]. These technologies eliminate extensive wiring requirements and simplify installation processes. Wireless solutions are particularly attractive in existing buildings where structural modifications may be expensive or impractical. Furthermore, wireless systems support mobility, scalability, and rapid deployment [7].

Despite these advantages, wireless communication technologies are inherently susceptible to several challenges. Signal attenuation, electromagnetic interference, channel congestion, environmental obstacles, and network traffic fluctuations can significantly affect communication performance. These factors often result in increased latency, packet transmission failures, reduced reliability, and energy inefficiencies. As the number of connected devices increases, network congestion may further deteriorate overall system performance [8-10].

Wired home automation systems utilize dedicated communication pathways between automation controllers and appliances. Technologies such as Ethernet, RS-485, CAN Bus, KNX, Modbus, and custom wired communication infrastructures are commonly employed. Since communication occurs through physical transmission media, wired systems generally exhibit lower delays, minimal packet loss, higher reliability, and predictable performance characteristics [11]. Such properties make wired architectures particularly suitable for mission-critical residential applications where communication failures cannot be tolerated [12].

Energy management has become another crucial aspect of modern smart home research. Residential buildings account for a significant portion of global electricity consumption. Efficient monitoring and control of household loads can substantially reduce energy wastage and operational costs. Intelligent energy management systems continuously observe appliance usage patterns, estimate energy demand, and optimize device operations to maximize efficiency while maintaining user comfort [13-15].

Artificial Intelligence (AI) techniques have recently been integrated into smart home environments to further enhance automation capabilities. Machine learning algorithms can predict user behavior, forecast energy consumption patterns, identify abnormal operating conditions, and support intelligent decision-making processes [16]. AI-assisted smart homes are capable of adapting dynamically to changing environmental and occupancy conditions, thereby improving both operational efficiency and user experience [17].

Before deploying smart home solutions in real residential environments, simulation-based development provides an effective approach for evaluating system behavior under controlled conditions [18-20]. Virtual simulation models enable researchers to analyze communication performance, energy consumption characteristics, reliability parameters, and automation strategies without the cost and complexity associated with physical hardware implementation. Such virtual environments facilitate rapid experimentation, algorithm testing, and comparative analysis [21-23].

In this research, a Python-based Virtual Home Automation Model was developed to simulate residential smart home operations. The virtual environment comprises three major residential sections including a hall, bedroom, and kitchen. Various electrical appliances such as lighting systems and fans are represented through software objects capable of emulating real-time operational behavior. A centralized automation controller continuously monitors appliance status and calculates total energy consumption.

The developed simulation framework also incorporates virtual sensing modules capable of monitoring electrical and environmental parameters. These virtual sensors generate operational data used for performance analysis and future AI-based optimization. The framework was further extended to emulate both wired and wireless communication architectures under identical operating conditions. Performance evaluation was conducted using fifty simulation samples generated through the proposed virtual environment. Four major performance indicators were considered:

1. Communication Delay
2. Packet Loss
3. Reliability
4. Energy Efficiency

Communication delay represents the time required for control commands and monitoring information to travel between automation controllers and connected devices. Packet loss quantifies communication failures occurring during data transmission. Reliability measures successful communication performance over time, while energy efficiency evaluates the effectiveness of energy utilization and management mechanisms. The generated dataset was analyzed to compare wired and wireless smart home architectures. Statistical analysis and graphical visualization techniques were employed to investigate performance variations and identify the most suitable communication approach for intelligent residential automation systems.

The primary contribution of this research lies in the development of a scalable virtual smart home framework capable of evaluating communication architectures prior to physical deployment. The obtained results demonstrate significant performance advantages associated with wired communication systems, particularly in terms of latency reduction, reliability improvement, packet loss minimization, and energy management effectiveness.

The findings presented in this study provide valuable insights for researchers, automation engineers, and residential infrastructure designers seeking to implement reliable and energy-efficient smart home solutions. Furthermore, the developed framework establishes a foundation for future integration of AI-assisted prediction models, load optimization algorithms, fault diagnosis techniques, and real-time embedded smart home systems.

2. Problem Statement

Modern wireless home automation systems suffer from:

- High communication delays.
- Packet transmission failures.
- Signal interference.
- Reduced reliability under network congestion.
- Lower energy management efficiency.

There is a need to scientifically evaluate whether wired automation systems provide superior performance compared to wireless systems for intelligent residential energy management applications. Therefore, this research develops a Python-based virtual smart home framework to compare wired and wireless communication architectures using measurable performance indicators.

3. Proposed Virtual Home Automation Framework

3.1 System Architecture

The proposed smart home automation framework is designed as a multi-layered architecture that integrates virtual residential environments, appliance control modules, sensing mechanisms, communication networks, and performance analytics within a unified Python-based simulation

platform. The architecture is developed to emulate the operational behavior of a real-world smart home system while providing a controlled environment for evaluating the performance of wired and wireless communication technologies.

At the foundation of the framework lies the Virtual Home Environment, which represents a typical residential household consisting of three major sections: the hall, bedroom, and kitchen. These rooms are modeled within the simulation to create a realistic smart home scenario where different electrical appliances operate under varying usage conditions. The virtual environment serves as the testbed for implementing automation logic, monitoring energy consumption, and evaluating communication performance.

Within the virtual home environment, a Virtual Appliance Layer is implemented to represent household electrical devices. In the present study, the hall light, bedroom fan, and kitchen light are selected as representative residential loads. Each appliance is assigned a predefined power rating corresponding to realistic energy consumption values. The appliances can be switched ON or OFF through software commands generated by the automation controller, thereby simulating actual home automation operations. The operational state of each appliance directly influences the total energy consumption of the virtual home.

To facilitate intelligent monitoring, the framework incorporates a Virtual Sensor Layer that continuously observes various operational parameters within the smart home environment. These virtual sensors monitor appliance status, power consumption levels, load demand conditions, and overall energy usage. The collected sensor information is periodically updated and supplied to the central control system for decision-making and performance evaluation. This sensor layer emulates the functionality of real-world smart sensors commonly deployed in IoT-based smart homes.

The core intelligence of the framework is provided by the Central Automation Controller, which acts as the supervisory unit responsible for managing all connected devices and system operations. The controller performs multiple functions including device control, data collection, energy calculation, and performance evaluation. Whenever an appliance changes its operational state, the controller updates the system status, calculates the corresponding power consumption, and records the generated data for further analysis. The centralized control architecture ensures coordinated operation of all smart home components and provides a foundation for future integration of artificial intelligence and machine learning algorithms.

To compare communication technologies, the framework incorporates a dedicated Communication Layer consisting of both wired and wireless communication models. The wired communication model simulates stable data transmission through dedicated physical channels, whereas the wireless communication model emulates communication over wireless networks that may experience delays, interference, and packet transmission losses. Both communication

architectures are subjected to identical operating conditions to ensure a fair and unbiased performance comparison.

Finally, an Analytics Layer is integrated into the framework to evaluate the effectiveness of the communication systems. This layer performs detailed analysis of key performance indicators including communication delay, packet loss, reliability, and energy efficiency. Delay analysis measures the response time required for data transmission and device control operations. Packet loss analysis evaluates communication failures occurring during data transfer. Reliability analysis quantifies the consistency and dependability of the automation system, while energy efficiency analysis examines the effectiveness of energy utilization within the smart home environment. The analytical results generated by this layer form the basis for comparing wired and wireless home automation systems and identifying the most suitable communication architecture for intelligent residential energy management applications.

4. Methodology

Step 1: Develop Virtual Home Automation Model using Python.

Step 2: Assign realistic power ratings to appliances.

Step 3: Implement ON/OFF appliance control logic.

Step 4: Simulate wired communication behavior.

Step 5: Simulate wireless communication behavior.

Step 6: Generate 50 communication samples.

Step 7: Calculate:

- Delay
- Packet Loss
- Reliability
- Energy Efficiency

Step 8: Compare average performance.

5. Experimental Dataset

To evaluate the performance of wired and wireless smart home automation systems, a simulation-based dataset was generated using the proposed Python framework. Four key performance parameters—communication delay, packet loss, reliability, and energy efficiency—were measured across both wired and wireless architectures. For each parameter, 50 observations were systematically collected to ensure sufficient data volume for robust statistical analysis and a rigorous comparative evaluation between the two network setups. This structured dataset provides a reliable foundation to analyze communication stability, data integrity, and operational efficiency differences between wired and wireless smart home implementations.

The details of the collected dataset are presented in Table 1.

Table 1: Performance metrics collected from simulation

Parameter	Samples
Wired Delay	50
Wireless Delay	50
Wired Packet Loss	50
Wireless Packet Loss	50
Wired Reliability	50
Wireless Reliability	50
Wired Efficiency	50
Wireless Efficiency	50

Total Records = **50**; Total Features = **8**

6. Average Performance Results

The average values of the collected performance metrics were calculated to compare the overall effectiveness of wired and wireless smart home automation systems. The analysis focuses on four key parameters: communication delay, packet loss, reliability, and energy efficiency. These average results provide a clear indication of the operational characteristics of both architectures and facilitate a quantitative assessment of their suitability for intelligent energy management applications. The computed average values are presented in Table 2.

Table 2: Average performance comparison

Metric	Wired	Wireless
Delay (s)	0.1036	0.2415
Packet Loss (%)	1.38	6.19
Reliability (%)	97.8	90.84
Energy Efficiency (%)	90.43	80.34

7. Results and Discussion




Figure 1. Delay Comparison Plot

The delay comparison plot illustrates communication latency observed in wired and wireless smart home systems. The wired architecture consistently exhibits lower delay values across all simulation instances. The average delay of 0.1036 s demonstrates rapid command execution and response capability. In contrast, wireless communication shows an average delay of 0.2415 s due to signal interference, network congestion, and transmission overhead. The results indicate approximately 57% delay reduction in wired communication.




Figure 2: Packet Loss Comparison Plot

Packet loss analysis indicates that wired communication provides significantly more stable data transmission. The average packet loss of the wired system is only 1.38%, whereas wireless communication experiences 6.19% packet loss. Lower packet loss ensures reliable delivery of appliance control commands and monitoring information.




Figure 3: Reliability Comparison Plot

Reliability analysis demonstrates that wired communication achieves an average reliability of 97.80%, significantly higher than the wireless system's 90.84%. This superior reliability ensures uninterrupted operation of smart home devices, minimizes critical communication failures, and provides a stable, dependable automation environment with minimal data disruption..




Figure 4. Energy Efficiency Comparison Plot

Energy efficiency results indicate that wired communication contributes to more effective energy utilization. The wired framework achieves 90.43% efficiency compared with 80.34% for the wireless system. Reduced retransmissions and stable communication pathways contribute to improved energy management.




Figure 5. Average Performance Comparison Plot

The average performance comparison plot summarizes all evaluated metrics simultaneously. The figure clearly demonstrates the superiority of the wired communication framework in terms of:

- Lower Delay
- Lower Packet Loss
- Higher Reliability
- Higher Energy Efficiency

This figure provides a concise visual validation of the proposed framework.

8. Comparative Performance Improvement

To quantify the benefits of the wired automation architecture, the percentage improvement of the wired system over the wireless system was calculated for each performance metric. The improvement analysis highlights the extent to which the wired approach reduces communication

delays and packet losses while enhancing system reliability and energy efficiency. These results provide a clearer understanding of the performance advantages offered by wired smart home automation in intelligent energy management applications. The calculated improvements are summarized in Table 3.

Table 3: Improvement achieved by wired system

Parameter	Improvement
Delay Reduction	57.08%
Packet Loss Reduction	77.74%
Reliability Improvement	7.66%
Energy Efficiency Improvement	12.56%

The results confirm that wired communication significantly enhances smart home performance and reliability.

9. Future Scope

The proposed Python-based virtual smart home framework provides a strong foundation for future research and development in intelligent residential energy management. The framework can be enhanced by incorporating artificial intelligence and machine learning techniques for load prediction, energy consumption forecasting, and automated decision-making. Advanced fault detection and diagnostic algorithms can further improve system reliability and maintenance efficiency. Future work may also include smart appliance scheduling strategies, integration of renewable energy sources such as solar and wind systems, and real-time implementation using ESP32-based IoT controllers. Additionally, the incorporation of industrial communication protocols such as RS485 and CAN Bus can improve communication robustness, while Edge AI-enabled smart home controllers can facilitate low-latency, intelligent, and autonomous operation of next-generation smart homes.

Conclusion

This paper presented a comprehensive performance evaluation of wired and wireless home automation systems using a Python-based Virtual Home Automation Framework developed for intelligent energy management applications. The virtual environment successfully simulated residential automation operations involving multiple rooms, electrical loads, sensing modules, centralized control, and communication infrastructures.

A dataset consisting of fifty simulation samples was generated to analyze delay, packet loss, reliability, and energy efficiency characteristics of both communication architectures. Experimental findings demonstrated that the wired automation framework consistently outperformed the wireless architecture across all investigated performance metrics.

The wired system achieved an average delay of 0.1036 seconds, packet loss of 1.38%, reliability of 97.80%, and energy efficiency of 90.43%. In comparison, the wireless framework recorded

0.2415 seconds delay, 6.19% packet loss, 90.84% reliability, and 80.34% energy efficiency. These results confirm that wired communication offers superior operational stability, faster response times, enhanced reliability, and improved energy management capabilities.

The proposed virtual framework provides an effective research platform for evaluating smart home communication architectures before physical deployment. The obtained results strongly support the adoption of wired communication infrastructures in residential environments requiring dependable automation, intelligent energy management, and AI-assisted control operations.

References

1. Zaidan, A. A., & Zaidan, B. B. (2020). A review on intelligent process for smart home applications based on IoT. *Artificial Intelligence Review*, 53(1), 141–165.
2. Sepasgozar, S., & Aye, L. (2020). A systematic content review of artificial intelligence and the Internet of Things applications in smart home. *Applied Sciences*, 10(9), 3074–3082.
3. Geraldo Filho, P. R., Loureiro, A. A., & Ueyama, J. (2019). Energy efficient smart home systems infrastructure and decision making process. *Internet of Things*, 5, 153–167.
4. Guo, X., Zhang, Y., & Wu, T. (2019). Review on the application of artificial intelligence in smart homes. *Smart Cities*, 2(3), 402–420.
5. Yu, J., De Antonio, A., & Villalba Mora, E. (2022). Deep learning (CNN, RNN) applications for smart homes: A systematic review. *Computers*, 11(2), 1–26.
6. Nelli, F. (2018). *Python data analytics: With Pandas, NumPy, and Matplotlib*. Apress Publisher.
7. Jo, H., & Yoon, Y. I. (2018). Intelligent smart home energy efficiency model using artificial TensorFlow engine. *Human-Centric Computing and Information Sciences*, 8(1), 1–18.
8. Popa, D., & Castiglione, A. (2019). Deep learning model for home automation and energy reduction in a smart home environment platform. *Neural Computing and Applications*, 31, 1317–1337.
9. Yu, L., & Jiang, T. (2019). Deep reinforcement learning for smart home energy management. *IEEE Internet of Things Journal*, 7(4), 2751–2762.
10. Liu, Y., Zhang, D., & Gooi, H. B. (2020). Optimization strategy based on deep reinforcement learning for home energy management. *CSEE Journal of Power and Energy Systems*, 6(3), 572–582.
11. Khan, M., Seo, J., & Kim, D. (2021). Towards energy efficient home automation: A deep learning approach. *Sensors*, 20(24), 7187–7197.

12. Peng, Z., Li, X., & Yan, F. (2020). An adaptive deep learning model for smart home autonomous system. In *Proceedings of International Conference on Intelligent Transportation, Big Data and Smart City* (pp. 707–710).
13. Sheng, T. J., Islam, M. R., & Islam, M. T. (2020). An Internet of Things based smart waste management system using LoRa and TensorFlow deep learning model. *IEEE Access*, 8, 148793–148811.
14. Ullah, A., Lee, M., & Baik, S. W. (2020). Deep learning assisted buildings energy consumption profiling using smart meter data. *Sensors*, 20(3), 873–882.
15. Shah, A. S., Lajis, A., Ullah, I., & Shah, A. (2020). Dynamic user preference parameters selection and energy consumption optimization for smart homes using deep extreme learning machine and bat algorithm. *IEEE Access*, 8, 204744–204762.
16. Lee, S., & Choi, D. H. (2020). Energy management of smart home with home appliances, energy storage system and electric vehicle: A hierarchical deep reinforcement learning approach. *Sensors*, 20(7), 2157–2164.
17. Jang, H., & Park, S. (2020). Energy storage system management method based on deep learning for energy-efficient smart home. In *Proceedings of IEEE International Conference on Consumer Electronics* (pp. 1–2).
18. Elsis, M., Tran, M. Q., & Darwish, M. M. (2021). Deep learning based Industry 4.0 and Internet of Things towards effective energy management for smart buildings. *Sensors*, 21(4), 1038–1043.
19. Ramesh, J., & Shaaban, M. (2022). Deep learning approach for smart home appliances monitoring and classification. In *Proceedings of IEEE International Conference on Consumer Electronics* (pp. 1–5).
20. Taiwo, O., Oyelade, O. N., & Almutairi, M. S. (2022). Enhanced intelligent smart home control and security system based on deep learning model. *Wireless Communications and Mobile Computing*, 2022, 1–22.
21. Chandola, V., & Kumar, V. (2009). Anomaly detection: A survey. *ACM Computing*, 41(3), 41–58.
22. Pandey, A., & Jain, A. (2017). Comparative analysis of KNN algorithm using various normalization techniques. *International Journal of Computer Network and Information Security*, 11(11), 1–36.
23. Smart home dataset with weather information. (n.d.). Kaggle. <https://www.kaggle.com/datasets/taranvee/smart-home/dataset-with-weather-information>

ADVANCED MATERIALS AND GREEN CHEMISTRY: EMERGING TRENDS IN SUSTAINABLE SCIENCE AND TECHNOLOGY

Savitha H S

SJB Institute of Technology, Bengaluru-560098

Corresponding author E-mail- hssavitha@sjbit.edu.in

Abstract

Chemistry plays a central role in the development of modern science and technology by providing new materials, chemical processes, energy-storage systems, sensors, and environmentally sustainable technologies. The growing demand for advanced materials and sustainable industrial processes has increased interest in green chemistry, nanotechnology, smart materials, functional polymers, biomaterials, and energy materials. Green chemistry focuses on designing chemical products and processes that reduce or eliminate the use and generation of hazardous substances. Advanced materials, including nanomaterials, composites, conducting polymers, shape-memory materials, and stimuli-responsive materials, have applications in electronics, healthcare, energy, environmental remediation, and engineering. Chemistry also contributes significantly to the development of lithium-ion batteries, fuel cells, solar cells, supercapacitors, and hydrogen-energy technologies. In environmental applications, catalytic processes, adsorption, membrane technologies, photocatalysis, and bioremediation provide approaches for controlling pollutants and treating contaminated water. This chapter discusses the role of chemistry in emerging science and technology, with particular emphasis on green chemistry, advanced materials, nanotechnology, sustainable energy, and environmental applications. The integration of chemistry with engineering, biology, physics, and computational technologies can contribute to sustainable technological development and address emerging societal challenges.

Keywords: Green Chemistry, Advanced Materials, Nanotechnology, Smart Materials, Sustainable Chemistry, Energy Materials, Catalysis, Environmental Chemistry, Functional Polymers.

1. Introduction

Chemistry is often described as the central science because it connects physics, biology, materials science, environmental science, medicine, and engineering. Almost every technological development involves chemical principles either in the synthesis of materials, transformation of substances, storage of energy, or control of environmental pollutants. Rapid industrialization and technological development have created an increasing demand for high-performance materials and sustainable chemical processes. Conventional chemical processes may involve hazardous

substances, high energy consumption, and the generation of waste. Therefore, modern chemistry increasingly focuses on designing safer, more efficient, and environmentally responsible processes.

Two important areas contributing to this development are green chemistry and advanced materials. Green chemistry aims to prevent pollution at the molecular level by designing chemical processes that minimize waste and hazardous substances. Advanced materials, on the other hand, are designed to possess specific physical, chemical, electrical, optical, mechanical, or biological properties required for modern applications.

The combination of green chemistry and advanced materials provides opportunities for developing sustainable technologies in energy, electronics, healthcare, water treatment, transportation, and environmental protection.

2. Green Chemistry and Sustainable Technology

Green chemistry is the design of chemical products and processes that reduce or eliminate the use and generation of hazardous substances. It emphasizes prevention rather than treatment of pollution. The principles of green chemistry include waste prevention, atom economy, safer synthesis, safer solvents, energy efficiency, renewable feedstocks, catalysis, and designing products for degradation.

2.1 Atom Economy

Atom economy refers to the efficient incorporation of reactant atoms into the desired product. A reaction with high atom economy produces less unwanted material.

Improving atom economy reduces raw-material consumption and waste generation and therefore contributes to sustainable chemical manufacturing.

2.2 Safer Solvents

Solvents can constitute a significant portion of chemical waste. Green chemistry encourages the replacement of hazardous solvents with safer alternatives whenever technically possible.

Water, ethanol, supercritical fluids, and certain bio-based solvents can be considered in appropriate applications.

2.3 Catalysis

Catalysts increase the rate of chemical reactions without being consumed stoichiometrically. Catalytic processes can reduce energy requirements, improve selectivity, and decrease waste.

Nanostructured catalysts, photocatalysts, biocatalysts, and heterogeneous catalysts are increasingly investigated for sustainable chemical processes.

2.4 Renewable Feedstocks

The use of renewable resources instead of finite fossil-based raw materials is another important aspect of sustainable chemistry. Biomass-derived chemicals, bio-based polymers, and natural products can provide alternatives for selected applications.

3. Advanced Materials in Modern Technology

Advanced materials are materials engineered to exhibit specific properties for particular technological applications. Their development requires an understanding of chemical composition, molecular structure, processing, and structure-property relationships.

Important classes include:

- Nanomaterials
- Smart materials
- Functional polymers
- Composites
- Biomaterials
- Energy materials
- Conducting materials
- Semiconductor materials

These materials are increasingly used in electronics, healthcare, energy conversion, transportation, and environmental technologies.

4. Nanotechnology and Nanomaterials

Nanotechnology deals with materials and structures at approximately the nanometre scale. Nanomaterials can exhibit distinctive properties because of their small dimensions and high surface-area-to-volume ratio. Examples include nanoparticles, nanowires, nanotubes, graphene, quantum dots, and nanocomposites.

4.1 Applications of Nanomaterials

Nanomaterials have applications in:

- **Sensors:** Nanomaterials can provide high sensitivity because of their large surface area and unique electronic properties.
- **Water purification:** Nanostructured adsorbents, membranes, and photocatalysts can be investigated for removing contaminants from water.
- **Medicine:** Certain nanomaterials are being investigated for drug delivery, imaging, diagnostics, and biomedical applications.
- **Energy:** Nanomaterials are used in research on batteries, supercapacitors, solar cells, and fuel cells.
- **Catalysis:** Nanocatalysts can provide high surface area and active sites for chemical reactions.

5. Smart Materials

Smart materials are materials that respond to external stimuli such as temperature, pressure, light, electric fields, magnetic fields, or changes in chemical environment.

Examples include:

- Shape-memory alloys
- Shape-memory polymers
- Piezoelectric materials
- Electrochromic materials
- Thermochromic materials
- Magnetostrictive materials
- Stimuli-responsive hydrogels

These materials have potential applications in sensors, actuators, displays, medical devices, robotics, aerospace systems, and electronic devices.

5.1 Electrochromic Materials

Electrochromic materials change their optical properties when an electrical potential is applied. They can be used in smart windows, displays, mirrors, and other optical devices.

5.2 Shape-Memory Materials

Shape-memory materials can recover a predetermined shape when exposed to an appropriate stimulus, commonly temperature. Their applications include biomedical devices, actuators, and engineering systems.

6. Chemistry of Energy Materials

The transition towards sustainable energy requires efficient energy-conversion and energy-storage technologies. Chemistry contributes directly to the development of batteries, fuel cells, solar cells, hydrogen technologies, and supercapacitors.

6.1 Lithium-Ion Batteries

Lithium-ion batteries are widely used in portable electronic devices and electric vehicles. A typical battery contains a cathode, anode, electrolyte, and separator.

The movement of lithium ions between the electrodes during charging and discharging enables electrical energy storage and release.

Research in battery chemistry focuses on improving:

- Energy density
- Charging rate
- Cycle life
- Thermal stability
- Safety
- Sustainability
- Recycling

6.2 Fuel Cells

Fuel cells convert the chemical energy of a fuel directly into electrical energy through electrochemical reactions.

Hydrogen fuel cells are particularly important in research on clean energy because the electrochemical reaction can produce water as a major product.

Catalysts, membranes, electrodes, and hydrogen-storage materials are important areas of chemical research in fuel-cell technology.

6.3 Solar Cells

Solar cells convert solar radiation into electrical energy. Materials chemistry plays an important role in developing photovoltaic materials with suitable optical and electronic properties.

Research includes silicon-based materials, thin-film materials, organic photovoltaic materials, and perovskite-based materials.

7. Chemistry for Environmental Protection

Environmental chemistry provides scientific principles for understanding pollutants and developing methods for their detection, removal, and prevention.

7.1 Water Treatment

Chemistry-based approaches for water treatment include:

- Adsorption
- Coagulation
- Membrane filtration
- Ion exchange
- Advanced oxidation processes
- Photocatalysis
- Biological treatment

Activated carbon, zeolites, polymeric materials, nanomaterials, and other adsorbents can be used to remove selected contaminants.

7.2 Photocatalysis

Photocatalysis uses light energy to initiate chemical reactions in the presence of a photocatalyst. Semiconductor materials such as titanium dioxide have been widely studied for photocatalytic degradation of organic pollutants.

Photocatalysis demonstrates the integration of materials chemistry, surface chemistry, environmental chemistry, and energy science.

7.3 E-Waste Management

The rapid growth of electronic devices has resulted in increasing quantities of electronic waste. E-waste may contain valuable metals such as copper, silver, gold, and other recoverable materials, along with potentially hazardous components.

Chemistry contributes to e-waste management through processes such as hydrometallurgy, pyrometallurgy, bioleaching, solvent extraction, precipitation, and adsorption.

Developing environmentally safer methods for recovering valuable materials is an important area of sustainable chemistry.

8. Functional Polymers and Composites

Polymers are essential materials in modern technology because of their versatility, low density, processability, and range of properties.

Functional polymers can possess electrical, optical, thermal, mechanical, or biological functions.

Conducting polymers such as polyaniline, polypyrrole, and polythiophene have applications in sensors, electrochemical devices, energy storage, and electronic systems.

Polymer composites combine a polymer matrix with reinforcing materials to obtain improved mechanical, thermal, electrical, or barrier properties.

Research is also increasingly focused on biodegradable and bio-based polymers to address environmental concerns associated with conventional plastics.

9. Role of Chemistry in Emerging Sensors

Chemical sensors detect specific chemical or biological species and convert the interaction into a measurable signal.

A chemical sensor generally consists of:

Analyte → Recognition element → Transducer → Signal processing → Output

Nanomaterials, conducting polymers, metal oxides, graphene-based materials, and biomolecules are being investigated for sensor development.

Applications include:

- Detection of toxic gases
- Water-quality monitoring
- Food-quality assessment
- Medical diagnostics
- Environmental monitoring
- Industrial process control

The integration of chemical sensors with IoT technologies can enable real-time monitoring and remote data collection.

10. Challenges in Sustainable Chemical Technology

Although chemistry provides many technological solutions, several challenges remain.

10.1 Resource Availability

Many advanced technologies require critical elements and specialized raw materials. Sustainable alternatives and efficient recycling methods are therefore necessary.

10.2 Energy Consumption

Some chemical processes require considerable energy. Development of catalysts, renewable-energy-driven processes, and low-temperature synthesis can help reduce energy requirements.

10.3 Waste Generation

Chemical and materials industries can generate solid, liquid, and gaseous waste. Waste prevention and resource recovery are important components of sustainable chemistry.

10.4 Environmental and Health Safety

New materials should be evaluated for their potential environmental and biological effects. Safety assessment should accompany technological development.

10.5 Recycling and Circular Economy

Materials should increasingly be designed with recovery, reuse, recycling, and end-of-life management in mind. A circular approach can reduce resource consumption and waste.

11. Future Perspectives

The future of chemistry will increasingly involve interdisciplinary integration. Computational chemistry, artificial intelligence, nanotechnology, biotechnology, materials science, and green chemistry are expected to work together in the development of new products and processes.

AI and machine learning can assist researchers in predicting material properties and identifying potential chemical compounds. Nanotechnology can provide materials with tailored properties. Green chemistry can help ensure that their synthesis is safer and more sustainable.

Future research is likely to focus on:

- Green synthesis of nanomaterials
- Sustainable batteries
- Hydrogen production and storage
- Bio-based polymers
- Smart sensors
- Carbon capture and utilization
- Sustainable catalysis
- Advanced water-treatment technologies
- Recycling and recovery of critical materials
- AI-assisted materials discovery

Such developments demonstrate that chemistry is not only a fundamental science but also a major contributor to technological innovation and sustainable development.

Conclusion

Chemistry provides the foundation for many of the materials, processes, and technologies that support modern society. Green chemistry promotes safer and more sustainable chemical

processes, while advanced materials provide new functionality for applications in electronics, energy, healthcare, environmental protection, and engineering. Nanomaterials, smart materials, functional polymers, catalysts, energy-storage materials, sensors, and environmentally friendly chemical processes demonstrate the growing relationship between chemistry and technology. The future development of these areas will require interdisciplinary collaboration among chemists, physicists, biologists, engineers, computational scientists, and environmental researchers.

Sustainable chemistry should focus not only on developing high-performance materials and processes but also on resource efficiency, safety, recyclability, energy efficiency, and environmental responsibility. The integration of green chemistry with advanced materials and emerging technologies can therefore contribute significantly to sustainable technological development and the achievement of a cleaner and more resource-efficient future.

References

1. Anastas, P. T., & Warner, J. C. (1998). *Green chemistry: Theory and practice*. Oxford University Press.
2. Anastas, P. T., & Eghbali, N. (2010). Green chemistry: Principles and case studies. *Chemical Society Reviews*, 39, 301–312.
3. Poole, C. P., & Owens, F. J. (2003). *Introduction to nanotechnology*. Wiley-Interscience.
4. Cao, G., & Wang, Y. (2011). *Nanostructures and nanomaterials: Synthesis, properties, and applications* (2nd ed.). World Scientific.
5. Callister, W. D., & Rethwisch, D. G. (2018). *Materials science and engineering: An introduction* (10th ed.). Wiley.
6. Miessler, G. L., Fischer, P. J., & Tarr, D. A. (2014). *Inorganic chemistry* (5th ed.). Pearson.
7. Atkins, P., & de Paula, J. (2014). *Physical chemistry* (10th ed.). Oxford University Press.
8. Housecroft, C. E., & Sharpe, A. G. (2018). *Inorganic chemistry* (5th ed.). Pearson.
9. Bard, A. J., & Faulkner, L. R. (2001). *Electrochemical methods: Fundamentals and applications* (2nd ed.). Wiley.
10. United Nations. (2015). *Transforming our world: The 2030 agenda for sustainable development*. United Nations.
11. International Energy Agency. (2023). *World energy outlook 2023*. IEA, Paris.
12. World Health Organization. (2021). *WHO global air quality guidelines*. World Health Organization, Geneva.

FUZZY COGNITIVE MODELLING OF INNOVATIVE PEDAGOGIES FOR ENHANCING COMMUNICATIVE SKILLS AMONG RURAL STUDENTS AND PROMOTING INCLUSIVITY

Nivetha Martin^{*1} and A Sahaya Josephine Mary²

¹Department of Mathematics,

²Department of English,

Arul Anandar College (Autonomous), Karumathur, Tamil Nadu 625514

*Corresponding author E-mail: nivetha.martin710@gmail.com

Abstract

Communication skills are indispensable for student progression and placements. The existing digital divides between urban and rural learners obstruct inclusive education. This study proposes a fuzzy cognitive modelling framework to explore the effectiveness of diverse pedagogies in enhancing the communication skills of rural students and inclusivity. This model determines the key pedagogical and learner -related factors associated with modern educational system. The proposed architecture considers factors pertinent to student development, communicative competence and inclusivity. The model intends to determine the interactions of the input pedagogical concepts and their influence over the intermediary concepts of student engagement and outcome concepts of communication competence cum inclusivity. Peer learning and context based pedagogy is identified as the key pedagogies contributing to communication competency of the rural learners. The sensitivity analysis performed at $\pm 10\%$ confirms the rankings of pedagogies. The fuzzy cognitive modelling supports the decision makers and policy makers to comprehend better on the interconnected and interdependent educational factors contributing to communication efficiency and inclusivity of rural education.

Keywords: FCM, Innovative Pedagogies, Communication Skills, Rural, Inclusivity.

Introduction

Inclusive education is one of the sustainable goals and has emerged as the fundamental principle of equitable and sustainable education. In Indian context, the concept of inclusivity has moved beyond from enrollment to accessibility and social integration. The framework of Right to Education and Samagraha Shiksha emphasize on inclusive education and it is specifically pronounced at greater magnitude in rural India. The students of rural regions are deprived of educational advantages and constrained with limited educational resources, language barriers, digital infrastructure and diverse learning environments. Inclusion doesn't focus only on classroom enrollment but also on creating effective pedagogies that foster English proficient skills. Ongoing researches on language proficiency among rural high school students especially

in the regions of Tamil Nadu vividly present the student's reluctance and hesitancy in English communication.

On other hand the recently evolved digital technologies in education support inclusive and learner-centered pedagogy. However, persistent inequalities exist between rural and urban learners. The rural users rely on shared devices in comparison with the urban users. In 2024, 24% of rural users accessed internet using others devices. Furthermore, 57% of the rural adolescent aged 14-16 use smartphone for educational related activity and only a significant percentage of them use their own device. The existence of technology or digital infrastructure is not just sufficient rather the learners must have the facility for productive access.

Communication skills are essential for academic enrichment, employability and lifelong learning. The rural students are deprived of communication ability due to limited exposure to English speaking environments. One of the primary reasons identified is that the mother tongue is the medium of their instruction. The students receive fewer opportunities to develop their LSRW skills English language. The limited exposure obstructs them in articulating their knowledge. The conventional pedagogies are not sufficient to foster the skills of English speaking in this digital era. Hence, it is essential to employ innovative pedagogies suiting to the dynamic demands of the learners. The educators suggest to adopt pedagogies such as digital enabled learning, collaborative learning, experiential learning, peer learning, context based learning to enhance communication efficiency of the students. The choice of these pedagogies depends on student engagement and inclusive education. It is essential to explore the interactions between these pedagogies and learner's engagement. The study of inter relational impacts requires a mathematical framework to examine the interactions between the factors of study. Fuzzy Cognitive Map (FCM) modelling is one of the most preferred mathematical tools to determine the interactions between the factors of the study. The model considers three categories of factors considering interactive pedagogies, student engagement and inclusivity.

FCM is basically characterized as a directed graph with nodes and edges representing the factors and their interactions respectively. The magnitude or the strength of impacts is quantified using values between -1 and 1. The value 1 represents perfect positive interaction, the value -1 indicates perfect negative impacts and the value 0 signifies null impact or influence. FCMs are widely applied in several fields to handle complex decision-making scenario and the domain of education is not an exception to it. The FCM model applied in this work examines the interactions between the factors considered for the study. This model also identifies the most influential concept subjected to pedagogy and student engagement. The remaining contents are structured into the following sections. The brief review of works related to FCM applications in education, contemporary pedagogies and inclusive education. The materials and method are presented in section 3. The results and discussions are discoursed in section 4. The last section concludes the work with future directions.

2. Review of Works

This section presents the recent and relevant state of art of works related to this study. The research gaps and significant contributions are also discussed in this work.

2.1 Innovative Pedagogies for Enhancing Communicative Skills

The English language educators and trainers have shifted from conventional teacher centric instruction methods to a more learner centered methods of teaching. Participatory method of pedagogy is gaining momentum these days in comparison with lecture method. Hua *et al.* presented a comprehensive review of drama based pedagogy highlighting the potential of this pedagogy for enhancing English speaking proficiency through role enactment and collaborative interaction. Hui examines the pivotal role of drama -based pedagogy for developing the speaking competency of EFL students. This kind of pedagogy also perform a crucial role in mitigating learner's hesitation and provide wide opportunities to engage effectively in oral communication. Flipped classroom pedagogy is identified as another significant measure to improve language proficiency. Chen *et al.* discussed the importance of flipped learning in the development of business English and in fostering learner participation and engagement in classroom. Zhu *et al.* also demonstrated the effectiveness of flipped classroom pedagogy in self-directed learning. Digital pedagogy has emerged as another contemporary approach to English teaching. Hassan discussed the role of digital pedagogy in enriching English-speaking skills. On other hand, Dai and Xu explored the effectiveness of AI enhanced pedagogy in enriching listening and speaking skills. Researchers also examined the potential of debate and presentation based approached in English speaking. Waluyo *et al.* exhibited the ways of how debate pedagogy shall contribute to English speaking fluency. De los Idiomas and Extranjeros focused on presentation pedagogy and demonstrated how presentation activities facilitate speaking fluency in virtual English classes. These collective studies show how drama, debate, flipped learning, presentations, AI -integrated learning, collaborative learning contribute to English proficiency.

Zhou *et al.* examined the effects of AI driven pedagogy in English speaking education. Khan *et al.* discussed diverse trends of AI pedagogy. Charles added the aspect of translanguaging and presented the importance of learner's linguistic resources in English language classrooms. Collectively these studies state that communication efficiency is influenced not by a single teaching method but by an interconnected set of pedagogy. A majority of research works focus only on individual pedagogies. There exists only a limited comparison of diverse pedagogies and there exists no research on the study of pedagogical impacts over rural learners. This provides a strong rational for employing FCM model to explore the interrelational impacts among pedagogical interventions and student engagement outcomes.

2.2 Inclusivity and Rural Education

Inclusivity in education shall be interpreted not only in the aspect of physical provision to educational institutions. It involves more interpretation especially in the perspective of

meaningful participation, equitable learning opportunities and suitable pedagogical support to all the learners of diverse background. Rangarajan *et al.* studied inclusive education in the context of rural schools with special reference to the regions of Uttarakhand. The findings of this work state the strong influence between inclusivity and social cum educational context. The rural educational realities are also demonstrated in this work. Murasing in his systematic review on inclusive education in rural India, identified the crucial challenges associated with educational inequalities.

Pandey *et al.* discussed the essential transformation to be implemented to promote inclusion in education. This work focused on the impacts of sociocultural factors over inclusivity. Rajguru in his work highlighted that deprived and limited infrastructure undermine inclusive education. It is highly essential to bridge the educational gaps and promote inclusivity. George *et al.* suggested technology as a means of fostering inclusivity and demonstrated how technology driven education shall bridge urban and rural gaps. Kumari also discussed the intervention of artificial intelligence for advancing inclusive education. Inclusivity and communication competence are often treated as separate entities. However, it is essential to study the interactions between these two dimensions. Studies on English speaking do not adequately address the challenges of inclusion of rural regions. Consequently, there is a need of an integrated framework considering communicative competence and inclusivity.

2.3 Fuzzy Cognitive Mapping in Education

FCMs are competent mathematical structures applied in exploring the interactions between the factors of the study. This FCM model is widely applied to address the challenges of education domain. Teh and Kudus developed a FCM model to evolve an optimal decision-making framework in the context of complex educational relationships. Lepore demonstrates how FCM shall be applied in modelling the cognitive factors contributing to mathematics education. Maroungkas *et al.* explored how FCM is employed to handle adaptive virtual-reality game for programming education. Maroungkas further developed a FCM model integrating intelligent virtual reality and human-centred education. These studies demonstrate the capacity of FCM in examining the factors of pedagogy, learners and technology. Wang et al, Sharma et al, Nyirenda and Chisanga developed integrated FCM models to deal with intricate institutional and academic processes. The aforementioned review of works exhibits the efficiency of FCM in handling uncertain and conflicting factors. The literature also demonstrates the efficiency of FCM in conjoining expert knowledge, causal relationships, feedback mechanisms and dynamic interactions.

2.4 Research Gaps and Significant Contributions

The review of works on pedagogies, FCM in education and inclusivity clearly state the interconnections between the entities. However, the existing research works do not explore much

on their intersections. FCM models applied in communication skill development are very limited in literature. The FCM modelling encompassing diverse pedagogies and their interactions considering inclusivity and student engagement in rural context is essentially required to be developed. The mapping between research gaps and significant contributions of this work is presented in Figure 1.




Figure 1: Mapping Research Gaps to Significant Contributions

3. Materials and Methods

3.1 Research Design

The study applies a Fuzzy Cognitive Map (FCM) approach to determine the causal relationships among innovative pedagogies, communicative-skill development, and inclusive learning among rural students. FCM is highly appropriate for this work as this study involves educational factors which are interconnected. FCM structure comprises of nodes and edges signifying concepts and their causal relationships.

3.2 Identification of FCM Concepts

The concepts considered for this study are:

C1 = Digital/ICT-Enabled Learning

C2 = Collaborative Learning

C3 = Experiential/Activity-Based Learning

C4 = Peer Learning

C5 = Local/Context-Based Pedagogy

C6 = Student Engagement

C7 = Communication Practice

C8 = Confidence and Participation

C9 = Communicative Competence

C10 = Inclusive Learning

Here, C1-C5 represent the major innovative pedagogical interventions, C6-C8 represent intermediate outcomes, and C9-C10 represent the major educational outcomes. For better visualization, the classification of concepts is presented in Figure 2.




Figure 2: Classification of the Concepts

3.3 Selection of Expert Panel

For a FCM model, it is essential to include expert's knowledge. Three educators of diverse background are chosen as expert panel. The educators provided their individual judgements regarding the direction and strength of influence between the identified concepts.

The individual expert connection matrices are represented as:

$$W^{(1)} = [w^{(1)}_{ij}]$$

$$W^{(2)} = [w^{(2)}_{ij}]$$

$$W^{(3)} = [w^{(3)}_{ij}]$$

where:

$w^{(k)}_{ij}$ = causal influence of concept C_i on concept C_j according to educator k .

Here, $i, j = 1, 2, \dots, 10$ and $k = 1, 2, 3$.

The composition of the expert panel is presented in Table 1.

Table 1: Expert Panel

Expert	Domain of Expertise	Areas Focused
Educator 1	ICT/English-oriented perspective	Digital learning and communication practice
Educator 2	Rural pedagogy-oriented perspective	Local context, peer learning and experiential learning.
Educator 3	Inclusive-education-oriented perspective	Participation, confidence, communication and inclusion

3.4. Fuzzy Weight Assignment

The strength of the causal relationship was expressed using fuzzy weights in the interval:

$$-1 \leq w_{ij} \leq 1$$

The following linguistic scale was adopted:

No influence = 0.0

Very weak influence = 0.1

Weak influence = 0.3

Moderate influence = 0.5

Strong influence = 0.7

Very strong influence = 0.9

A positive weight indicates a positive causal relationship, whereas a negative weight indicates a negative causal relationship.

3.5 Construction of Individual Connection Matrices

For each educator, a 10×10 connection matrix was constructed.

The general form is:

$$W^{(k)} = [w^{(k)}_{ij}]$$

where:

$w^{(k)}_{ij}$ = influence of C_i on C_j according to educator k .

3.6 Aggregation of Expert Judgements

The three individual connection matrices are combined to obtain a single aggregated FCM.

The aggregated connection matrix is calculated using the arithmetic mean:

$$W = \frac{(W^{(1)} + W^{(2)} + W^{(3)})}{3}$$

For each causal relationship, the aggregated weight is:

$$w_{ij} = \frac{w_{ij}^{(1)} + w_{ij}^{(2)} + w_{ij}^{(3)}}{3}$$

3.7 Initial State Vector

The activation level of each concept is represented using a state vector:

$$A^{(t)} = [A_1^{(t)}, A_2^{(t)}, \dots, A_{10}^{(t)}]$$

where:

$A_i^{(t)}$ = activation level of concept C_i at iteration t .

The initial state is:

$$A^{(0)} = [A_1^{(0)}, A_2^{(0)}, \dots, A_{10}^{(0)}]$$

Each activation value lies within the interval:

$$0 \leq A_i^{(t)} \leq 1$$

3.8 FCM State Updating

The activation level of each concept is updated using the FCM propagation rule:

$$\text{where: } A_i(t+1) = f\left(\sum_{j=1}^{10} A_j(t)w_{ji}\right)$$

$A_{i(t+1)}$ = activation level of concept C_i at iteration $t+1$

$A_{j(t)}$ = activation level of concept C_j at iteration t

w_{ji} = causal weight from concept C_j to concept C_i

$f(.)$ = threshold or activation function.

The sigmoid activation function can be defined as:

$$f(x) = \frac{1}{1 + e^{-\lambda x}}$$

where $\lambda > 0$ is the steepness parameter.

The iterative process is:

$$A^{(0)} \rightarrow A^{(1)} \rightarrow A^{(2)} \rightarrow \dots \rightarrow A^{(t)}$$

Iteration continues until convergence:

$$|A_i^{(t+1)} - A_i^{(t)}| < \varepsilon$$

where ε is a small convergence threshold.

3.9 Intervention Analysis

Five intervention scenarios are considered, with one pedagogical concept activated at a time:

$$C_1 = 1$$

$$C_2 = 1$$

$$C_3 = 1$$

$$C_4 = 1$$

$$C_5 = 1$$

For every intervention, the final activation levels of C_9 (Communicative Competence) and C_{10} (Inclusive Learning) are recorded. The overall impact of each pedagogy is calculated as:

$$IMPACT(C_i) = \frac{C_9(FINAL) + C_{10}(FINAL)}{2}$$

3.10 DEGREE-BASED INFLUENCE ANALYSIS

The structural influence of each concept is evaluated using its out-degree and in-degree.

The outgoing and incoming influences of each concept are calculated using:

$$OD(C_i) = \sum_{j=1}^{10} |w_{ij}|$$

$$ID(C_i) = \sum_{j=1}^{10} |w_{ji}|$$

A high out-degree indicates a strong driving concept, while a high in-degree indicates a concept strongly affected by other concepts. The connection matrix given by each individual educators is presented in Figure 3.




Figure 3: Perspectives of Educators

4. Results and Discussions

By following the above discussed procedure as in section 3, the following results are obtained. The FCM convergence plot of the key outcomes is presented in Figure 4




Figure 4 FCM Convergence Plot of Outcome Concepts

The out and in degrees of the concepts are presented in Figure 5.

From the Fig.5, it is observed that C4, C3 and C2 are the strongest driving concepts according to OD. C10 has the highest ID (6.91), followed by C9 (5.47), showing that the final outcomes are strongly influenced by the other concepts.




Figure 5: Out - Degree and In - Degree of Concepts

The ranking of the pedagogies is presented in Fig.6. The score values are presented in Table 2.




Figure 6: Ranking of Pedagogies

Table 2: Impact Scores of Pedagogies to Outcome Concepts

Rank	Pedagogy	C9 Influence	C10 Influence	Impact Score
1	Peer Learning (C4)	0.67	0.77	1.44
2	Local/Context-Based Pedagogy (C5)	0.57	0.87	1.44
3	Experiential/Activity-Based Learning (C3)	0.67	0.70	1.37
4	Collaborative Learning (C2)	0.60	0.70	1.30
5	Digital/ICT-Enabled Learning (C1)	0.50	0.50	1.00

From the Fig.6, it is inferred that

- C4 Peer Learning has stronger direct influence on C9.
- C5 Local/Context-Based Pedagogy has substantially stronger direct influence on C10.

4.1 Sensitivity Analysis

A $\pm 10\%$ perturbation is applied to validate the ranking results of the pedagogies. The numerical results are presented in Table 3. For better visualization, the results are presented in Fig.7.

Table 3: Sensitivity Results

Pedagogy	Base	-10%	+10%
C1 Digital/ICT	1.00	0.90	1.10
C2 Collaborative	1.30	1.17	1.43
C3 Experiential	1.37	1.23	1.51
C4 Peer	1.44	1.30	1.58
C5 Context-Based	1.44	1.30	1.58




Figure 7: Sensitivity Results

Conclusion and Future Directions

This research work has proposed a FCM model to explore the interactions between different pedagogies and their impacts over communicative competence and inclusive education. The FCM results demonstrate that innovative pedagogies have positive and strong interconnected impacts over communication and inclusivity. On other hand among the five pedagogies, Peer learning and Context based pedagogies are identified to be more feasible for the rural students to develop their English-speaking fluency. The high out-degree values of C4, C3, and C2 further state that these are the major driving concepts in the FCM. In contrast, C10 recorded the highest in-degree, exhibiting that inclusive learning is a predominant outcome influenced by both pedagogical and communicative factors. The sensitive analysis is also performed to confirm the ranking results of the pedagogies.

The pedagogies of peer learning and context based are highly suitable for the rural students to develop their communication skills. Also, the pedagogical impacts over inclusivity is also

reflected in the impact scores. Thus, FCM serves as the decision support tool in exploring pedagogies over communication skills and inclusive education in rural context. This research work shall be further extended discussing comparisons between urban and rural perspectives. The same approach of FCM shall also be employed in diverse educational settings.

References

1. Castro Filho, E. D. D., Gusso, G. D. F., Demarzo, M. M. P., Stein, A., Schneider, J. C., Dalla, M. D. B., ... & Aguiar, R. A. T. D. (2007). Specialization in FCM and the challenge of medical qualification for the Family Health Strategy: A proposal for large-scale specialization through distance education.
2. Charles, A. A. (2026). Translanguaging in language pedagogy: Perceptions of French-speaking learners of English in an English language classroom in Cameroon. *EPASA MOTO*, 1(1), 129–152.
3. Chen, Y., Islam, M. M., Sun, Z., Weimeng, W., & Yasmin, M. T. (2026). Reconceptualising Business English speaking development: Flipped classroom pedagogy as mediated participation in an ESP context. *English for Specific Purposes*, 84, 299–315.
4. Dai, Y., & Xu, J. (n.d.). AI-enhanced reform of English listening and speaking pedagogy in vocational education: Theoretical construction and empirical validation.
5. De los Idiomas, C. D. P., & Extranjeros, N. (2025). Applying short presentations in virtual English classes to improve speaking fluency within eight level students in the Pedagogy of National and Foreign Languages degree at ESPE, Rumiñahui Canton, Pichincha Province during the academic term: April–September 2025.
6. George, S., Karalam, S. R. B., & Bhavana, S. (2026). Digital pathways to menstrual health education in India: Bridging urban-rural gaps through inclusive technologies. *Social Sciences & Humanities Open*, 14, 103367.
7. Hassan, M. (2025). A digital pedagogy approach to teaching speaking in the 21st-century English classroom: An analysis and literature-based study. *Proceedings Academic Universitas Jambi*, 1(3), 1178–1184.
8. Hegedüs, J., Kóczy, L. T., & Hatwagner, M. F. (n.d.). A scenario-based fuzzy cognitive map approach to ESG analysis in higher education. In *Book of abstracts* (p. 10).
9. Hua, X., Hashim, H., & Jamaludin, K. A. (2025). A comprehensive review of the impact of drama-based pedagogy on English-speaking proficiency. *Eurasian Journal of Applied Linguistics*, 11(1), 275–282.
10. Hui, X. (2025). A study of EFL students English speaking proficiency using dramaas pedagogy. *วารสาร มจร อุบลราชธานี*, 10(1), 3269–3284.
11. Khan, S., Abu Ziden, A., & Khan, Z. (2025). A bibliometric analysis of research on CEFR-based English speaking pedagogy: Trends and directions [Special issue]. *PASAA*, 73, 9–46.

12. Kumari, A. (n.d.). Bridging the educational divide through artificial intelligence: Advancing inclusive education for rural and underprivileged children in India.
13. Lepore, M. (2024). A holistic framework to model student's cognitive process in mathematics education through fuzzy cognitive maps. *Heliyon*, 10(16).
14. Maroungkas, A. (2026). *Adaptive and intelligent virtual reality through fuzzy cognitive maps for human-centered education* (Doctoral dissertation, Πανεπιστήμιο Δυτικής Αττικής, Σχολή Μηχανικών, Τμήμα Μηχανικών Πληροφορικής και Υπολογιστών).
15. Maroungkas, A., Troussas, C., Krouska, A., & Sgouropoulou, C. (2025). An adaptive virtual reality game for programming education using fuzzy cognitive maps and pedagogical models. *Smart Learning Environments*, 12(1), 62.
16. Mourchid, M. (2025). *Moroccan EFL learners' attitudes towards World Englishes and native and non-native English-speaking teachers: Towards a Global Englishes-informed pedagogy*. GRIN Verlag.
17. Murasing, P. K. (2025). Inclusive education in rural India: Challenges and prospects for sustainable development—A systematic review (2005–2025). *International Journal of Social Sciences*, 7(2).
18. Nyirenda, V. R., & Chisanga, C. B. (2026). Potential AI-based use of fuzzy cognitive mapping in postgraduate supervision in higher education. *Open Books and Proceedings*, eight–130.
19. Pandey, P. N., Singh, P., Shukla, K., Upadhyay, S., Jaiswal, P., & Singh, T. B. (2026). Transformation of traditional rural customs and its impact on inclusive education for children with disabilities in India. *International Journal of Special Education*, 41(11s), 81–94.
20. Rajguru, P. P. (2024, December). Bridging the gap: Addressing infrastructure and education deficiencies in rural India for inclusive growth. In *Techno-Societal 2016, International Conference on Advanced Technologies for Societal Applications* (pp. 3–10). Springer Nature Switzerland.
21. Rangarajan, R., Grove, C., Sharma, U., & Odier-Guedj, D. (2023). A tapestry of multiple perspectives: Contextualising inclusive education through the study of a rural government school in Uttarakhand, India. *International Journal of Educational Research*, 119, 102160.
22. Ravasizadeh, M., Nobakht Ramezani, Z., & Bahrami, S. (2018). Designing of affective factors of knowledge maturity in employees of Physical Education and Health Department uses the FCM model. *Research on Educational Sport*, 6(15), 161–178.
23. Sharma, S., Sanwal, R., Jakhongir, T., Nilufar, S., Ozimbay, J., & Abdunabiyev, H. (2026, February). Redefining learning efficacy with fuzzy logic algorithm: Cognitive load analysis and decision making in special education classroom. In *2026 International*

- Conference on Intelligent Processing, Hardware, Electronics, and Radio Systems (CIPHER)* (pp. 1–6). IEEE.
24. Solano, M. (2025). *Pedagogical insights: A phenomenological study of non-native English speaking teachers' second language identity, experiences, and pedagogy* (Doctoral dissertation, Southern New Hampshire University).
 25. Teh, C. S., & Kudus, A. R. (2024). The application of fuzzy cognitive mapping in education: Trend and potential. *TEM Journal*, 13(2), 976.
 26. Waluyo, G. H., Sulistiyo, U., Abrar, M., & Wulandari, B. A. (2025). Debate pedagogy in semi-military law enforcement classrooms: A case study of English-speaking fluency. *Ta'dib*, 28(2), 297–312.
 27. Wang, Q., Fang, Z., & Ye, X. (2025). Construction of a multi-objective optimization labor education course model based on fuzzy cognitive orientation. *International Journal for Housing Science & Its Applications*, 46(4), 4736.
 28. Zhou, B., Pek, L. S., Kabir, N., Yang, J., Bouteraa, M., & He, X. (2026). Artificial intelligence in English-speaking education: A WoS-based bibliometric analysis of technology, pedagogy, and ethics (2021–2025). *IJLTER.ORG*, 25(3), 853–873.
 29. Zhu, Q., Jamaludin, K. A., & Li, J. (2025). Development of flipped classroom module FCM for music theory instruction: An innovative approach to music education. *PLOS ONE*, 20(11), e0337590.

CLIMATE CHANGE AND ENVIRONMENTAL SUSTAINABILITY

D. Ashok Kumar and D. Mahhamitra*

Department of Civil Engineering,

Jai Shriram Engineering College, Tamil Nadu 638660

*Corresponding author E-mail: mahhamitra24@gmail.com

Abstract

Climate change and systemic ecosystem degradation represent the defining, interconnected existential challenges of the twenty-first century. As atmospheric carbon dioxide (CO₂) concentrations surge past 420 parts per million (ppm) and global mean temperatures track at 1.44°C above pre-industrial baselines, the limitations of traditional, reactive "end-of-pipe" environmental engineering frameworks have become critically apparent. Linear models of pollution containment and retrospective remediation are structurally unsuited to match the compounding velocity of modern ecological decay. This paper presents a comprehensive critical analysis of the paradigm shift sweeping across science and technology research, specifically investigating the convergence of Artificial Intelligence (AI), Quantum Engineering, and Synthetic Biology, and evaluates their structural impact on global environmental sustainability. By fundamentally re-engineering the scientific method itself, these emerging paradigms shift the sustainability frontier away from passive damage mitigation toward predictive, autonomous, and molecular-scale interventions. We examine how deep learning architectures accelerate global climate downscaling and optimize macroeconomic energy distribution; how quantum chemical modeling bypasses classical computing boundaries to discover novel carbon-capture materials; and how synthetic biology platforms apply CRISPR-Cas9 genetic tools to construct active, carbon-negative bio-manufactured loops. This study synthesizes current technological trajectories and models their cross-disciplinary impacts through an integrated Systems Engineering and Lifecycle Assessment (LCA) methodology. Ultimately, this paper outlines how this digital-molecular convergence provides scalable pathways to compress century-long material discovery timelines into days, offering the technological leverage required to maintain planetary boundaries below the critical 1.5°C warming threshold. Finally, we address the critical socio-technical trade-offs, immense computational infrastructure footprints, and complex regulatory and biosecurity bottlenecks that must be resolved to ensure an equitable, globally just deployment of these next-generation environmental defenses.

Keywords: Artificial Intelligence (AI), Climate Change Mitigation, Environmental Sustainability, Green Environmental Engineering, Quantum Engineering, Synthetic Biology, Systems Engineering.

1. Introduction and Historical Context

The macroscale stability of the Earth's biosphere is currently failing under the cumulative weight of industrial externalized impacts. For over two centuries, human civilization has relied on a linear economic engine—characterized by extraction, transformation, consumption, and unmitigated disposal. The physical manifestations of this paradigm are no longer localized or easily segmentable; they have combined into a global atmospheric crisis. With atmospheric carbon dioxide (CO₂) concentrations firmly fixed above 420 ppm and baseline ambient warming tracking at 1.44°C, humanity has exited the stable parameters of the Holocene and entered the high-entropy unpredictability of the Anthropocene. The contemporary global climate crisis is characterized by unprecedented shifts in atmospheric composition, with carbon dioxide (CO₂) concentrations exceeding 420 parts per million (ppm) and global mean temperatures tracking at 1.44°C above pre-industrial baselines. This anthropogenic warming amplifies positive climate feedbacks—such as cryosphere melting, albedo loss, and permafrost thawing—which directly undermine the foundational pillars of **environmental sustainability**: ensuring that current resource use does not compromise the survival needs of future generations.

Historically, environmental engineering has operated on a linear, reactive model: pollutants were generated, quantified, and subsequently filtered, contained, or remediated. However, the scale, speed, and compounding nature of modern ecological degradation render these legacy frameworks inadequate. To achieve true systemic resilience, a fundamental paradigm shift is required. Science and technology research is transitioning away from isolated, single-discipline investigations toward highly convergent, compute-intensive, and molecularly precise methodologies. Understanding how these emerging frameworks can be systematically deployed to overhaul energy, water, urban infrastructure, and waste systems is essential to building an actionable roadmap for planetary survival.

Historically, the discipline of environmental engineering emerged as a branch of civil and sanitary engineering, specifically tasked with safeguarding public health from localized industrial offcasts. This legacy operational framework was inherently reactive, giving rise to what is technically termed "end-of-pipe" remediation. Under this model, chemical pollutants, municipal sewage, or toxic particulate effluents were treated only after their generation and prior to environmental discharge. Common interventions relied heavily on brute-force mechanical and chemical isolation, such as chemical precipitation, gravity sedimentation, electrostatic precipitation, and localized physical encapsulation within multi-layered secure landfills.

While these legacy systems were highly successful in addressing acute, point-source pollution during the mid-to-late twentieth century, they are structurally unsuited to address the compounding, non-linear planetary crises of today. Modern ecological degradation is distributed, multi-variable, and self-amplifying. Brute-force filters and local containment shields cannot scale

to offset ocean-wide acidification, continent-scale biodiversity loss, or the systemic failure of international water cycles. Consequently, the research frontier must undergo a radical epistemological and practical realignment. Science and tech research must shift from physical, localized mitigation to compute-intensive, multi-scale, and predictive interventions that capture, manipulate, and optimize resource flows at the atomic and macro-infrastructure levels simultaneously.

2. Accelerating Environmental Changes and Ecosystem Stressors

To understand the urgency of this transition, the precise acceleration profile of modern environmental disruptions must be scrutinized. The saturation of planetary carbon sinks has led to an exponential thermal accumulation within the troposphere and oceans. This thermal energy acts as a systemic destabilizer across the global climate system. According to the Clausius-Clapeyron relation, the water-holding capacity of the atmosphere increases by approximately 7% for every 1°C of atmospheric warming. The physical consequence is a profound polarization of the hydrological cycle: regions prone to aridity experience accelerated soil desiccation and prolonged multi-year droughts, while humid zones are subjected to catastrophic, hyper-localized flash flooding and unprecedented precipitation anomalies.

Simultaneously, the marine biosphere is experiencing an unprecedented dual crisis of thermal stress and ocean acidification. The oceans have absorbed over 90% of the excess heat trapped by anthropogenic greenhouse gases, leading to widespread marine heatwaves that trigger rapid, catastrophic coral bleaching and ecosystem collapse. Furthermore, approximately 30% of anthropogenically emitted CO₂ is absorbed directly into surface waters, driving a chemical shift that has reduced ocean pH by 0.1 units since the pre-industrial era—representing a 30% increase in acidity. This acidification severely compromises the calcification kinetics of marine organisms, threatening the foundational links of global marine food chains.

On land, these atmospheric and marine stressors interact with existing systemic disruptions, including aggressive deforestation, intensive monoculture agriculture, and rapid urban sprawl. The result is an alarming acceleration in biodiversity loss and structural alterations to global biomes. Forests, which historically served as net carbon sinks, are converting into net carbon sources due to widespread wildfire propagation and drought-induced diebacks. The compounding speed of these ecological feedback loops threatens to push key planetary subsystems past irreversible tipping points, necessitating immediate, next-generation technological intervention.

3. Sectoral Impact of Emerging Tech Paradigms

The integration of deep-tech architectures into environmental engineering fundamentally redefines how sustainability is actioned across industrial sectors:

3.1 Artificial Intelligence & Environmental Informatics

Artificial Intelligence has evolved past simple statistical analysis to become the primary engine of predictive macro-environmental control. Deep learning models allow for the downscaling of global climate projections into hyper-localized simulations. In the energy sector, reinforcement learning algorithms handle the integration of highly volatile renewable sources into smart national grids, dynamically matching industrial loads with real-time solar and wind variations. Within municipal infrastructure, AI-driven operations allow wastewater treatment plants to process complex, shifting influent streams autonomously, optimizing biological aeration and chemical dosing schedules to reduce municipal energy overheads by up to 30%. In circular waste economics, advanced computer vision networks automate high-speed mechanical sorting of municipal solid waste, segregating complex, multi-layered polymers and valuable e-waste with superhuman precision.

3.2 Quantum Engineering & Subatomic Materials Modeling

Quantum computing circumvents the physical limits of classical supercomputers by utilizing superposition and entanglement to model molecular interactions at an atomic scale. In the domain of carbon capture, quantum algorithms simulate electron configurations within complex Metal-Organic Frameworks (MOFs), accelerating the discovery of materials that bond with and extract CO₂ from ambient air with minimal thermal energy penalties. For the green hydrogen economy, quantum chemistry accelerates the synthesis of non-precious metal catalysts for high-efficiency water splitting, eliminating reliance on expensive, scarce platinum-group elements. Furthermore, quantum-precision simulations guide the development of solid-state lithium-metal batteries, optimizing electrolyte-electrode interfaces to double the range of electric vehicles while eliminating thermal runaway risks.

3.3 Synthetic Biology & Bio-Digital Convergence

Synthetic biology shifts environmental remediation from passive reliance on natural processes to the engineering of targeted biological mechanisms. By applying CRISPR-Cas9 genetic tools, synthetic biologists alter the metabolic pathways of marine cyanobacteria, enabling them to directly convert captured atmospheric CO₂ into high-purity, biodegradable bioplastics. In contaminated land zones, transgenic plant-microbe symbioses are engineered to hyper-accumulate toxic heavy metals—such as lead, cadmium, and arsenic—from industrial brownfields, isolating the toxins within the plant tissue for easy harvesting and mineral recovery. For public health, paper-based cell-free gene networks act as highly sensitive biosensors, changing color or producing an electrical signal within minutes when exposed to trace endocrine disruptors or pharmaceutical residues in remote rural water networks.

3.4 Space Technology & Hyperspectral Remote Sensing

The deployment of high-density small satellite constellations equipped with advanced hyperspectral imaging sensors has transformed global climate accountability. Operating with on-satellite edge-AI processing, these orbital networks observe greenhouse gas emissions with unparalleled spatial and chemical resolution. Instead of relying on retrospective, self-reported emission estimates from national industries, these systems perform real-time, automated fingerprinting of point-source methane leaks from deep-sea oil rigs, agricultural zones, and aging distribution pipelines. This continuous data stream feeds directly into international compliance frameworks, transforming climate policy from voluntary commitments into an objective, legally binding discipline verified by real-time orbital telemetry.

4. Sustainable Design Methodologies and Built Environment Integration

Transitioning from theoretical deep-tech discovery to practical environmental stability requires translating these breakthroughs into actionable engineering design methodologies. In urban planning, the historical approach of channeling stormwater through rigid concrete culverts is being replaced by Sponge City hydro-informatics. This methodology uses predictive fluid dynamics and sensor-driven green architecture to design interconnected matrices of permeable pavements, subterranean retention aquifers, bioswales, and vegetative green roofs. These urban systems naturally absorb, filter, and store severe storm runoff, simultaneously mitigating urban flash floods, recharging local water tables, and reducing the urban heat island effect through natural evapotranspiration.

In chemical and industrial manufacturing, green engineering design relies on the principles of process intensification. Traditional batch processing plants, which generate immense thermal waste and toxic solvent byproducts, are replaced by continuous-flow microfluidic reactors. These systems optimize mass and heat transfer at a microscopic scale, minimizing the energy required for chemical synthesis and enabling the widespread utilization of bio-based, non-volatile solvents like supercritical CO₂ or ionic liquids. By integrating closed-loop zero-liquid discharge (ZLD) systems into these industrial designs, factories can recover and reuse virtually 100% of their chemical process water, extracting valuable dissolved mineral salts from the remaining solid waste for reuse in secondary manufacturing pipelines.

On a broader ecological scale, sustainable engineering design leverages advanced Digital Twins—highly detailed, real-time virtual replicas of vulnerable regional ecosystems. By feeding continuous data from IoT sensor arrays and satellite telemetry into a centralized computational twin, engineers can run predictive simulations to model how an ecosystem will respond to proposed interventions before execution. Whether optimizing the microbiome composition of a constructed wetland to treat agricultural runoff or adjusting urban coastal barrier designs against rising sea levels, Digital Twins allow for the rigorous testing and validation of green engineering systems under varied climate scenarios.



5. Cross-Paradigm Technical Analysis Matrix

The strategic deployment of these advanced technology paradigms requires balancing their computational and laboratory resource demands against their anticipated impacts and regulatory hurdles:

Paradigm Cluster	Primary Technical Bottleneck	Anticipated Sustainability Impact	Regulatory / Ethical Hurdle
Artificial Intelligence & Informatics	High computational energy overhead; sensor degradation in harsh industrial environments.	Reduces municipal energy demands by up to 30%; enables autonomous resource balancing.	Establishing legal liability frameworks for autonomous pollution discharge errors.
Quantum Molecular Engineering	Hardware limitations of current intermediate-scale quantum processors.	Compresses century-long material science research timelines into days or weeks.	Mitigating the technological divide between nations with advanced quantum assets.
Synthetic Biology & Convergence	Ensuring strict biocontainment within open-loop, dynamic ecosystems.	Enables targeted, molecular-scale remediation of legacy industrial contaminants.	Navigating biosecurity and ecological safety regulations for altered organisms.
Geospatial Systems & Remote Sensing	Processing massive datasets at the edge; satellite orbital maintenance costs.	Provides global, unfalsifiable point-source tracking of emissions and leaks.	Balancing international sovereignty concerns against global climate transparency.

Conclusion and Future Strategic Horizons

Mitigating climate change and achieving lasting environmental sustainability can no longer rely on incremental adjustments to legacy industrial processes. The emerging paradigms of AI-driven informatics, quantum molecular engineering, synthetic biology, and advanced earth observation systems offer the technological leverage necessary to compress century-long discovery timelines into months or days. These deep-tech frameworks change the nature of environmental action, enabling humanity to move past reactive cleanup and build an inherently predictive, molecularly precise, and circular industrial infrastructure.

However, these advanced technological tools do not represent a simple solution. As documented in the analysis, their deployment introduces unique socio-technical bottlenecks. The immense energy footprints required to train next-generation AI models and operate quantum infrastructure present a potential paradox: if the energy to run these advanced systems is derived from fossil fuels, the technological solution itself will exacerbate atmospheric warming. Furthermore, the

intentional deployment of engineered biological organisms into open, natural ecosystems requires a level of regulatory oversight and biosecurity tracking that is currently unmapped in international law.

Ultimately, technology functions as an accelerator, not a surrogate for systemic socio-economic change. To fully realize the impact of these scientific breakthroughs, global governance must evolve concurrently. This involves establishing global data commons to guarantee equitable access to computational infrastructure, preventing an international "technological divide" between the Global North and Global South. By merging advanced deep-tech innovation with a steadfast commitment to climate justice, environmental equity, and circular economic design, humanity can engineer a resilient civilization capable of enduring, adapting to, and ultimately reversing the anthropogenic degradation of the planet.

References

1. Crutzen, P. J. (2002). Geology of mankind. *Nature*, 415(6867), 23.
2. Intergovernmental Panel on Climate Change (IPCC). (2023). *Climate change 2023: Synthesis report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. IPCC, Geneva, Switzerland.
3. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., ... & Hassabis, D. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, 596(7873), 583–589.
4. MacArthur, E., & Waughray, D. (2016). *Intelligent assets: Unlocking the circular economy potential of the Internet of Things*. World Economic Forum & Ellen MacArthur Foundation.
5. National Academies of Sciences, Engineering, and Medicine. (2019). *A research agenda for transforming separation science*. The National Academies Press.
6. Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F. S., Lambin, E. F., ... & Foley, J. A. (2009). A safe operating space for humanity. *Nature*, 461(7263), 472–475.
7. Steffen, W., Richardson, K., Rockström, J., Cornell, S. E., Fetzer, I., Bennett, E. M., ... & Sorlin, S. (2015). Planetary boundaries: Guiding human development on a changing planet. *Science*, 347(6223), 1259855.
8. Venter, J. C., Remington, K., Heidelberg, J. F., Halpern, A. L., Rusch, D., Eisen, J. A., ... & Smith, H. O. (2004). Environmental genome shotgun sequencing of the Sargasso Sea. *Science*, 304(5667), 66–74.

BIOREMEDIATION: MICROBIAL SOLUTIONS FOR ENVIRONMENTAL SUSTAINABILITY

Arulmozhi R.* and Sudharani P.

PG and Research Department of Microbiology,
Sengamala Thayaar Educational Trust Women's College (Autonomous)
Sundarakkottai, Mannargudi, Thiruvarur (dt) - 614 016, Tamil Nadu.

*Corresponding author E-mail: amozhi72@gmail.com

Abstract

Environmental pollution resulting from industrialization, urbanization, agricultural activities, mining, and improper waste disposal has emerged as a significant global environmental concern. Conventional pollution-control methods may be costly, energy-intensive, and, in some cases, capable of generating secondary pollutants. Bioremediation is an environmentally friendly approach that employs living organisms, particularly microorganisms, to remove, degrade, transform, or detoxify pollutants present in soil, water, and other environmental systems. Microorganisms, including bacteria, fungi, algae, and certain archaea, possess diverse metabolic capabilities that enable them to utilize organic pollutants as sources of nutrients or transform toxic substances into less harmful forms. Microbial bioremediation has been applied to the treatment of petroleum hydrocarbons, pesticides, dyes, heavy metals, industrial wastes, and other environmental contaminants. Techniques such as bioaugmentation, biostimulation, biosorption, phytoremediation-assisted bioremediation, and microbial degradation provide sustainable approaches for environmental pollution management. This chapter presents the principles of microbial bioremediation, the microorganisms involved, mechanisms of pollutant removal, major bioremediation techniques, applications, advantages, limitations, and future perspectives. The integration of microbiology, biotechnology, and environmental science can improve the effectiveness of bioremediation and make an important contribution to environmental sustainability.

Keywords: Bioremediation, Microorganisms, Biodegradation, Bioaugmentation, Biostimulation, Environmental Sustainability, Pollution Control.

Introduction

Environmental pollution is a major challenge that affects ecosystems, human activities, and biodiversity. Rapid industrialization, intensive agricultural practices, excessive chemical usage, petroleum contamination, plastic waste, mining activities, and the discharge of untreated wastewater contribute significantly to the accumulation of harmful substances in the environment. Pollutants can be broadly classified as organic and inorganic substances. Organic pollutants include petroleum hydrocarbons, pesticides, dyes, phenols, and solvents, whereas

inorganic contaminants include heavy metals such as lead, mercury, cadmium, chromium, and arsenic. Many of these pollutants can persist in the environment and may eventually enter food chains.

Bioremediation offers a natural and sustainable method for managing environmental pollution. It involves the use of microorganisms or their biological activities to decrease the concentration or toxicity of contaminants. Microorganisms are particularly valuable in bioremediation because of their metabolic diversity and their capacity to survive and function under different environmental conditions. The basic principle of microbial bioremediation can be represented as:

Pollutant → Microbial action → Transformation/Degradation → Less harmful products

Thus, bioremediation establishes a direct connection between microbial activity, environmental protection, and sustainable development.

Principle of Microbial Bioremediation

Microorganisms obtain energy and nutrients through a variety of biochemical reactions. During these metabolic processes, certain microorganisms are capable of utilizing pollutants as sources of carbon or energy. For instance, some bacteria can degrade petroleum hydrocarbons into simpler compounds, which under appropriate aerobic conditions may ultimately be converted into carbon dioxide, water, and microbial biomass. The general process can be represented as:

Contaminated environment → Pollutant identification → Selection of suitable microorganisms → Microbial degradation/transformation → Reduction of pollutant toxicity → Environmentally safer products

The effectiveness of bioremediation is influenced by several factors, including temperature, pH, oxygen availability, moisture, nutrient availability, pollutant concentration, and the size and activity of the microbial population.

Microorganisms Involved in Bioremediation

A broad range of microorganisms contributes to the removal, degradation, transformation, or immobilization of environmental pollutants.

Bacteria

Bacteria are among the most widely utilized microorganisms in bioremediation. Their rapid growth and diverse metabolic pathways enable them to degrade a variety of environmental contaminants. Important bacterial genera include *Pseudomonas*, *Bacillus*, *Rhodococcus*, *Acinetobacter*, *Alcaligenes* and *Sphingomonas*.

These bacteria are capable of degrading hydrocarbons, pesticides, dyes, and several industrial pollutants.

Fungi

Fungi produce extracellular enzymes that are capable of degrading complex organic molecules. Important fungal genera involved in bioremediation include *Phanerochaete*, *Pleurotus*, *Aspergillus*, *Trametes* and *Penicillium*

White-rot fungi are particularly significant because they produce extracellular enzymes such as laccase, lignin peroxidase, and manganese peroxidase, which can degrade complex organic compounds.

Algae

Algae can contribute to the removal of nutrients, metals, and certain organic pollutants from wastewater. In addition, algal activity contributes to oxygen production and carbon fixation.

Archaea

Certain archaea are capable of surviving under extreme environmental conditions and can participate in the transformation of pollutants. They are particularly relevant in anaerobic environments where conventional aerobic microorganisms may be less effective.

Mechanisms of Microbial Bioremediation

Microorganisms remove or transform environmental pollutants through several biological mechanisms.

Biodegradation

Biodegradation refers to the microbial breakdown of complex organic compounds into simpler substances.

A general representation is:



When an organic compound is completely degraded into inorganic end products, the process is referred to as mineralization.

Biosorption

Biosorption involves the binding of pollutants, particularly heavy metals, to the surfaces of microbial cells. Functional groups present in microbial cell-wall components, including carboxyl, hydroxyl, phosphate, and amino groups, can interact with and bind metal ions.

Bioaccumulation

Bioaccumulation occurs when microorganisms absorb or take up pollutants into their cells. This mechanism can contribute to the removal of certain contaminants from the surrounding environment.

Biotransformation

During biotransformation, microorganisms convert a toxic compound into a different chemical form. The resulting compound may be less toxic, more stable, or easier to remove from the environment.

Mineralization

Mineralization refers to the complete microbial conversion of organic compounds into simple inorganic substances, such as carbon dioxide, water, and mineral nutrients.

Major Microbial Bioremediation Techniques

Biostimulation

Biostimulation involves modifying or improving environmental conditions to enhance the growth and metabolic activity of naturally occurring microorganisms. Nutrients such as nitrogen and phosphorus, oxygen, moisture, or appropriate electron acceptors may be supplied to improve microbial activity.

Example: Addition of nutrients to petroleum-contaminated soil can stimulate indigenous hydrocarbon-degrading bacteria.

Bioaugmentation

Bioaugmentation involves introducing selected microorganisms into a contaminated environment to enhance pollutant degradation. The introduced microorganisms may possess specific metabolic capabilities that are insufficiently represented within the native microbial population.

Bioventing

Bioventing involves supplying air or oxygen to contaminated soil in order to stimulate the activity of aerobic microorganisms. It is mainly applied for the treatment of volatile and semi-volatile petroleum hydrocarbons.

Biosparging

Biosparging involves the injection of air into groundwater systems to increase oxygen availability and stimulate microbial degradation of contaminants.

Landfarming

In landfarming, contaminated soil is spread over a prepared treatment area and periodically mixed. This promotes aeration and microbial activity, thereby supporting the degradation of pollutants.

Composting

Composting involves mixing organic wastes with suitable materials and allowing microbial decomposition to occur under controlled conditions. This process can reduce certain organic contaminants while simultaneously generating a useful soil amendment.

Applications of Microbial Bioremediation

Petroleum Hydrocarbon Pollution

Oil spills and petroleum leakage can contaminate both terrestrial and aquatic ecosystems. Hydrocarbon-degrading bacteria and fungi are capable of breaking down various petroleum components. Bacterial genera such as *Pseudomonas*, *Rhodococcus*, and *Bacillus* have been investigated for their ability to degrade hydrocarbons.

Heavy Metal Contamination

Heavy metals cannot be biologically destroyed because they are chemical elements. However, microorganisms can alter their chemical forms, immobilize them, precipitate them, or adsorb them onto their cell surfaces. Important heavy-metal contaminants include:

- Lead (Pb)
- Mercury (Hg)
- Cadmium (Cd)
- Chromium (Cr)
- Arsenic (As)

Pesticide Contamination

Microorganisms can degrade certain pesticides and transform them into less harmful compounds. This capability is particularly useful for agricultural soils that have been affected by excessive or prolonged pesticide application.

Textile Dye Wastewater

Textile industries generate wastewater containing synthetic dyes that can be difficult to remove using conventional methods. Certain bacteria and fungi can degrade or decolorize dyes through enzymatic activities. Fungal enzymes, particularly laccases and peroxidases, have considerable importance in dye treatment.

Domestic and Industrial Wastewater

Microorganisms have a central role in wastewater treatment. Biological treatment systems employ microbial communities to remove organic matter and nutrients from wastewater. Activated sludge is one of the major examples of microbial wastewater treatment.

Plastic Pollution

Certain microorganisms have demonstrated the ability to interact with, modify, or degrade particular polymers and plastic-associated compounds. Microbial cells and microbial enzymes are therefore being investigated as potential approaches for managing plastic waste.

Important Microbial Enzymes in Bioremediation

Microbial enzymes play a major role in the degradation and transformation of environmental pollutants.

Table 1: Microbial enzymes and its role

Enzyme	Major Role
Laccase	Oxidation of phenolic compounds and dyes
Lignin peroxidase	Degradation of complex aromatic compounds
Manganese peroxidase	Oxidation of organic pollutants
Dehalogenases	Transformation of halogenated compounds
Oxygenases	Oxidation of hydrocarbons and aromatic compounds
Hydrolases	Breakdown of various organic compounds

These enzymes may function within microbial cells or may be secreted outside the cells as extracellular enzymes.

Factors Affecting Bioremediation

The efficiency of microbial bioremediation is influenced by a combination of environmental and biological factors.

Temperature

Temperature strongly affects microbial growth and enzyme activity. Most environmental microorganisms show optimal activity within a specific temperature range.

pH

Extremely acidic or alkaline conditions can inhibit microbial growth and enzymatic activity. Maintaining a suitable pH can therefore improve the efficiency of pollutant degradation.

Oxygen Availability

Aerobic microorganisms require oxygen for many degradation reactions, whereas anaerobic microorganisms can perform their metabolic activities in the absence of molecular oxygen.

Nutrient Availability

Nutrients such as nitrogen, phosphorus, sulfur, and other essential elements are required to support microbial growth and metabolic activity.

Moisture

Adequate moisture is essential for microbial metabolism and facilitates the movement and availability of nutrients and pollutants within the treatment environment.

Pollutant Concentration

Very high concentrations of pollutants may inhibit microbial activity because of their toxic effects. Therefore, pollutant concentration is an important factor in determining treatment efficiency.

Microbial Population

The presence, abundance, and metabolic activity of appropriate pollutant-degrading microorganisms are essential for successful bioremediation.

Advantages of Microbial Bioremediation

Microbial bioremediation offers several important advantages:

- It is an environmentally friendly approach.
- It is generally less expensive than many physical and chemical treatment methods.
- It can be carried out either in situ or ex situ.
- It reduces dependence on harsh chemicals.
- It can transform certain pollutants into less harmful products.
- It supports sustainable waste management.
- It can be applied to large contaminated areas.
- It utilizes naturally occurring biological processes.
- In many applications, it produces fewer secondary pollutants.

Limitations of Bioremediation

Although bioremediation provides several environmental benefits, it also has certain limitations:

- Microbial activity is strongly influenced by environmental conditions.
- Some pollutants are resistant to microbial degradation.
- Very high pollutant concentrations may inhibit microbial growth and activity.
- Bioremediation may proceed more slowly than certain physical or chemical treatment methods.
- Complete degradation of pollutants cannot always be achieved.
- Introduced microorganisms may fail to survive or compete effectively with indigenous microorganisms.
- Continuous monitoring is necessary to assess treatment efficiency.
- Heavy metals cannot be destroyed biologically and therefore require processes such as transformation, adsorption, precipitation, or immobilization rather than degradation.

***In Situ* and *Ex Situ* Bioremediation**

***In Situ* Bioremediation**

In situ bioremediation involves treating the contaminated material directly at the site without removing it from the contaminated environment. Examples include:

- Bioventing
- Biosparging
- *In-situ* biostimulation
- *In-situ* bioaugmentation

***Ex Situ* Bioremediation**

Ex situ bioremediation involves removing contaminated soil or water from the original location and treating it at another site or within a controlled treatment system.

Examples include:

- Landfarming
- Biopiles
- Composting
- Bioreactors

Table 2: Comparison of *In Situ* and *Ex Situ* Bioremediation

Feature	<i>In Situ</i>	<i>Ex Situ</i>
Treatment location	At the contaminated site	Outside the contaminated site
Excavation	Usually not required	Usually required
Cost	Often lower	Often higher
Control	Relatively less	Greater control
Application	Large contaminated areas	Controlled treatment systems

Role of Bioremediation in Environmental Sustainability

Environmental sustainability refers to meeting present environmental requirements without compromising the ability of future generations to meet their own needs. Microbial bioremediation supports this principle by utilizing natural biological processes for pollution control. The overall contribution can be represented as:

Pollution reduction → Ecosystem protection → Resource recovery → Reduced chemical use → Sustainable environment

Microbial treatment can also contribute to the circular economy by converting certain organic wastes into useful products, including compost, biogas, and microbial biomass.

Recent and Future Perspectives

The future development of microbial bioremediation is closely associated with advances in biotechnology, molecular biology, and related technologies.

Metagenomics

Metagenomic approaches enable researchers to investigate microbial communities directly from environmental samples. These techniques can help identify microorganisms and genes associated with pollutant degradation.

Genomics and Proteomics

Genomic and proteomic approaches can be used to identify degradation pathways, enzymes, and proteins involved in microbial bioremediation. These approaches provide deeper insights into the molecular basis of pollutant transformation.

Microbial Consortia

Rather than relying on a single microorganism, researchers increasingly investigate microbial consortia containing multiple microorganisms that can work cooperatively to degrade complex pollutants.

Immobilized Microorganisms

Microbial cells can be immobilized on appropriate support materials to improve their stability, enhance their application in treatment systems, and facilitate their reuse.

Enzyme-Based Bioremediation

Purified or engineered microbial enzymes can provide targeted approaches for the degradation or transformation of specific environmental pollutants.

Synthetic Biology

Synthetic biology offers the possibility of developing microorganisms with specifically designed metabolic pathways for targeted environmental applications. However, such approaches require appropriate environmental risk assessment, biosafety measures, and effective containment strategies.

Conclusion

Microbial bioremediation is an important biological strategy for managing environmental pollution and promoting environmental sustainability. Bacteria, fungi, algae, and archaea possess diverse metabolic capabilities that enable them to degrade, transform, immobilize, or remove a wide range of environmental contaminants. Techniques such as biostimulation, bioaugmentation, bioventing, biosparging, landfarming, and composting have important applications in the treatment of contaminated soil and water. However, the effectiveness of bioremediation can be limited by environmental conditions, pollutant toxicity, and slow degradation rates. Advances in molecular biology, metagenomics, microbial biotechnology, enzyme technology, and synthetic biology provide new opportunities for improving the efficiency and applicability of bioremediation. Integrating microbial biotechnology with sustainable environmental management can therefore contribute to safer, more effective, and environmentally responsible approaches to pollution control.

References

1. Vidali, M. (2001). Bioremediation: An overview. *Pure and Applied Chemistry*, 73(7), 1163–1172.
2. Boopathy, R. (2000). Factors limiting bioremediation technologies. *Bioresource Technology*, 74(1), 63–67.
3. Alexander, M. (1999). *Biodegradation and bioremediation*. Academic Press.
4. Gadd, G. M. (2004). Microbial influence on metal mobility and application for bioremediation. *Geoderma*, 122, 109–119.
5. Lovley, D. R. (2003). Cleaning up with genomics: Applying molecular biology to bioremediation. *Nature Reviews Microbiology*, 1, 35–44.
6. Das, N., & Chandran, P. (2011). Microbial degradation of petroleum hydrocarbon contaminants: An overview. *Biotechnology Research International*, 2011, 1–13.
7. Verma, J. P., Jaiswal, D. K., & others. (n.d.). Microbial approaches for sustainable environmental remediation.
8. Maier, R. M., Pepper, I. L., & Gerba, C. P. (n.d.). *Environmental microbiology*. Academic Press.

SHIFTING PARADIGMS IN SCIENCE AND TECHNOLOGY: FROM DATA-DRIVEN DISCOVERY TO CONVERGENT SOLUTIONS

Umarani

Department of Chemistry (Science and Humanities),

Hindusthan Institute of Technology, Coimbatore-641032, Tamil nadu, India

Corresponding author E-mail: umarani.kalidas@gmail.com

Abstract

The paradigmatic change in science and technology research takes place due to the appearance of various technologies, multidisciplinary approaches, digital tools, and paradigms built on sustainability which alter the way of knowledge production, validation, distribution, and implementation. Besides the traditional research conducted in a specific field, there exist data-based, computational, collaborative, and convergent approaches. Artificial Intelligence, machine learning, big data, Internet of things, quantum technologies, biotechnology, nanotechnology, advanced materials, robotics, digital twins, and open science have become important elements of contemporary research environments. However, besides the increase in scientific discoveries' pace, the impact of these innovations is related to changing the relations between researchers, technologies, industries, government, and society. Technology convergence is especially valuable for new research paradigms because of its ability to link knowledge and skills of various fields [1].

In this chapter, a number of major paradigms that are becoming apparent within the realm of science and technology studies are introduced and assessed for their consequences with regard to science, technology, economy, environment, education, and society. Apart from many other paradigms, the emphasis will be on the AI-assisted discovery, data-driven research, intelligent sensing, quantum science, synthetic biology, nanotechnology, digital twins, autonomy, interdisciplinary research, and open science. On top of that, the importance of sustainability, responsible innovation, research integrity, data management, cybersecurity, privacy, and fair access to emerging technologies gains increasing significance. The convergence of scientific disciplines and technologies of the digital world can possibly be able to solve many of the problems humanity is facing in terms of climate change, environmental pollution, health care, food supply, energy, and sustainable development. But for this to be done successfully, it needs to be accompanied by ethics and human intervention.

Keywords: Emerging Paradigms, Science and Technology Research, Artificial Intelligence, Big Data, Internet of Things, Quantum Technology, Biotechnology, Nanotechnology, Digital Twins, Open Science, Sustainability, Responsible Innovation.

1. Introduction

Over time, science and technology have developed in response to various changes in observation techniques, experimental methods, computations, communications, and information dissemination. The development of scientific research has moved from observation-based research to experimental science, theory-based modelling, computation science, and research using increasing amounts of data. Nowadays, the research landscape is shaped by the process of rapid technological convergence when artificial intelligence, advanced computations, biotechnology, materials science, sensors, robotics, and digital platforms combine to open up new areas of science [1].

Today's research landscape differs significantly from the traditional disciplinary research. Scientists now work with large and heterogeneous datasets produced by sensors, satellites, labs, electronic documents, simulations, social media, and other devices. Ability to gather and analyze these datasets creates new chances for revealing correlations that could not be revealed using traditional experimental methods [2]. Machine learning is able to reveal patterns in complex datasets, computational models are able to simulate physical and biological processes, and digital platforms make international cooperation possible.

The next key change is the shift towards interdisciplinary and transdisciplinary research. Today many of the challenges that we face cannot be effectively tackled by the methods of one particular scientific discipline. Challenges like climate change, sustainable water management, public health, food security, energy transitions, biodiversity conservation, and urban planning contain an interconnected array of environmental, technological, economic, and social components [3]. Modern research therefore involves not just scientific knowledge but also engineering, computing, social sciences, policy studies, and other perspectives.

Today technology convergence is a key research paradigm. According to the OECD, technology convergence between fields such as artificial intelligence, synthetic biology, neurotechnology, quantum technologies, and earth observation is a key trend in modern science and technology [1]. Technology convergence may result in the emergence of innovative technologies, disciplines, and innovations.

However, technology development poses certain challenges as well. AI-based systems could replicate any biases that were found in training datasets; automation of scientific processes could produce false certainty; and growing reliance on digital systems makes scientists concerned about cybersecurity, privacy, reproducibility, and even energy consumption [4]. Hence, development of science responsibly is possible only when technological skills are combined with ethics, transparency, research integrity, and social responsibility.

In this chapter, we will look at the major paradigms that are emerging within science and technology research and their impact on scientific discoveries and innovations.

2. Evolution of Research Paradigms

The evolution of research paradigms may be considered as a shift from observation towards increasing integration of experiments, computation and data-based research.

2.1 Experimental and Theoretical Research

Traditionally, scientific research is based on observations, experiments, formulation of hypotheses, measurements and theoretical explanations. Experimentation is essential in scientific research since the empirical validation is a way to verify the existence of scientific knowledge. Theoretical research is based on the use of mathematics and other theoretical models to explain the observation.

2.2 Computational Research

The advent of supercomputers created a new stage in scientific research. Computational modelling allows scientists to study the objects that are expensive, dangerous or just impossible to investigate using experiments. Examples of computational science include weather forecast, molecular simulations, computational fluid dynamics, structural modelling and astrophysics.

2.3 Data-Driven Research

With the rapid development of information technologies, there appeared a new scientific environment where the amount of both structured and unstructured data is huge. Data-driven science is based on observation, experiments, computation, statistics and machine learning [2]. Nowadays, the modern research instruments produce data constantly, which requires automated ways of storing, processing, analysing and visualisation.

2.4 Convergence Research

While the prevailing paradigm is evolving from merely merging experimentation and computation to convergence within disciplines and technologies, convergence involves integrating diverse scientific disciplines, technologies, methodologies, and viewpoints [1]. Convergence is especially applicable in new and rapidly developing domains like biological applications of artificial intelligence, quantum sensing, smart materials, precision medicine, and digital twins of the environment.

3. Major Emerging Paradigms in Science and Technology Research

3.1 Artificial Intelligence and Machine Learning

Artificial intelligence is one of the major emerging paradigms in science research. Machine learning allows computer systems to detect patterns, make predictions, classify data, optimize processes, and aid in making decisions based on the available information. AI has become an integral part of scientific methodology, including the analysis of scientific literature, image interpretation, molecular design, materials discovery, modeling, and experimentation [5].

In chemistry and materials science, machine learning algorithms may be applied to predict molecular properties, discover new materials, optimize chemical reactions, and analyze spectral

data. In healthcare research, artificial intelligence is used to perform medical imaging, disease predictions, develop drugs, and find personalized treatment options [6]. In environmental science, machine learning may be applied to satellite images, air-quality data, climate data, and ecological data.

AI generates another dimension through the aid in scientific writing, programming, interpretation of data, hypothesis creation, and research planning. Nonetheless, researchers need to differentiate the role of assistance from that of scientific validation since information generated using AI may not always be accurate, based on false claims, or even falsified [4].

Consequently, the effect of AI in research becomes twofold. While AI may help increase efficiency and speed up the research process, an overreliance on AI may result in an “illusion of understanding,” where researchers blindly believe in the outputs of computations while failing to understand the assumptions behind them [4].

3.2 Big Data and Data-Driven Science

Use of big data is becoming an integral part of contemporary science. Data sets used in scientific applications can be described through characteristics such as size, speed, diversity and quality. Proper handling of scientific data in the era of big data includes data management, metadata, computing power, statistics and reproducibility.

One area where we can see an excellent example is the Earth-system sciences. Combination of satellite data, environmental sensors, physical models and machine learning algorithms allows researchers to develop more advanced environmental prediction models. Digital Earth that combines big data with physics and computing was recently suggested [7].

It is possible for research that uses big data to find out connections and patterns which may be hard to discern with the traditional method. Nevertheless, correlation alone cannot prove causation. The interpretation of science, therefore, needs to be done using not only statistical information but also domain knowledge and reasoning, among other things.

The ability to use data effectively will be essential in future scientific work. Abilities in handling data, programming, statistics, visualization, and artificial intelligence have become useful skills in scientific domains.

3.3 Internet of Things and Intelligent Sensing

IoT connects sensors, devices, machines, communications and software platforms. IoT-based studies allow for constant monitoring of physical systems as well as data acquisition in real-time. Some applications of IoT are smart agriculture, environment monitoring, healthcare, industrial automation, energy, transport and smart cities. For instance, in agriculture, there could be sensors measuring soil moisture, temperature, humidity, nutrient content and crop conditions. The acquired data could then be sent to analysis platforms.

The integration of IoT with artificial intelligence and edge computing is becoming more prevalent. In lieu of all data transmission to the central cloud, edge devices can perform computations nearer the location of data gathering. This may lead to reductions in latency, communication, and dependence on the centralized architecture.

There are, however, many issues that need to be overcome in future IoT research such as interoperability, cybersecurity, dependability of the sensors, power usage, data sovereignty, and privacy [8].

3.4 Quantum Science and Technology

Quantum science is yet another emerging research paradigm that could have relevance in computing, communications, sensing, chemistry, and materials science. The technologies enabled by quantum science are based on such phenomena as superposition, entanglement, and coherence to enable the execution of certain processes that might otherwise prove challenging using conventional technology.

Quantum computers are under development to be used in areas like molecular modeling, optimization, cryptography, and machine learning.

Quantum technology developments remain to be characterized by many difficult technical problems, such as quantum error correction, scalability issues, stabilization issues, complexity of hardware, and necessary specialized infrastructure [9]. However, quantum science demonstrates increasing convergence of physics, mathematics, computer science, engineering, chemistry, and materials science.

3.5 Biotechnology and Synthetic Biology

Modern biotechnology is becoming more and more focused on designing biological parts rather than observing biological phenomena. In synthetic biology, engineering concepts like modularity, standardization, design, modeling, and optimization are applied to biological processes [16,17].

Some areas of application of synthetic biology include medicine, agriculture, food technology, environmental cleanup, and industrial biotechnology [10]. Designing biological circuits and cells can be used for diagnostics, delivering medicines, biosensors, and chemical production.

One very promising trend in synthetic biology is its convergence with AI and automation technologies. AI technologies can help analyze and predict biological sequences, whereas automated laboratories can speed up experiments.

However, there are certain issues associated with synthetic biology such as biosafety, biosecurity, ecological impacts, intellectual property, and ethical aspects. Therefore, there should be governance mechanisms developed as well.

3.6 Nanotechnology and Advanced Materials

Nanotechnology deals with materials and structures of the nano scale range in which size-dependent properties give unique optical, electrical, mechanical, catalytic, and biological properties. Nanomaterials are used in energy storage, sensing, catalysis, environmental remediation, medicinal purposes, coating, and electronics.

Advanced materials science is becoming more computational and data-oriented. The researchers can use simulation, high-throughput experiments, machine learning, and characterization techniques for speeding up the discovery process of materials. Nanotechnology is also very useful for sustainable purposes. The nanostructured catalysts can enhance the reaction efficiencies, whereas the advanced membranes are helpful in the purification of water. Moreover, nanomaterials can help in improving the batteries and energy-storage systems.

On the other hand, the environmental and health effects of engineered nanomaterials need to be assessed properly. Life cycle analysis, toxicology studies, safe by design philosophy, and proper disposal are very essential for the research projects.

3.7 Digital Twins and Simulation-Based Research

The digital twin refers to the computational replication of a physical system using observation, models, and feedback. Digital twins are being developed for applications in engineering, manufacturing, healthcare, city design, Earth-system sciences, and disaster response [18].

For environmental research, digital twins can combine observation, models, machine learning, and simulations in order to facilitate monitoring and predictions [7]. Similar methods can be applied for modeling equipment, buildings, transport systems, and energy systems.

Digital twins are a big leap forward from modeling to research based on dynamic and interactive analysis. The success of this process requires high-quality data, effective models, computing capabilities, interconnectivity, and updating.

3.8 Robotics and Autonomous Systems

Robotics is transitioning from programmed robots to autonomous and adaptive systems. Robots can be equipped with AI, computer vision, sensors, machine learning, and sophisticated control systems in today's world. Some of the scientific uses of robots include laboratory automation, exploration of hazardous environments, agricultural robotics, underwater study, robotics in medicine, and space exploration. Laboratory automation allows for the completion of repetitive experiments while scientists concentrate on planning the experiment and analysis of its results.

However, autonomous systems can bring new issues related to responsibility, human control, employment, and reliability, thus the scientific research should cover human-robot interaction.

4. Interdisciplinary and Transdisciplinary Research

Due to the increasing complexity of modern science issues, cooperation between different disciplines becomes essential. Interdisciplinary research uses ideas and approaches from several

disciplines, while transdisciplinary research involves cooperation not only among scholars but also practitioners, policymakers, companies, communities, and other stakeholders [3,19].

The sustainability science is a case in point. Environmental problems are both scientific, technological, economic, cultural, and political. Transdisciplinary approaches may help to integrate scientific knowledge and experiences of stakeholders and local communities in solving these problems [3].

To conduct successful interdisciplinary research, cooperation between researchers from different disciplines is not sufficient. Common goals, shared language, compatible approaches, data sharing, and approaches to dealing with differences in assumptions need to be developed.

Table 1: Emerging Research Paradigms and Their Potential Impacts

Research Paradigm	Key Technologies / Methods	Major Applications	Potential Impact
AI-driven research	Machine learning, generative AI	Discovery, prediction, automation	Faster analysis and knowledge generation
Big-data science	Cloud computing, analytics	Climate, health, engineering	Evidence-based decision-making
IoT research	Sensors, edge computing	Agriculture, cities, environment	Real-time monitoring
Quantum technology	Quantum computing, sensing	Simulation, security, metrology	New computational capabilities
Synthetic biology	Gene circuits, genome engineering	Medicine, agriculture, industry	Biological innovation
Nanotechnology	Nanomaterials, nanostructures	Energy, sensors, water treatment	Advanced functional materials
Digital twins	Simulation, sensors, AI	Industry, Earth systems, cities	Predictive and adaptive management
Robotics	Autonomous systems, computer vision	Laboratories, agriculture, healthcare	Automation and improved safety
Open science	Open data, open access, repositories	All disciplines	Greater accessibility and reproducibility

5. Digitalization of Scientific Research

The digital transformation touches upon practically all phases of the research process. The digital methods can be used for literature search, experiment planning, data collection, data analysis, modeling, visualization, publication, and collaboration.

Cloud computing allows geographically distributed research groups to use computational resources and shared databases. Digital storage helps preserve data over the long run, and electronic lab notebooks are useful for keeping records.

The development of scientific software is yet another important trend. Research work can use different programming languages, statistical software, simulators, machine learning methods, databases, and visualization tools. As a result, reproducibility requires not only dissemination of research papers but also the description of computational infrastructure, datasets, algorithms, and software code.

Finally, digital technology can facilitate automation of experiments. Laboratory robots can conduct repetitive measurements, and intelligent algorithms can select conditions of further experiments. In this way, automated experimentation can boost the pace of research work.

6. Open Science and Collaborative Research

Open science is a significant shift in the process of producing and disseminating scientific knowledge. It encourages free access to publications, data, software, infrastructure, and other research products, and also stimulates broader participation in the process of scientific research [11].

The Recommendation on Open Science by UNESCO focuses on the aspects of accessibility, inclusiveness, transparency, cooperation, and reduction of knowledge and technological gaps [6]. Open science may increase the reproducibility of results since researchers can review datasets, methods, software, and other evidence.

Open research enables cooperation across organizations and countries through common database and repository that allows researchers to use existing knowledge and avoid unnecessary replication of results.

However, there is always the need to find the right balance between openness and privacy, intellectual property, ethical issues, security of nation and responsible management of data. Some research data cannot be made openly available. Data related to the health care of individuals, industrial information, and some security-related data should not be disclosed publicly.

The future of open science will lie in this compromise.

7. Science and Technology for Sustainable Development

The concept of sustainability is now becoming an important aim of research in science and technology. New technologies can help sustainable development through increased resource efficiency, waste reduction, renewable energy systems, environmental monitoring, and well-informed decisions making.

Artificial intelligence and IoT technologies can help to develop intelligent systems of energy management. Energy consumption sensors will allow predicting the needs of consumers and

optimizing its distribution. Precision technologies will help to decrease water and fertilizer use in agriculture.

New advanced materials can be helpful in developing systems of energy storage, solar technologies, water purification, and pollution treatment. Biotechnology will assist in creating sustainable production systems and biological analogs to petroleum technologies.

Digital twins and Earth observation technologies can be used in environmental monitoring and adapting to climate change. The integration of environmental data and physical models with AI opens new possibilities of predictive environment management [7].

Table 2: Technology Trends and Sustainability Contributions

Technology	Sustainability Contribution
Artificial intelligence	Resource optimization and predictive management
IoT	Real-time environmental and energy monitoring
Big data	Evidence-based sustainability planning
Nanotechnology	Water treatment, catalysis and pollution control
Biotechnology	Sustainable production and bioremediation
Digital twins	Simulation and optimization of complex systems
Robotics	Precision agriculture and hazardous-waste handling
Quantum technologies	Advanced sensing and future computational applications

Technology, on its own, does not always constitute sustainable development. The operations of data centers and artificial intelligence require a lot of energy and computing power. E-waste is another environmental problem caused by electronic gadgets. For this reason, the analysis of sustainability should look at the entire life cycle of technology.

8. Ethical, Social and Environmental Challenges

New paradigms not only present plenty of opportunities, but also pose new types of challenges.

8.1 Data Privacy

Connected devices and data science lead to an increase in the amount of personal and behavioral information that is collected. Personal health, location, biometric, and consumer data must be secured.

8.2 Algorithmic Bias

AI algorithms learn from historical datasets. In case there are any biases in those datasets, the model may amplify them. Proper research should include dataset evaluation, bias tests, methodology, and human intervention [20,22].

8.3 Reproducibility and Research Integrity

Automated computational processes allow for faster research but may make errors harder to notice. Reproducible research entails proper dataset documentation, algorithm, version of software, experiments performed, and analysis [12].

8.4 Cybersecurity

Increased connectivity of scientific infrastructure poses new cyber security risks. Laboratories, hospitals, industries, and IoT systems need to have secure communication, authentication, access control, and data security.

8.5 Technology Disparity

Disparities in infrastructure could result in technological disparities among different organizations and geographical locations. It is not considered to be a transformative research paradigm where there is an uneven distribution of technological capabilities among the users of the research.

8.6 Environmental Impact

Developing technologies use energy and raw materials. There could be significant environmental impacts for AI computing, data centers, semiconductors, and electronics production. Hence, green computing, energy-efficient algorithms, renewable-powered infrastructures, recycling and life cycle assessment must be included in technology research [23].

8.7 Responsible Innovation

There must be responsible innovation where researchers assess the possible impact of new technologies even before its large-scale deployment. Issues related to ethics, laws and society must be incorporated in the research process and not after technology creation [20,21].

9. Effect On Education, Industry, and Society

Changes in research approaches have had an effect on education as well. Future graduates will need to be equipped with skills in multidisciplinary domains of knowledge along with data analysis, programming, communication, critical thinking, and ethics.

Engineering and sciences can benefit from project-based learning, where students solve real-life problems based on different technologies. A sustainability project can include such disciplines as chemistry, internet of things sensors, data analytics, artificial intelligence, and environmental sciences.

Changes also affect the industry, where research ecosystems that include universities, startups, government organizations, technological firms, and communities prevail. Innovations today happen not within organizations but via networks.

Effect on society can also be quite high – with the help of AI, healthcare can be improved, precision agriculture can contribute to food production, environmental sensors can detect pollution early, and new materials can be used for clean energy.

However, social acceptance is very important. Efficient technologies can fail to work if communities are not willing to use them. Communication, engagement, transparency, and innovation must be integrated into the process of technological development [13].

10. Research Gaps and Future Directions

While new technologies offer great promise for research, there still exists a number of gaps.

First of all, the link between AI and fundamental science should be strengthened. Models of AI should integrate more physical laws, chemistry, biology, and domain-specific knowledge beyond the statistics [24].

Secondly, new organizational forms should be developed for interdisciplinary research. University departments and funding organizations usually operate within disciplinary framework. Thus, new criteria for assessing collaborative works, datasets, software and societal impact should be developed.

Thirdly, the research infrastructure should be interoperable. All instruments, databases, software and archives should cooperate to create integrated research process.

Fourthly, responsible innovations require greater implementation. Ethics assessment should be incorporated into the process of emerging technology research, especially in the areas of AI, biotechnology, neurotechnology, quantum technology and autonomous systems.

Fifthly, sustainability should be introduced as an explicit criterion for the technological development. Energy costs, resource requirements, ecological impact, reparability and recyclability should be assessed.

Table 3: Future Research Priorities

Research Priority	Required Development
AI-enabled science	Explainable, reliable and domain-aware AI
Interdisciplinary research	Shared frameworks and collaborative infrastructure
Open science	Reproducible data, software and research workflows
Sustainable technology	Life-cycle assessment and green design
Digital twins	Real-time data-model integration
Quantum research	Scalable and reliable quantum systems
Biotechnology	Safe and responsible biological engineering
IoT	Secure, interoperable and energy-efficient networks
Research ethics	Integrated ethical and societal assessment
Global research	Equitable access to knowledge and infrastructure

Future research endeavors will increasingly shift towards autonomous and quasi-autonomous scientific systems in which the AI will detect patterns, form hypotheses, design experiments, interpret outcomes, and formulate further experiments. This type of research could speed up discoveries to an incredible degree, but scientists have to ensure human control over the process.

The other important trend is the creation of “convergence spaces” where people from various fields can collaborate in the same physical space using similar computational means, databases,

and research goals [1]. Convergence spaces are able to give rise to entirely new sciences and technologies.

The future of research thus does not seem to be dependent on a single technology. It will rather consist of the interaction between many technologies, disciplines, organizations, and social actors.

Conclusion

There is a new era coming for science and technology research which will feature the emergence of such traits as convergence, digitalization, automation, interdisciplinarity and data-driven discoveries. Artificial intelligence, big data, IoT, quantum technologies, biotechnologies, nanotechnologies, robotics, digital twins and open science are transforming research methodologies and changing the scope of scientific studies.

The most significant contribution of all these new paradigms is not the technologies themselves but the possibility of their combination. AI and biotechnology will facilitate the discovery of biology-related facts and phenomena; IoT and data analysis – intelligent environmental monitoring; advanced materials and nanotechnologies – sustainable technologies; and digital twins along with Earth observation technologies – environmental prediction and management.

In turn, scientific advancement should be accompanied by responsible governance as privacy issues, cybersecurity problems, research ethics, algorithmic bias, environmental problems, inequality of access to research and others cannot be overlooked. Science should become more and more related to transparency, inclusivity, sustainability and social responsibility.

Open science and interdisciplinary collaboration may contribute to increasing involvement in the knowledge-making process, and convergence may lead to new scientific and technological opportunities. Hence, future studies must target the creation of technologies that are innovative, but at the same time, explainable, safe, eco-friendly, socially inclusive, and human-oriented.

The emerging research landscape shows that science and technology are not evolving along independent disciplinary paths. The future will be determined by research ecosystems interconnected with people, information, computation, experimentation platforms, technologies, and societal knowledge. The key task of researchers thus lies in taking advantage of convergence and translating scientific and technological breakthroughs into social benefits.

References

1. OECD. (2025). *OECD science, technology and innovation outlook 2025: Driving change in a shifting landscape*. OECD Publishing. <https://doi.org/10.1787/5fe57b90-en>
2. Kitchin, R. (2014). Big data, new epistemologies and paradigm shifts. *Big Data & Society*, 1(1), 1–12. <https://doi.org/10.1177/2053951714528481>
3. Lang, D. J., Wiek, A., Bergmann, M., Stauffacher, M., Martens, P., Moll, P., Swilling, M., & Thomas, C. J. (2012). Transdisciplinary research in sustainability science: Practice,

- principles, and challenges. *Sustainability Science*, 7, 25–43. <https://doi.org/10.1007/s11625-011-0149-x>
4. Messeri, L., & Crockett, M. J. (2024). Artificial intelligence and illusions of understanding in scientific research. *Nature*, 627, 49–58. <https://doi.org/10.1038/s41586-024-07146-0>
5. Jumper, J., Evans, R., Pritzel, A., et al. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, 596, 583–589. <https://doi.org/10.1038/s41586-021-03819-2>
6. Topol, E. J. (2019). High-performance medicine: The convergence of human and artificial intelligence. *Nature Medicine*, 25, 44–56. <https://doi.org/10.1038/s41591-018-0300-7>
7. Li, X., Feng, M., Ran, Y., et al. (2023). Big data in Earth system science and progress towards a digital twin. *Nature Reviews Earth & Environment*, 4, 319–332. <https://doi.org/10.1038/s43017-023-00409-w>
8. Hajjaji, Y., Boulila, W., Farah, I. R., Romdhani, I., & Hussain, A. (2021). Big data and IoT-based applications in smart environments: A systematic review. *Computer Science Review*, 39, 100318. <https://doi.org/10.1016/j.cosrev.2020.100318>
9. Preskill, J. (2018). Quantum computing in the NISQ era and beyond. *Quantum*, 2, 79. <https://doi.org/10.22331/q-2018-08-06-79>
10. Zhao, N., Song, Y., Xie, X., et al. (2023). Synthetic biology-inspired cell engineering in diagnosis, treatment and drug development. *Signal Transduction and Targeted Therapy*, 8, 112. <https://doi.org/10.1038/s41392-023-01375-x>
11. UNESCO. (2021). *UNESCO recommendation on open science*. UNESCO.
12. National Academies of Sciences, Engineering, and Medicine. (2019). *Reproducibility and replicability in science*. National Academies Press. <https://doi.org/10.17226/25303>
13. Bornmann, L. (2013). What is societal impact of research and how can it be assessed? A literature survey. *Journal of the American Society for Information Science and Technology*, 64(2), 217–233. <https://doi.org/10.1002/asi.22803>
14. Wyborn, C., Datta, A., Montana, J., Ryan, M., Leith, P., Chaffin, B., Miller, C. A., & van Kerkhoff, L. (2019). Co-producing sustainability: Reordering the governance of science, policy, and practice. *Annual Review of Environment and Resources*, 44, 319–346. <https://doi.org/10.1146/annurev-environ-101718-033103>
15. Wiek, A., Withycombe, L., & Redman, C. L. (2011). Key competencies in sustainability: A reference framework for academic program development. *Sustainability Science*, 6, 203–218. <https://doi.org/10.1007/s11625-011-0132-6>
16. Yan, X., Liu, X., Zhao, C., & Chen, G. Q. (2023). Applications of synthetic biology in medical and pharmaceutical fields. *Signal Transduction and Targeted Therapy*, 8, 199.
17. Davies, J., & Levin, M. (2023). Synthetic morphology with agential materials. *Nature Reviews Bioengineering*, 1, 46–59. <https://doi.org/10.1038/s44222-022-00001-9>

18. Dal Zilio, L., Giardini, D., Carbonell, R., et al. (2023). Harnessing the potential of digital twins in seismology. *Nature Reviews Earth & Environment*, 4, 510–512. <https://doi.org/10.1038/s43017-023-00469-y>
19. National Research Council. (2014). *Convergence: Facilitating transdisciplinary integration of life sciences, physical sciences, engineering, and beyond*. National Academies Press.
20. European Commission. (2019). *Ethics guidelines for trustworthy AI*. High-Level Expert Group on Artificial Intelligence, European Commission.
21. UNESCO. (2021). *Recommendation on the ethics of artificial intelligence*. UNESCO.
22. National Institute of Standards and Technology. (2023). *Artificial intelligence risk management framework (AI RMF 1.0)*. NIST. <https://doi.org/10.6028/NIST.AI.100-1>
23. World Economic Forum. (2025). *The global risks report 2025*. World Economic Forum.
24. OECD. (2023). *Artificial intelligence in science: Challenges, opportunities and the future of research*. OECD Publishing.
25. UNESCO. (2023). *Open science outlook 1: Status and trends around the world*. UNESCO.

WSe₂ THIN FILMS: STRUCTURAL AND OPTICAL PROPERTIES

Prashant A. Chate^{1*} and Dattatray J. Sathe²

¹Department of Chemistry, J.S.M. College, Alibag (M.S.) India

²Department of Chemistry, KIT College of Engg (Autonomous), Kolhapur (M.S.) India.

Corresponding author E-mail: pachate04@rediffmail.com, djsathe77@gmail.com

1. Introduction

Two-dimensional (2D) materials have emerged as an important class of functional materials because their physical properties can differ substantially from those of their bulk counterparts. The discovery and rapid development of graphene stimulated extensive research into layered materials with properties that cannot be readily obtained from conventional three-dimensional semiconductors. Among these materials, transition-metal dichalcogenides (TMDs), represented by compounds such as MoS₂, WS₂, MoSe₂, and WSe₂, have received particular attention [1-2].

Tungsten diselenide, WSe₂, is a semiconducting member of the layered transition-metal dichalcogenide family. It is composed of tungsten atoms coordinated by selenium atoms in a Se–W–Se sandwich configuration. The individual layers are strongly bonded internally but are weakly coupled to adjacent layers through van der Waals interactions. This structural arrangement permits mechanical exfoliation and enables thin-film formation down to the monolayer regime. WSe₂ is particularly interesting because its electronic structure changes with thickness. Bulk and multilayer WSe₂ generally possess an indirect band gap, while monolayer WSe₂ exhibits a direct band gap. This thickness-dependent transition strongly influences optical absorption, photoluminescence, carrier dynamics, and device performance.

In addition to its electronic properties, WSe₂ offers advantages associated with its relatively strong environmental stability, chemical tunability, and compatibility with van der Waals heterostructures. The presence of tungsten and selenium also produces significant spin–orbit coupling, which is important for spintronic and valleytronic applications.

The properties of WSe₂ thin films are strongly dependent on synthesis conditions. Parameters such as substrate type, deposition temperature, precursor concentration, Se/W ratio, pH, complexing agent, growth time, annealing temperature, and atmosphere can influence crystallinity, grain size, stoichiometry, thickness, defect concentration, and phase composition. Consequently, systematic studies of WSe₂ thin films are essential for understanding the relationship between synthesis, structure, and properties and for optimizing the material for specific applications [3-4].

2. Crystal Structure and Properties of WSe₂

WSe₂ belongs to the transition-metal dichalcogenide family with the general chemical formula MX₂, where M represents a transition metal and X represents a chalcogen. In WSe₂: M = W; X = Se.

A single WSe₂ layer consists of a plane of tungsten atoms sandwiched between two selenium planes. Each tungsten atom is coordinated by six selenium atoms in a trigonal-prismatic arrangement in the stable semiconducting 2H phase. The layers are held together by weak van der Waals forces. This weak interlayer interaction is fundamentally different from the strong covalent bonding within an individual layer. The most commonly encountered polytypes include:

2.1 2H-WSe₂

The 2H phase is the thermodynamically stable semiconducting phase under ordinary conditions. It is characterized by trigonal-prismatic coordination of tungsten. This phase is particularly important for field-effect transistors, photodetectors, photovoltaic devices, optoelectronic devices, valleytronics.

2.2 1T-WSe₂

The 1T phase possesses octahedral coordination and can display substantially different electronic behavior. Depending on structural conditions and defects, distorted or metastable phases may also occur. Phase engineering can therefore be used to modify the electrical and catalytic properties of WSe₂.

The layered structure is one of the most important characteristics of WSe₂. Within each layer, W–Se bonds are strong, whereas neighboring layers interact primarily through van der Waals forces. As a result, the material can be separated into atomically thin sheets. This structure provides several advantages like atomic-scale thickness control, large surface-to-volume ratio, reduced dielectric screening, strong interaction with incident light, efficient surface functionalization, compatibility with van der Waals heterostructures, and potential flexibility in ultrathin-film form. The weak interlayer bonding also permits the fabrication of few-layer structures without necessarily introducing dangling bonds at the surfaces.

The electronic properties of WSe₂ are strongly thickness dependent. Bulk WSe₂ generally behaves as an indirect-gap semiconductor. When this material is reduced to a monolayer, the locations of the valence-band maximum and conduction-band is minimum change, resulting in a direct electronic transition. This transition is extremely important for optoelectronic applications. The electronic structure is also influenced by strong spin–orbit coupling arising from the heavy tungsten atoms. Consequently, the valence-band states exhibit significant spin splitting. These characteristics make WSe₂ particularly attractive for valleytronics, spintronics, quantum optoelectronics, excitonic devices, photodetection, and nanoscale transistors.

One of the remarkable characteristics of atomically thin WSe₂ is its strong excitonic behavior. An exciton is a bound state consisting of an electron and a hole attracted through Coulomb interaction. In conventional bulk semiconductors, dielectric screening reduces electron–hole interaction. In atomically thin materials, however, dielectric screening is substantially reduced. Consequently, WSe₂ can exhibit strongly bound excitons. The optical response of WSe₂ is therefore dominated by excitonic transitions rather than simple independent electron–hole transitions. Important excitonic species include neutral excitons, charged excitons or trions, localized excitons, defect-bound excitons and interlayer excitons in heterostructures. These states provide opportunities for developing exciton-based optoelectronic and quantum devices [5-6].

3. Chemical and Solution-Based Deposition

Solution-based approaches are particularly attractive for researchers interested in inexpensive and scalable thin-film deposition. Chemical bath deposition (CBD), dip coating, successive ionic layer adsorption and reaction (SILAR), electrodeposition, and related techniques have been widely investigated for other metal chalcogenides. For WSe₂, the chemistry is more complicated because tungsten precursors generally require controlled reduction and selenium incorporation. A generalized solution-growth process may involve preparation of a tungsten precursor, stabilization or complexation of tungsten species, preparation of a selenium precursor, controlled mixing, substrate immersion, nucleation of W-containing species, selenium incorporation, growth of WSe₂ nuclei, washing, drying, and thermal treatment. The possibility of adapting low-temperature chemical methods to WSe₂ is scientifically interesting because it could enable deposition on inexpensive and temperature-sensitive substrates.

The growth of WSe₂ thin films can generally be understood in terms of nucleation followed by crystal growth. Initially, precursor species approach or form near the substrate surface. When the local concentration exceeds the critical supersaturation, stable nuclei form. The nuclei subsequently grow through surface diffusion, attachment of reactive species, lateral growth, and coalescence. The final morphology depends on the relative rates of nucleation and growth. A high nucleation rate generally produces smaller grains, greater grain density, more continuous films. A lower nucleation density with rapid crystal growth may produce larger grains, thicker crystallites, island-like morphology. Controlling this balance is critical for obtaining uniform WSe₂ thin films.

Substrate temperature influences precursor decomposition, surface diffusion, nucleation, and crystallinity. An increase in temperature can initially improve crystallinity by increasing surface mobility. However, excessive temperatures may cause selenium loss or phase instability. Increasing deposition time generally increases film thickness until precursor depletion or surface saturation occurs. Very long deposition times can produce excessive thickness, rough surfaces, secondary phases, nonuniform morphology. Precursor concentration controls supersaturation and

nucleation density. Low concentration may result in incomplete coverage, whereas excessively high concentration may produce rapid nucleation and poorly crystalline deposits. The Se/W ratio is particularly important. Selenium deficiency can generate Se vacancies, W-rich regions, altered carrier concentration, defect states. Adequate selenium chemical potential is therefore important for obtaining stoichiometric WSe₂ [7-8].

4. X-Ray Diffraction

The structural properties of WSe₂ thin films were investigated by X-ray diffraction using CuK α radiation with a wavelength of 1.5406 Å. The XRD pattern confirms the crystalline nature of the deposited WSe₂ films and can be indexed to the hexagonal 2H phase of tungsten diselenide. The characteristic diffraction peaks are generally observed at approximately $2\theta = 13.6^\circ, 31.5^\circ, 37.0^\circ, 47.5^\circ, 56.0^\circ, \text{ and } 65.5^\circ$, corresponding to the crystallographic planes of WSe₂. Among these reflections, the low-angle peak at about 13.6° is assigned to the (002) plane and is characteristic of the layered structure of WSe₂. The presence of the (002) reflection is associated with the periodic stacking of Se–W–Se layers along the crystallographic c-axis. The other reflections arise from the in-plane and mixed crystallographic arrangements of tungsten and selenium atoms. The absence of additional prominent diffraction peaks attributable to elemental tungsten, selenium, tungsten oxide, or other tungsten selenide phases indicates the formation of the desired WSe₂ phase. The interplanar spacing of the different crystallographic planes was calculated using Bragg's law, $n\lambda = 2d \sin\theta$. For the (002) reflection observed at $2\theta = 13.6^\circ$, the corresponding d-spacing is approximately 6.50 Å. The relatively large interplanar spacing is characteristic of the layered structure and weak van der Waals interaction between adjacent WSe₂ layers. Similarly, the reflection at 31.5° gives a d-spacing of 2.84 Å. For a hexagonal crystal system, the lattice parameters were calculated using the relationship $1/d^2 = 4(h^2 + hk + k^2)/(3a^2) + l^2/c^2$. The calculated lattice constants are approximately $a = 3.28 \text{ Å}$ and $c = 13.0 \text{ Å}$, which are consistent with the hexagonal 2H structure of WSe₂. The unit-cell volume calculated using $V = (\sqrt{3}/2)a^2c$ is 121.1 Å^3 . Small deviations in the lattice parameters from standard values may be associated with residual strain, selenium vacancies, nonstoichiometry, substrate-induced stress, and defects generated during film growth.

The broadening of the diffraction peaks provides information about the crystallite dimensions and structural disorder. The average crystallite size can be estimated using the Scherrer equation, $D = K\lambda/\beta\cos\theta$, where K is the crystallite shape factor, λ is the wavelength of the X-ray radiation, β is the full width at half maximum of the diffraction peak in radians, and θ is the Bragg angle. For example, an FWHM of 0.20° for the (002) reflection gives an estimated crystallite size of 40 nm. The relatively small crystallite dimensions are characteristic of nanocrystalline thin films and may result from the high nucleation density during deposition.

The XRD peak broadening may originate not only from finite crystallite size but also from lattice strain. Therefore, Williamson–Hall analysis can be employed to distinguish the contributions of crystallite size and microstrain. The Williamson–Hall equation, $\beta \cos \theta = K\lambda/D + 4\epsilon \sin \theta$, allows the crystallite size to be obtained from the intercept and the microstrain from the slope of the $\beta \cos \theta$ versus $4 \sin \theta$ plot. The presence of microstrain can be attributed to lattice imperfections, dislocations, grain boundaries, vacancies, and mismatch between the WSe₂ film and the underlying substrate. Selenium vacancies are particularly significant in WSe₂ because selenium deficiency can produce local lattice distortion and modify the electronic properties of the film. The dislocation density, which provides an approximate measure of the density of crystallographic defects, can be calculated using $\delta = 1/D^2$. For a crystallite size of 40 nm, the corresponding dislocation density is approximately $6.25 \times 10^{10} \text{ cm}^{-2}$. A relatively low dislocation density generally indicates improved crystalline quality, whereas a higher value suggests a greater concentration of structural imperfections. The preferred orientation of WSe₂ crystallites can also be evaluated using the texture coefficient. A texture coefficient greater than unity for a particular crystallographic plane indicates preferential orientation along that plane. Because WSe₂ possesses a layered structure, preferential orientation of the basal planes parallel to the substrate is commonly important and can strongly influence charge transport and surface properties. The structural characteristics obtained from XRD are closely related to the morphology and functional properties of WSe₂ thin films. Larger crystallites, improved crystallinity, reduced microstrain, and lower defect density generally favor efficient carrier transport by reducing grain-boundary and defect scattering. In contrast, excessive selenium vacancies, lattice distortion, and grain boundaries may introduce localized electronic states and act as recombination centers. Thus, the XRD analysis provides an important foundation for correlating the deposition conditions and crystal structure of WSe₂ thin films with their optical, electrical, photoelectrochemical, and device-related properties [9-10].

5. Optical Properties

The optical properties of WSe₂ thin films are strongly influenced by their layered crystal structure, film thickness, crystallinity, surface morphology, defect concentration, and number of layers. WSe₂ exhibits significant interaction with visible and near-infrared radiation, making it an attractive semiconductor for photodetectors, photovoltaic devices, optical sensors, light-emitting devices, and other optoelectronic applications. The optical characteristics of WSe₂ thin films can be evaluated from absorption, transmission, reflection, absorption coefficient, optical band gap, Urbach energy, extinction coefficient, refractive index, dielectric constant, optical conductivity, and optical penetration depth. The absorption spectrum of WSe₂ thin films is generally investigated in the wavelength range of about 300–1000 nm. A pronounced absorption edge is normally observed in the visible region, followed by strong absorption toward shorter

wavelengths. The absorption coefficient, α , can be calculated from the measured absorbance using the relation; $\alpha = 2.303A/t$, where A is the absorbance and t is the thickness of the film. The absorption coefficient represents the extent to which incident photons are absorbed within the material. WSe₂ thin films can exhibit absorption coefficients of the order of 10^4 – 10^5 cm⁻¹ in the strongly absorbing region, demonstrating their strong light-absorbing capability.

The optical transmittance of WSe₂ films generally decreases in the region of strong absorption and increases toward longer wavelengths. The magnitude of transmittance depends on film thickness, crystallinity, surface roughness, grain size, and defect concentration. Increasing the thickness of the WSe₂ film generally increases the optical path length and consequently enhances absorption, resulting in lower transmittance. Thin WSe₂ films can therefore provide a useful combination of light absorption and partial optical transparency, which is advantageous for thin-film optoelectronic and photodetection applications. The optical band gap is one of the most important parameters for understanding the electronic structure of WSe₂. It can be determined using the Tauc relation; $(\alpha h\nu)^n = B(h\nu - E_g)$ where $h\nu$ is the photon energy, B is a constant, E_g is the optical band gap, and n represents the nature of the electronic transition. For an allowed direct transition, $n = 2$, and E_g is obtained by extrapolating the linear portion of the plot of $(\alpha h\nu)^2$ against $h\nu$ to the photon-energy axis. The optical band gap of WSe₂ is highly dependent on thickness. Bulk and multilayer WSe₂ generally exhibit an indirect band gap of approximately 1.2–1.3 eV, whereas monolayer WSe₂ exhibits a direct optical transition with an energy close to 1.6–1.7 eV. This variation is associated with quantum confinement, reduced dimensionality, and changes in interlayer electronic coupling. The wavelength corresponding to the optical band gap can be estimated from $\lambda_g = 1240/E_g$, where λ_g is the absorption-edge wavelength in nm and E_g is the band gap in eV. For example, an optical band gap of 1.65 eV corresponds to an absorption edge of approximately 752 nm. This demonstrates that WSe₂ can absorb a substantial portion of the visible and near-infrared regions of the electromagnetic spectrum.

The absorption edge of WSe₂ is also influenced by structural disorder and localized electronic states. The Urbach energy is therefore an important parameter for evaluating the degree of disorder in the film. The Urbach relation is given by $\alpha = \alpha_0 \exp(h\nu/E_u)$, where α_0 is a constant and E_u is the Urbach energy. The Urbach energy can be obtained from the inverse slope of the linear portion of the $\ln \alpha$ versus $h\nu$ plot. A smaller Urbach energy corresponds to a sharper absorption edge and generally indicates better structural order and lower defect concentration. Conversely, a larger Urbach energy is associated with increased disorder, grain boundaries, vacancies, strain, and localized states near the band edges.

The extinction coefficient, k , represents the attenuation of electromagnetic radiation within the WSe₂ film and is calculated using, $k = \alpha\lambda/4\pi$, where λ is the wavelength of incident radiation. The extinction coefficient normally increases in the region of strong optical absorption and

decreases in the transparent region. A high extinction coefficient indicates strong interaction between the incident electromagnetic radiation and the electronic states of WSe₂. The refractive index is another important optical parameter that describes the propagation of electromagnetic radiation through the WSe₂ film. It is related to the dielectric properties of the material and can be determined from optical transmission and reflection measurements. The refractive index generally exhibits anomalous dispersion near the fundamental absorption edge because of strong electronic transitions. Away from the absorption region, normal dispersion is generally observed. The magnitude of the refractive index depends on film thickness, crystallinity, porosity, density, and wavelength. The complex dielectric constant of WSe₂ is expressed as $\epsilon = \epsilon_1 + i\epsilon_2$, where ϵ_1 is the real part of the dielectric constant and ϵ_2 is the imaginary part. These quantities can be obtained from the refractive index and extinction coefficient using $\epsilon_1 = n^2 - k^2$ and $\epsilon_2 = 2nk$.

The real part of the dielectric constant represents the polarization and energy-storage capability of the material, whereas the imaginary component is associated with optical absorption and energy dissipation. Strong variations in ϵ_1 and ϵ_2 near the absorption edge are related to interband electronic transitions. The optical conductivity provides information about the response of charge carriers to an electromagnetic field. It can be determined from the absorption coefficient and refractive index and generally increases in the region where photon-induced electronic transitions become significant. The optical conductivity is therefore influenced by the electronic band structure, carrier concentration, crystallinity, defects, and film thickness. The optical penetration depth, or skin depth, represents the distance over which the intensity of incident radiation decreases within the WSe₂ film. It can be expressed as $\delta = 1/\alpha$. For an absorption coefficient of 10^5 cm^{-1} , the penetration depth is approximately 10^{-5} cm , equivalent to 100 nm. Thus, a WSe₂ film with a high absorption coefficient can effectively absorb incident radiation within a relatively small thickness. This property is particularly beneficial for thin-film photodetectors, photovoltaic devices, and other light-harvesting applications. The optical density of WSe₂ films can be expressed as $OD = \alpha t$, where α is the absorption coefficient and t is the film thickness. A higher optical density corresponds to stronger attenuation of incident radiation. The optical density generally increases with increasing film thickness and absorption coefficient.

The optical response of WSe₂ is particularly sensitive to the number of layers. In bulk and multilayer WSe₂, interlayer interactions influence the electronic band structure, whereas the electronic structure changes significantly when the material approaches the monolayer limit. Monolayer WSe₂ exhibits strong excitonic effects because of reduced dielectric screening and enhanced Coulomb interactions. Consequently, pronounced excitonic absorption and photoluminescence features can occur in atomically thin WSe₂. These properties make WSe₂ highly attractive for nanoscale optical and optoelectronic devices. Defects can substantially modify the optical behavior of WSe₂ thin films. Selenium vacancies, tungsten vacancies, antisite

defects, grain boundaries, and structural imperfections can introduce localized energy states within the forbidden energy gap. These defect states may broaden the absorption edge, increase the Urbach energy, modify photoluminescence intensity, and influence electron–hole recombination. Improved crystallinity and controlled stoichiometry generally produce a sharper absorption edge and more stable optical characteristics. The optical properties of WSe₂ films can also be controlled through deposition parameters such as substrate temperature, deposition time, precursor concentration, film thickness, annealing temperature, and chemical environment. Thermal treatment can improve crystallinity and modify defect concentration, thereby changing the optical band gap, absorption coefficient, extinction coefficient, refractive index, and dielectric response. Similarly, changes in thickness can modify interlayer coupling and produce measurable changes in the optical band gap. WSe₂ thin films possess strong and tunable optical absorption, thickness-dependent electronic transitions, pronounced excitonic effects, and significant visible-to-near-infrared optical activity. The ability to control their optical properties through thickness, crystallinity, strain, defects, and layer number makes WSe₂ a promising material for photodetectors, photovoltaic devices, optical sensors, field-effect transistors, and flexible optoelectronic systems [11-14].

7. Defects and Doping in WSe₂

Defects play a major role in determining the properties of WSe₂. Common defects include selenium vacancies, tungsten vacancies, antisite defects, grain boundaries, dislocations, substitutional impurities, interstitial atoms. Among these, selenium vacancies are particularly important because selenium can be lost during high-temperature processing or under selenium-deficient conditions. Defects can influence carrier concentration, carrier mobility, optical absorption, photoluminescence, catalytic activity, and chemical stability. Defect engineering can therefore be considered both a challenge and an opportunity.

Doping provides an effective strategy for tuning WSe₂. Dopants can modify carrier concentration, Fermi level, conductivity, optical properties, catalytic activity, contact resistance. Possible dopants include transition metals, alkali metals, chalcogen substitutions, nonmetal dopants. Doping must be carefully controlled because excessive dopant incorporation can introduce defect states and degrade crystallinity.

References

1. Thakur, S., Thanigai, A., Chou, S., Dong, C., & Kandasami, A. (2025). *Optical Materials*, 162, 116772.
2. Kadiwala, K., Butanovs, E., Ogurcovs, A., Zubkins, M., & Polyakov, B. (2022). *Journal of Crystal Growth*, 593, 126764.
3. Thakur, S., Thanigai, A., Kushvaha, S., Meena, R., Dong, C., Muthusamy, S., & Kandasami, A. (2025). *Applied Physics A*, 131, 198.

4. Thomas, S., Greenacre, V., Zhang, J., Zhelev, N., Ramadan, S., Han, Y., Beanland, R., Abdelazim, N., Noori, Y., Groot, K., Reid, G., & Bartlett, P. (2024). *Journal of Materials Chemistry C*, 12, 19191.
5. Brune, V., Hartl, F., Fischer, T., Aytuna, Z., Patrun, D., Hillebrand, D., & Mathur, S. (2024). *Materials Advances*, 5, 3490.
6. Cheng, Q., Pang, J., Sun, D., Wang, J., Zhang, S., Liu, F., Chen, Y., Yang, R., Liang, N., Lu, X., Ji, Y., Wang, J., Zhang, C., Sang, Y., Liu, H., & Zhou, W. (2020). *InfoMat*, 2, 656.
7. Sathe, D., Chate, P., Hankare, P., Manikshete, A., & Sankpal, U. (2016). *Applied Nanoscience*, 6, 191.
8. Ghods, S., Esfandiar, A., Zad, A., & Vardast, S. (2022). *ACS Omega*, 7, 2091.
9. Hankare, P., Manikshete, A., Sathe, D., Chate, P., & Rathod, K. (2009). *Materials Chemistry and Physics*, 113, 183.
10. Boscher, N., Carmalt, C., & Parkin, I. (2006). *Journal of Materials Chemistry*, 16, 122.
11. Gujari, S., Hernández, A., Morales, P., Duran-Ledezma, A., Pérez, D., & Rocha, M. (2026). *Physica B*, 732, 418530.
12. Ma, Q., Kyureghian, H., Banninga, J., & Ianno, N. (2014). *MRS Online Proceedings Library*, 1670, 77.
13. Gu, H., Song, B., Fang, M., Hong, Y., Chen, X., Jiang, H., Ren, W., & Liu, S. (2019). *Nanoscale*, 11, 22762.
14. Kim, D., Lee, D., Moon, Y., Choi, J., & Kim, T. (2026). *Current Applied Physics*, 88, 111.

Emerging Paradigms in Science and Technology Research: Trends and Impact (ISBN: 978-81-69967-10-5)

About Editors



Prof. Yogita Shinde is an Assistant Professor at Vishwakarma University with 10 years of academic experience and specialized expertise in pharmaceutical sciences, particularly nutraceuticals, herbal drug delivery systems, and herbal-based pharmaceutical and nutraceutical formulations. Her research focuses on integrating the therapeutic potential of natural products with advanced drug delivery technologies to develop innovative, effective, and sustainable healthcare solutions. She has contributed significantly to scholarly research through 21 publications indexed in Scopus and Web of Science, one authored book, and five book chapters. Prof. Shinde has also guided and mentored 25 undergraduate students, promoting academic development and research excellence. Her work emphasizes translating scientific knowledge into practical applications addressing contemporary healthcare challenges. Through academic and industry collaborations, she continues to contribute to pharmaceutical sciences and healthcare innovation, with particular emphasis on sustainability, therapeutic efficacy, and patient-centred solutions.



Dr. Jagendra D. Punde, M.Sc. (Physics), SET, Ph.D., is an Associate Professor at S. S. Girls' College, Gondia, with 17 years of teaching experience at the undergraduate level. His research specialization is Solid Oxide Fuel Cells (SOFCs), reflecting his interest in advanced energy technologies and materials research. He has authored four books and published 13 research papers in national and international journals. Dr. Punde has also been granted two design patents, highlighting his contributions to innovation and applied research. He received a travel grant from the University Grants Commission (UGC), New Delhi, to present his research at the 19th International Conference on Solid State Ionics held in Japan. His academic profile combines teaching, research, scholarly writing, and innovation, with continued contributions to the fields of physics, energy technology, and solid-state materials.



Mujahid M. Hussain is an Assistant Professor in the Department of Physics at Maharashtra College of Arts, Science & Commerce, Nagpada, Mumbai. He holds an M.Sc. in Physics with specialization in Nuclear Physics and has qualified NET, SET, JEST, and GATE examinations. With nine years of teaching experience at the undergraduate level, he has developed a strong academic foundation in physics education. His research interests primarily focus on Nonlinear Dynamics and Chaos Theory, with an emphasis on understanding complex dynamical systems and their underlying behaviour. Alongside his teaching and research activities, he has contributed to academic content development and previously worked with Target Publications on solving previous-year questions for their entrance examination preparation book. His professional interests encompass physics education, research, academic writing, and the application of theoretical concepts to complex physical systems.



Dr. R. Elavarasi, M.E., Ph.D., is an Associate Professor in the Department of Electrical and Electronics Engineering at AMET Deemed to be University. She obtained her B.E. in Electrical and Electronics Engineering and M.E. from the University of Madras, Chennai, and her Ph.D. from AMET University. She secured the University Third Rank in her M.E. examinations. With 26 years of teaching and research experience across reputed institutions, she has published more than 35 research papers in SCI, Scopus-indexed, UGC, and peer-reviewed journals. She is a lifetime member of ISTE, IAENG, and SCIEI. She has received a funded project from the Tamil Nadu State Council for Science and Technology and has three patent publications to her credit. Her research interests include power converters, special machines, and renewable energy sources.

